



A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000-2010



A Profile of Kentucky Medicaid Mental Health Diagnoses, 2000-2010

BY
Michael T. Childress

RESEARCH SUPPORTED BY
The Foundation for a Healthy Kentucky

SEPTEMBER 2012

College of Communication and Information
308B Lucille Caudill Little Fine Arts Library
University of Kentucky
Lexington, KY 40506-0224

Center for Business and Economic Research
335AV Gatton College of Business and Economics
University of Kentucky
Lexington, KY 40506-0034

859.257.2912 office
859.257.7671 fax
michael.childress@uky.edu

PREFACE

Good policy is dependent upon good data. This is especially true in health policy. Here we provide data on the number of Kentucky Medicaid beneficiaries who have received a mental or behavioral health diagnosis from 2000 to 2010. *A Profile of Kentucky Medicaid Mental Health Diagnoses, 2000-2010* provides information on the total number of individuals who have been diagnosed with a mental health disorder as well as a year-by-year count of the 15 broad categories used to classify these diagnoses. Presented for children (age 18 and younger) and adults (19 and older), these data are organized at the state, regional, and county levels—which should enable leaders and citizens to compare mental health diagnoses between different communities. These comparisons should provoke important public policy and public health questions, such as what accounts for the different patterns across the state overall, between genders, and among races with respect to, for example, ADHD, developmental disorders, and substance-related disorders. Moreover, used in conjunction with a companion report on Medicaid pharmaceutical utilization, the *Kentucky Medicaid Pharmaceutical Utilization Guide, 2000-2010*, the strategic allocation of resources dedicated to improving health literacy can be advanced—among patients, health care providers, and the community at large.

This work is a collaborative effort between the Foundation for a Healthy Kentucky, the University of Kentucky College of Communication and Information, and the Center for Business and Economic Research (CBER) in the Gatton College of Business and Economics.

Foundation for a Healthy Kentucky

This research is funded by a grant from the Foundation for a Healthy Kentucky. For more information about the Foundation, please visit <http://www.healthy-ky.org>. Inquiries about the Foundation and its various initiatives should be directed to:

Susan G. Zepeda, Ph.D., President/CEO
Foundation for a Healthy Kentucky
9300 Shelbyville Road, Suite 1305
Louisville, KY 40222
Voice: (502) 326-2583
Toll Free: (877) 326-2583
E-mail: info@healthy-ky.org
Web: www.healthy-ky.org

College of Communication and Information

Research has connected poor health literacy—the ability of individuals to understand basic health information and make appropriate decisions—to poor health outcomes and increased costs for healthcare. There are many indicators that point to poor health literacy in Kentucky: our citizens frequently make poor health choices, they suffer from high levels of chronic disease and disability, and they have low levels of prose literacy. The College of Communication and Information has launched a Health Literacy Initiative to help improve the health literacy and health outcomes of our citizens. Information about the College's Health Literacy Initiative is available at <http://cis.uky.edu/hl/>. Inquiries about the College and its various initiatives should be directed to:

Dan O'Hair, Ph.D.
Dean & Professor
308 Lucille Caudill Little Fine Arts Library
College of Communications and Information Studies
University of Kentucky
Lexington, KY 40506-0224
Voice: (859) 218-0290
E-mail: ohair@uky.edu
Web: cis.uky.edu

Center for Business and Economic Research

The Center for Business and Economic Research (CBER) is the applied economic research branch of the Carol Martin Gatton College of Business and Economics at the University of Kentucky. Its purpose is to disseminate economic information and provide economic and policy analysis to assist decision makers in Kentucky's public and private sectors. In addition, CBER performs research projects for federal, state, and local government agencies, as well as for private-sector clients nationwide. The primary motivation behind CBER's research agenda is the belief that systematic and scientific inquiries into economic phenomena yield knowledge that is indispensable to the formulation of informed public policy. Inquiries about the Center and its various initiatives should be directed to:

Chris Bollinger, Ph.D.
Professor of Economics and Director CBER
Department of Economics
335A Gatton Business and Economics BLDG
University of Kentucky
Lexington, KY 40506-0034
Voice: (859) 257-7675
E-mail: crboll@uky.edu
Web: cber.uky.edu

Institute for Pharmaceutical Outcomes and Policy (IPOP)

The Institute employs over 40 professional staff, researchers and students engaged in pharmaceutical outcomes and policy projects either grant-funded or contracted with Kentucky Medicaid, UK HealthCare, the Kentucky Cabinet for Health & Family Services, the Urban Institute, the Robert Wood Johnson Foundation, UK Human Resources and the UK Center for Clinical and Translational Science. The Institute's projects include the development of decision support technology, data warehousing, business analysis tools, policy analysis and program evaluations. Inquiries about the Institute should be directed to:

Jeffery Talbert, Ph.D.
Director, Institute for Pharmaceutical Outcomes and Policy (IPOP)
College of Pharmacy
Pharmacy Practice and Science
University of Kentucky
Lexington, KY 40536-0596
Voice: (859) 323-7141
E-mail: Jeff.Talbert@uky.edu
Web: pharmacy.mc.uky.edu

Institutional Review Board Approvals

The University of Kentucky, Office of Research Integrity, Institutional Review Board, authorized this research with Exemption Certification for Protocol No. 11-0641-X2B (September 2011), as did the Kentucky Cabinet for Health and Family Services Institutional Review Board (CHFS IRB) (November 2011).

TABLE OF CONTENTS

PREFACE	iii
ACKNOWLEDGEMENTS	vii
INTRODUCTION	1
Data and Method	2
How to Use This Report	5
Organization and Presentation	5
Limitations.....	5
A Profile of Kentucky Medicaid Mental Health Diagnoses, 2000-2010: Tables.....	7
Regions.....	10
Area Development Districts	18
BRFSS Groups	48
Counties	126

ACKNOWLEDGEMENTS

This project would not have been possible without the support of the Foundation for a Healthy Kentucky. Sarah Walsh, the project program officer at the Foundation, provided important feedback and review. Darren Henderson, with the University of Kentucky's College of Pharmacy, performed an invaluable function in making this project possible—he organized and provided the data. Jeff Talbert, the director of the Institute for Pharmaceutical Policy and Outcomes in the College of Pharmacy deserves credit for providing the initial idea for this project. While many others provided input, review, and inspiration, the author accepts sole responsibility for any errors.

INTRODUCTION

This report provides data on mental health diagnoses for individuals on Medicaid in Kentucky. Medicaid is a state-federal partnership to provide health care coverage for people with lower incomes, older people, people with disabilities, and some families and children. The Medicaid program is jointly funded by states and the federal government, but the states administer Medicaid within broad federal rules and have a lot of flexibility to design their programs. The eligibility rules for Medicaid are different for each state, but most states offer coverage for adults with children at some income level. In Kentucky, the Department for Medicaid Services, within the Cabinet for Health and Family Services, administers the Medicaid program—which in FY2010 was a \$5.6 billion program. There are many types of services provided—from inpatient hospitalization to long-term care to prescription drugs for acute care.

In the wider context of Kentucky's state budget, Medicaid constitutes a significant portion of total state government spending. According to the National Association of State Budget Officers, *State Expenditure Report: Fiscal Year 2010*,¹ 21.9 percent of Kentucky state government expenditures were for Medicaid, which was second only to higher education (22.4 percent) and slightly higher than elementary and secondary education (19.4 percent). Within the Medicaid budget, mental health—also known as behavioral health—is an important category. According to a recent study in *Health Affairs*, behavioral health spending accounts for 11.5 percent of overall Medicaid spending.² Beyond the percentages and dollars, there are other, perhaps more compelling reasons, why understanding the trajectory of various mental health disorders is important.

Individuals diagnosed with mental health disorders are at a higher risk for physical illness and the cost of their medical care is much higher compared to those without a mental health disorder. A 2011 study by the United Hospital Fund found that Medicaid beneficiaries evidence rates of hypertension, heart disease, asthma, and chronic obstructive pulmonary disease (COPD) 30 to 60 percent higher than individuals on Medicaid without a mental health condition or diagnosis.³ Moreover, those who have been diagnosed with substance abuse disorders demonstrate conditions like heart disease, asthma/COPD, and HIV/AIDS at rates from 50 to 300 percent higher than individuals on Medicaid who do not have a substance abuse disorder.⁴ Finally, average Medicaid spending for mental health beneficiaries (\$28,451) was nearly double the amount for those without a mental health diagnosis (\$15,964), and only 25 percent of the spending for the mental health beneficiaries was actually for mental health treatment, including inpatient hospital care, prescription drugs, and outpatient psychiatric services.⁵

The Centers for Disease Control and Prevention (CDC) states that while mental health and mental illness are related, “they represent different psychological states.”⁶ The CDC notes that:

- **Mental health** is “a state of well-being in which the individual realizes his or her own abilities, can cope with the normal stresses of life, can work productively and fruitfully, and is able to make a contribution to his or her community.” It is estimated that only about 17% of U.S. adults are considered to be in a state of optimal mental health.
- **Mental illness** is defined as “collectively all diagnosable mental disorders” or “health conditions that are characterized by alterations in thinking, mood, or behavior (or some combination thereof) associated with distress and/or impaired functioning.” Depression is the most common type of mental illness, affecting more than 26% of the U.S. adult population.

The CDC states that “evidence has shown that mental disorders, especially depressive disorders, are strongly related to the occurrence, successful treatment, and course of many chronic diseases including diabetes, cancer,

¹ *State Expenditure Report: 2010*, National Association of State Budget Officers, 2011, available at <www.nasbo.org>.

² This is for 2005 spending. See Mark TL, Levit KR, Vandivort-Warren R, Buck JA, Coffey RM. Changes in U.S. Spending on Mental Health and Substance Abuse Treatment, 1986 2005, And Implications for Policy. *Health Affairs* 30, No 2. (2011) (abstract available at : <http://content.healthaffairs.org/content/30/2/284>).

³ Elizabeth M. Patchias and Michael Birnbaum, *Providing Care to Medicaid Beneficiaries with Behavioral Health Conditions*, Medicaid Institute at United Hospital Fund, Feb. 2011. Available online at <http://www.uhfny.org/assets/880>.


⁴ Ibid.

⁵ Ibid.

⁶ Mental Health Basics, CDC Web site, available online at <http://www.cdc.gov/mentalhealth/basics.htm>.

cardiovascular disease, asthma, and obesity and many risk behaviors for chronic disease; such as, physical inactivity, smoking, excessive drinking, and insufficient sleep.”⁷

Survey results from the CDC Behavioral Risk Factor Surveillance System (BRFSS) provide some insight about the mental health status of Kentucky adults. The BRFSS is an annual state-based telephone survey of the U.S. civilian, non-institutionalized adult population. While the sample is drawn from the general population—not just the Medicaid population—the numbers provide some context for understanding the state of mental health in Kentucky and the United States. Figure 1 illustrates the survey results from the question, “*Now thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?*” At nearly 15 percent (14.8), a higher percentage of Kentucky adults are potentially at risk for frequent mental distress compared to the U.S. average (11.1).⁸



Data and Method

We present data on the overall number of individuals—both children (18 and younger) and adults (19 and older)—with at least one mental health diagnosis.⁹ There are 15 categories within the Medicaid billing data used to indicate if a beneficiary has been diagnosed with a mental health disorder. It is possible for someone to have more than one of these 15 disorders (e.g., anxiety disorders and substance-related disorders). While a patient is designed with a primary diagnosis, we do not limit inclusion to the primary diagnosis; if a Medicaid recipient has ever had one of the diagnoses in a given category, in any of their billing data, then they are included in these data. The categories as well as their subcategories, if applicable, are listed below:

1. Adjustment disorders
2. Anxiety disorders

⁷ Ibid.

⁸ The CDC uses this question and 3 others to calculate frequent mental distress (FMD=14 to 30 mentally unhealthy days in the past 30 days) which has been used as a proxy for poor mental health. Refer to the CDC Web site for more information: <http://www.cdc.gov/mentalhealth/data_stats/nspd.htm>.

⁹ A patient is included in the data set if they are coded with one of the 15 mental health disorder codes and they have at least two encounters within a year. An encounter is defined as the patient interacting with a health provider that generates a billing for Medicaid. These encounters do not have to be for mental health services. The encounters are both inpatient and outpatient and represent, for instance, dental visits, hospital stays, or regular doctor visits which are not necessarily related to a mental health disorder. Also, we do not present data if the cell size is less than 30 individuals. Typically this is an issue when reporting numbers for minorities at the county level. When this is the case, we also suppress the data for Whites when it might be possible to “back out” the number for minorities using the total number and the White number.

3. Attention deficit, conduct, and disruptive behavior disorders
 - a. Conduct disorder
 - b. Oppositional defiant disorder
 - c. Attention deficit disorder and Attention deficit hyperactivity disorder
4. Delirium, dementia, and amnesic and other cognitive disorders
5. Developmental disorders
 - a. Communication disorders
 - b. Developmental disabilities
 - c. Intellectual disabilities
 - d. Learning disorders
 - e. Motor skill disorders
6. Disorders usually diagnosed in infancy, childhood, or adolescence
 - a. Elimination disorders
 - b. Other disorders of infancy childhood or adolescence
 - c. Pervasive developmental disorders
 - d. Tic disorders
7. Impulse control disorders not elsewhere classified
8. Mood disorders
 - a. Bipolar disorders
 - b. Depressive disorders
9. Personality disorders
10. Schizophrenia and other psychotic disorders
11. Alcohol-related disorders
12. Substance-related disorders
13. Suicide and intentional self-inflicted injury
14. Screening and history of mental health and substance abuse codes
 - a. Codes related to mental health disorders
 - b. Codes related to substance-related disorders
15. Miscellaneous mental disorders
 - a. Dissociative disorders
 - b. Eating disorders
 - c. Factitious disorders
 - d. Psychogenic disorders
 - e. Sexual and gender identity disorders
 - f. Sleep disorders
 - g. Somatoform disorders
 - h. Mental disorders due to general medical conditions not elsewhere classified
 - i. Other miscellaneous mental conditions

We present these data at the state, region,¹⁰ Area Development District (Figure 2), Behavioral Risk Factor Surveillance System Region¹¹ (Figure 3), and county levels. Most individuals will be counted only one time, but if a beneficiary moves from one county to another and uses Medicaid to pay for services in each county then they will be counted in each county.

¹⁰ The West region is comprised of the counties in the three most western Area Development Districts (ADDs). The East region is comprised of the six most eastern ADDs. The Urban Triangle is comprised of the Bluegrass, KIPDA, and N. KY ADDs, and the South Central region is Barren River, Lincoln Trail, and Lake Cumberland ADDs.

¹¹ These county groups were developed by researchers at the University of Kentucky Markey Cancer Control Program and College of Public Health under the direction of the Kentucky Department for Public Health. The thirty-nine county groups were developed using a methodology which considered socioeconomic, demographic, and health-related statistics at the county level, together with geographical proximity and number of BRFSS respondents, to cluster counties based on similar characteristics. Aggregating counties in this manner provides more reliable estimates for areas with small populations. Figure 2 shows counties belonging to the same groups. Note that 14 counties had enough respondents from the BRFSS to produce stable estimates on their own: Boyd, Daviess, Fayette, Greenup, Hardin, Jefferson, Kenton, Mason, McCracken, Montgomery, Pike, Pulaski, Rowan, Warren.

To facilitate comparisons between regions as well as changes over time within a region we show the number of individuals with a mental health diagnosis as a percentage of the total number of Medicaid-eligible individuals (children or adults). For example, in 2010 there were 244,089 Medicaid eligible children in the billing data with at least one of the mental health disorders. This represents a number that is 46% of the total number of Medicaid-eligible children for that year and is a decrease from 53% in 2000. The Urban Triangle, by comparison, was lower than the state average in both 2000 (51%) and 2010 (42%).

FIGURE 2
Kentucky Area Development Districts



Source: Kentucky State Data Center (<http://ksdc.louisville.edu/1maps.htm>)

FIGURE 3
Behavioral Risk Factor Surveillance System (BRFSS) Regions



Source: University of Kentucky Markey Cancer Control Program and College of Public Health under the direction of the Kentucky Department for Public Health.

How to Use This Report

Comparing Medicaid mental health disorders between different geographic regions, such as county-to-county or county-to-state, or between gender and racial/ethnic groups within a region, provides a view into community-level health indicators that might not otherwise be readily available. It should generate discussion within a community about size differences between respective populations and whether they are changing over time. This information that can be used for the strategic allocation of health literacy resources. We provide examples below on how the data in this report might be used by state and local leaders:

- In the companion report on pharmaceutical utilization, *Kentucky Medicaid Pharmaceutical Utilization Guide, 2000-2010*, we show that important differences between counties across Kentucky in the use of medication for Attention Deficit Hyperactivity Disorder (ADHD) (Therapeutic Class H2V). Some of the drugs included in this class are Focalin, Metadate, and Concerta. Overall, 244.4 grams per 1,000 member-years were dispensed to Medicaid-eligible children in Kentucky. However, there is considerable variation from one county to the next—evidenced by 57.5 grams in Leslie County and 553.3 grams in Henderson County. Likewise, in 2010 about 45 percent of Medicaid-eligible children with a mental health disorder were classified as having ADHD, while Henderson County children evidenced a much higher rate at nearly 63 percent and Leslie County children showed a much lower rate at just under 32 percent.
- Medicaid-eligible adults categorized with mood disorders, which includes bipolar disorder and depressive disorders, as a percentage of all individuals with a mental health disorder increased in Kentucky from 58.8 percent in 2000 to 63.5 percent in 2010—representing an increase of 4.7 percentage points. Adults in Muhlenberg County increased by 10.4 percentage points—from 55.4 to 65.8 percent—but during the same time period Magoffin County adults remained at 56.2 percent, showing no change.

The examples above are illustrative of how the data in *A Profile of Kentucky Medicaid Mental Health Diagnoses, 2000-2010* can be used—and there are many other examples of large differences between regions and demographic groups across the state. Health, community, and education leaders in these counties might want to know why their county numbers are so different from the state average, or why they are changing over time.

Organization and Presentation

This report should be of interest to public health officials, health care providers, and community leaders. In the tables that follow we show the year-by-year numbers for the individuals with a mental health diagnosis. There are two pages for each geographic region—one for children and another for adults. We present total numbers as well as by gender and race. We then use the total number of individuals in the region who are eligible for Medicaid in that year as the denominator to calculate percentages. This facilitates comparisons between regions and through time within a region. Finally, we show the total number of individuals categorized as having any of the 15 mental health disorders. At the bottom of each page we show a chart that compares the percentage point difference between that region and the state average for the percentage of individuals with at least one mental health diagnosis as a percent of the total number of Medicaid-eligible individuals.

Limitations

First, all of the data are derived from fully adjudicated paid claims, not all claims submitted. Second, it is possible, indeed likely, that not all claims for 2010 are included in the data presented here—as they continue to be submitted and adjudicated at the time of this report. Consequently, the drop in numbers we see for 2010 *could* be a function of unprocessed claims instead of a *real* decrease in the number of individuals with a particular disorder.

A Profile of Kentucky Medicaid Mental Health Diagnoses, 2000-2010: Tables

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Kentucky											
CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	206,705	226,735	239,191	250,790	256,732	260,172	261,566	248,589	257,974	258,714	244,089
Female	102,037	110,517	115,151	119,408	121,064	121,500	121,454	114,260	118,159	117,448	109,932
Male	104,668	116,218	124,040	131,382	135,668	138,672	140,112	134,329	139,815	141,266	134,157
White	166,108	181,893	190,758	199,586	203,727	206,158	206,461	195,738	202,843	203,247	191,828
Minority	29,628	32,164	34,070	35,254	35,536	35,439	35,231	32,982	34,224	34,148	32,155
CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	55%	57%	57%	57%	57%	56%	52%	53%	50%	46%
Female	54%	55%	56%	56%	55%	54%	54%	50%	50%	47%	42%
Male	53%	55%	58%	59%	59%	59%	58%	55%	56%	53%	49%
White	53%	55%	57%	57%	57%	56%	55%	52%	52%	50%	45%
Minority	39%	39%	40%	40%	39%	38%	37%	34%	34%	32%	29%
CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	45,222	49,817	52,895	55,204	55,920	55,946	55,168	50,950	51,033	49,287	45,149
Alcohol-related disorders	9,656	9,802	9,622	9,334	8,897	8,268	7,551	6,411	5,880	5,150	4,166
Anxiety disorders	71,601	77,189	80,717	84,226	86,000	86,464	86,163	81,373	82,010	79,802	73,970
Attention-deficit, conduct, and disruptive behavior disorders	93,708	104,180	111,876	118,886	123,117	125,126	125,184	117,669	119,125	117,026	109,718
Delirium, dementia, and amnestic and other cognitive disorders	3,265	3,407	3,517	3,560	3,580	3,560	3,527	3,315	3,341	3,168	2,879
Developmental disorders	46,535	53,923	60,782	67,372	72,851	77,021	80,481	80,117	85,320	87,473	84,734
Disorders usually diagnosed in infancy, childhood, adolescence	16,116	18,479	20,565	22,513	24,124	25,239	26,050	25,206	26,020	25,871	24,822
Impulse control disorders, NEC	12,519	13,343	13,822	13,977	13,782	13,244	12,612	11,232	10,571	9,664	8,326
Miscellaneous disorders	31,506	33,268	34,053	34,590	34,642	34,204	33,463	31,028	31,081	29,977	27,582
Mood disorders	74,296	77,497	78,123	77,769	75,710	72,339	68,426	60,993	58,620	54,585	47,865
Personality disorders	7,758	7,925	7,910	7,776	7,586	7,093	6,548	5,768	5,331	4,768	4,026
Schizophrenia and other psychotic disorders	8,659	8,818	8,923	8,989	8,858	8,562	8,211	7,476	7,257	6,679	5,825
Screening and history of mental health and substance abuse	126,878	135,736	139,524	140,731	137,991	133,672	129,282	118,037	117,855	113,552	103,154
Substance-related disorders	24,462	24,659	23,991	23,144	21,815	20,163	18,450	15,984	15,143	13,641	11,375
Suicide and intentional self-inflicted injury	6,575	6,727	6,752	6,694	6,531	6,287	6,083	5,574	5,462	5,143	4,439

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Kentucky											
ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	225,450	243,303	256,716	267,776	276,437	282,089	287,249	272,093	293,459	298,589	288,956
Female	158,620	170,679	179,831	187,427	193,374	196,993	200,332	188,061	203,085	205,347	196,888
Male	66,830	72,624	76,885	80,349	83,063	85,096	86,917	84,032	90,374	93,242	92,068
White	190,390	205,588	217,053	226,574	234,081	239,134	243,872	230,392	248,919	252,625	242,958
Minority	23,803	25,661	26,840	27,691	28,150	28,040	27,470	25,675	28,197	29,318	28,518
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	66%	67%	69%	69%	69%	69%	65%	69%	68%	64%
Female	66%	68%	70%	71%	71%	71%	71%	66%	71%	70%	66%
Male	59%	61%	63%	64%	64%	64%	65%	62%	65%	64%	61%
White	64%	67%	68%	69%	70%	70%	70%	65%	69%	69%	65%
Minority	40%	42%	43%	43%	43%	42%	42%	39%	41%	41%	38%
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	22,771	25,143	27,260	29,165	30,746	32,209	33,577	32,722	35,594	36,924	36,266
Alcohol-related disorders	23,476	25,829	27,576	29,309	30,694	31,261	32,094	30,692	32,424	32,609	31,342
Anxiety disorders	129,110	140,625	150,323	158,878	165,425	170,157	175,095	169,572	180,105	182,533	177,450
Attention-deficit, conduct, and disruptive behavior disorders	11,888	13,527	15,238	17,337	19,499	21,542	23,979	24,723	28,588	31,350	32,177
Delirium, dementia, and amnesic and other cognitive disorders	52,230	53,717	53,780	52,575	51,249	50,245	49,039	46,002	45,488	42,947	39,316
Developmental disorders	21,490	22,712	23,806	24,785	25,542	26,212	26,999	26,818	28,068	28,898	29,184
Disorders usually diagnosed in infancy, childhood, adolescence	2,586	2,876	3,083	3,342	3,611	3,904	4,248	4,471	5,040	5,597	5,990
Impulse control disorders, NEC	4,135	4,599	5,039	5,502	5,994	6,418	6,955	7,040	7,803	8,364	8,574
Miscellaneous disorders	52,695	58,183	62,915	67,781	71,705	74,416	76,191	72,017	76,226	76,745	73,311
Mood disorders	132,572	144,884	155,086	164,218	171,210	176,045	181,025	174,152	185,850	188,637	183,432
Personality disorders	18,094	19,716	21,148	22,382	23,398	23,815	24,225	23,352	24,245	24,247	23,465
Schizophrenia and other psychotic disorders	49,939	52,829	54,402	55,092	55,335	55,466	55,564	53,359	53,781	52,443	49,549
Screening and history of mental health and substance abuse	126,386	140,398	152,691	163,182	171,696	176,718	181,553	172,642	187,822	191,672	185,627
Substance-related disorders	40,783	45,404	49,211	53,458	56,939	59,588	62,327	60,217	65,607	67,614	65,810
Suicide and intentional self-inflicted injury	8,506	9,435	10,333	11,210	12,114	12,983	14,204	14,259	15,757	16,285	16,154


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Western Kentucky

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	30,415	33,342	35,355	37,078	37,854	38,404	38,289	35,680	36,767	36,738	34,628
Female	14,905	16,247	16,944	17,573	17,700	17,852	17,715	16,355	16,738	16,583	15,486
Male	15,510	17,095	18,411	19,505	20,154	20,552	20,574	19,325	20,029	20,155	19,142
White	23,409	25,651	27,092	28,390	28,887	29,180	29,086	27,121	27,908	27,853	26,289
Minority	5,225	5,572	5,841	6,007	6,020	6,074	5,884	5,292	5,451	5,441	5,078

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	56%	58%	58%	57%	57%	56%	52%	53%	50%	46%
Female	54%	56%	57%	56%	55%	54%	53%	49%	49%	46%	42%
Male	54%	56%	59%	60%	59%	59%	59%	55%	56%	53%	49%
White	54%	55%	57%	57%	56%	55%	54%	51%	51%	48%	44%
Minority	41%	41%	42%	42%	41%	40%	40%	36%	36%	35%	32%


CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	5,222	5,711	6,005	6,172	6,166	6,158	6,086	5,542	5,480	5,274	4,876
Alcohol-related disorders	956	970	937	892	827	778	706	597	552	488	385
Anxiety disorders	7,812	8,301	8,627	8,950	8,963	8,950	8,770	7,992	7,869	7,472	6,783
Attention-deficit, conduct, and disruptive behavior disorders	15,493	17,146	18,316	19,479	20,155	20,432	20,185	18,709	18,817	18,432	17,348
Delirium, dementia, and amnestic and other cognitive disorders	442	468	473	488	503	501	488	458	469	446	404
Developmental disorders	5,927	6,995	8,042	9,030	9,842	10,561	11,159	11,064	11,772	12,143	11,776
Disorders usually diagnosed in infancy, childhood, adolescence	1,832	2,091	2,331	2,582	2,780	2,967	3,061	2,946	3,051	3,081	2,962
Impulse control disorders, NEC	2,658	2,809	2,883	2,914	2,900	2,829	2,673	2,385	2,251	2,076	1,817
Miscellaneous disorders	5,047	5,346	5,450	5,507	5,504	5,428	5,214	4,705	4,658	4,439	4,058
Mood disorders	11,408	12,024	12,287	12,327	12,113	11,695	11,097	9,850	9,522	8,920	7,912
Personality disorders	855	874	849	813	792	730	692	619	573	491	411
Schizophrenia and other psychotic disorders	1,134	1,134	1,157	1,160	1,132	1,087	1,032	931	887	804	710
Screening and history of mental health and substance abuse	17,921	19,329	20,098	20,209	19,732	19,322	18,670	16,930	16,922	16,405	14,966
Substance-related disorders	2,602	2,667	2,579	2,531	2,365	2,186	2,000	1,714	1,636	1,448	1,207
Suicide and intentional self-inflicted injury	832	860	844	823	819	756	728	628	597	549	481



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Western Kentucky											
ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	30,506	33,128	34,984	36,406	37,560	38,408	39,027	36,314	39,547	40,018	38,460
Female	22,564	24,371	25,668	26,712	27,480	27,965	28,321	26,014	28,264	28,498	27,092
Male	7,942	8,757	9,316	9,694	10,080	10,443	10,706	10,300	11,283	11,520	11,368
White	25,028	27,161	28,708	29,928	30,906	31,669	32,210	29,823	32,707	33,051	31,473
Minority	4,278	4,648	4,852	4,954	4,991	4,959	4,886	4,519	4,829	4,890	4,788
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	65%	67%	68%	68%	68%	68%	63%	68%	67%	63%
Female	64%	67%	69%	70%	70%	70%	70%	65%	70%	69%	64%
Male	56%	59%	61%	63%	63%	63%	63%	60%	64%	63%	59%
White	63%	66%	67%	68%	69%	69%	68%	63%	68%	67%	63%
Minority	46%	48%	50%	50%	50%	49%	48%	45%	47%	46%	43%
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,721	3,057	3,371	3,558	3,691	3,856	4,003	3,799	4,107	4,246	4,077
Alcohol-related disorders	2,593	2,853	3,054	3,266	3,462	3,497	3,620	3,448	3,692	3,738	3,611
Anxiety disorders	15,254	16,765	17,975	19,234	20,144	20,873	21,540	20,636	22,183	22,422	21,719
Attention-deficit, conduct, and disruptive behavior disorders	2,183	2,457	2,751	3,154	3,556	3,904	4,332	4,464	5,127	5,639	5,749
Delirium, dementia, and amnesic and other cognitive disorders	8,010	8,199	8,236	7,980	7,689	7,484	7,308	6,866	6,949	6,496	5,971
Developmental disorders	3,255	3,415	3,538	3,594	3,665	3,714	3,813	3,791	3,891	3,931	3,947
Disorders usually diagnosed in infancy, childhood, adolescence	261	289	323	355	390	425	455	486	543	594	646
Impulse control disorders, NEC	613	680	744	812	884	973	1,093	1,091	1,217	1,341	1,374
Miscellaneous disorders	6,216	6,967	7,519	8,048	8,574	8,906	9,142	8,466	9,128	9,269	8,875
Mood disorders	17,398	19,262	20,774	22,174	23,244	24,026	24,684	23,538	25,421	25,787	24,958
Personality disorders	2,009	2,216	2,391	2,482	2,616	2,673	2,710	2,664	2,834	2,877	2,735
Schizophrenia and other psychotic disorders	7,124	7,549	7,776	7,891	7,898	7,917	7,967	7,696	7,836	7,682	7,316
Screening and history of mental health and substance abuse	15,554	17,659	19,295	20,736	21,870	22,662	23,195	21,515	23,622	24,092	23,100
Substance-related disorders	4,037	4,575	5,042	5,462	5,852	6,200	6,498	6,189	6,806	7,019	6,665
Suicide and intentional self-inflicted injury	1,162	1,306	1,435	1,567	1,737	1,871	2,045	2,094	2,403	2,530	2,446



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

South Central Kentucky

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	37,486	41,517	44,010	46,265	47,324	48,033	48,300	46,058	47,793	48,177	45,376
Female	18,461	20,303	21,278	22,166	22,500	22,629	22,648	21,397	22,229	22,189	20,728
Male	19,025	21,214	22,732	24,099	24,824	25,404	25,652	24,661	25,564	25,988	24,648
White	32,595	36,024	38,052	39,887	40,636	41,185	41,216	39,157	40,555	40,860	38,457
Minority	3,129	3,423	3,589	3,730	3,727	3,725	3,694	3,490	3,662	3,676	3,509

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	55%	56%	57%	57%	56%	56%	53%	53%	50%	46%
Female	53%	55%	56%	56%	55%	55%	54%	50%	50%	47%	43%
Male	52%	55%	57%	58%	58%	58%	58%	55%	56%	53%	48%
White	53%	54%	56%	56%	56%	56%	55%	52%	52%	49%	45%
Minority	35%	35%	35%	35%	33%	32%	32%	30%	30%	27%	25%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	7,868	8,699	9,286	9,673	9,759	9,753	9,565	8,828	8,949	8,795	8,048
Alcohol-related disorders	1,540	1,559	1,536	1,495	1,378	1,252	1,152	953	850	718	539
Anxiety disorders	14,087	15,444	16,314	17,091	17,620	17,899	18,060	17,283	17,592	17,204	15,995
Attention-deficit, conduct, and disruptive behavior disorders	17,896	19,845	21,242	22,382	23,085	23,348	23,254	21,792	22,054	21,722	20,229
Delirium, dementia, and amnestic and other cognitive disorders	554	586	607	606	606	602	590	541	545	521	481
Developmental disorders	7,860	9,155	10,397	11,569	12,544	13,303	14,018	14,033	15,041	15,556	15,111
Disorders usually diagnosed in infancy, childhood, adolescence	2,971	3,404	3,807	4,147	4,405	4,574	4,678	4,527	4,689	4,697	4,516
Impulse control disorders, NEC	2,842	3,040	3,150	3,178	3,093	2,948	2,794	2,436	2,314	2,133	1,822
Miscellaneous disorders	5,427	5,769	5,896	5,971	5,927	5,804	5,609	5,162	5,084	4,811	4,344
Mood disorders	14,427	15,150	15,285	15,170	14,660	13,856	13,079	11,682	11,168	10,393	9,055
Personality disorders	1,377	1,397	1,386	1,370	1,344	1,237	1,152	1,023	946	861	709
Schizophrenia and other psychotic disorders	1,434	1,455	1,495	1,497	1,474	1,392	1,341	1,199	1,158	1,057	909
Screening and history of mental health and substance abuse	22,840	24,661	25,304	25,407	24,541	23,526	22,543	20,389	20,335	19,621	17,912
Substance-related disorders	3,996	4,034	3,966	3,844	3,554	3,256	2,947	2,445	2,248	1,981	1,582
Suicide and intentional self-inflicted injury	1,060	1,077	1,087	1,037	996	942	911	815	785	746	628



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

South Central Kentucky

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	44,362	47,626	50,004	51,783	53,237	54,077	54,839	52,011	55,486	56,482	54,311
Female	30,818	33,041	34,593	35,828	36,835	37,400	37,896	35,738	38,183	38,556	36,809
Male	13,544	14,585	15,411	15,955	16,402	16,677	16,943	16,273	17,303	17,926	17,502
White	38,716	41,600	43,767	45,398	46,763	47,605	48,304	45,824	48,983	49,869	47,734
Minority	3,339	3,535	3,629	3,660	3,612	3,469	3,353	3,000	3,275	3,354	3,258

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	66%	68%	69%	69%	69%	69%	65%	68%	67%	63%
Female	66%	69%	70%	71%	71%	71%	71%	66%	70%	69%	65%
Male	59%	62%	63%	64%	64%	64%	64%	62%	63%	63%	59%
White	64%	67%	68%	69%	69%	69%	69%	65%	68%	67%	63%
Minority	37%	38%	38%	38%	38%	36%	35%	31%	34%	32%	30%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	4,132	4,578	4,923	5,261	5,468	5,735	5,947	5,794	6,316	6,507	6,384
Alcohol-related disorders	4,183	4,638	4,934	5,137	5,391	5,427	5,485	5,163	5,385	5,459	5,238
Anxiety disorders	25,835	28,000	29,972	31,487	32,815	33,702	34,590	33,539	35,459	36,105	34,994
Attention-deficit, conduct, and disruptive behavior disorders	2,154	2,452	2,748	3,143	3,570	3,857	4,350	4,473	5,201	5,772	5,932
Delirium, dementia, and amnesic and other cognitive disorders	11,032	11,282	11,196	10,846	10,484	10,207	9,890	9,302	9,037	8,458	7,741
Developmental disorders	4,491	4,669	4,858	5,020	5,148	5,234	5,373	5,338	5,531	5,685	5,694
Disorders usually diagnosed in infancy, childhood, adolescence	583	633	675	725	778	823	898	912	1,018	1,131	1,179
Impulse control disorders, NEC	864	952	1,074	1,177	1,282	1,351	1,479	1,482	1,631	1,736	1,783
Miscellaneous disorders	9,561	10,531	11,325	12,135	12,673	13,159	13,551	12,931	13,603	13,777	13,170
Mood disorders	27,253	29,693	31,616	33,196	34,530	35,284	36,145	34,764	36,884	37,508	36,177
Personality disorders	3,554	3,800	4,038	4,218	4,332	4,347	4,393	4,212	4,385	4,383	4,283
Schizophrenia and other psychotic disorders	10,266	10,797	11,125	11,177	11,138	11,057	11,001	10,615	10,637	10,245	9,602
Screening and history of mental health and substance abuse	23,021	25,484	27,607	29,241	30,672	31,238	31,910	30,277	32,496	33,153	31,868
Substance-related disorders	6,393	7,114	7,636	8,167	8,665	8,901	9,266	8,868	9,581	9,943	9,657
Suicide and intentional self-inflicted injury	1,541	1,682	1,860	1,988	2,144	2,253	2,448	2,429	2,657	2,738	2,752



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Urban Triangle

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	72,154	80,890	86,779	92,132	95,444	97,543	99,181	95,039	99,870	101,377	96,624
Female	35,142	38,805	41,220	43,223	44,375	44,886	45,312	42,917	44,968	45,255	42,706
Male	37,012	42,085	45,559	48,909	51,069	52,657	53,869	52,122	54,902	56,122	53,918
White	48,173	54,454	58,229	62,062	64,474	66,241	67,314	64,466	67,950	69,273	66,298
Minority	20,106	21,934	23,380	24,278	24,579	24,456	24,482	23,084	23,931	23,870	22,465

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	51%	53%	55%	55%	55%	54%	53%	49%	49%	46%	42%
Female	51%	52%	53%	53%	52%	51%	50%	46%	46%	43%	38%
Male	52%	54%	56%	57%	57%	56%	56%	52%	53%	50%	45%
White	52%	54%	56%	56%	55%	54%	53%	49%	49%	46%	41%
Minority	42%	42%	44%	43%	42%	41%	41%	38%	38%	35%	32%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	19,981	22,474	24,098	25,370	25,939	25,999	25,778	23,843	23,964	23,104	21,266
Alcohol-related disorders	3,107	3,258	3,249	3,170	3,089	2,946	2,737	2,352	2,207	1,977	1,645
Anxiety disorders	22,754	25,060	26,602	27,976	28,721	28,907	28,891	27,254	27,639	27,089	25,135
Attention-deficit, conduct, and disruptive behavior disorders	35,717	40,389	43,974	47,211	49,360	50,591	51,220	48,602	49,567	49,155	46,481
Delirium, dementia, and amnestic and other cognitive disorders	1,353	1,447	1,532	1,569	1,608	1,619	1,633	1,568	1,603	1,520	1,393
Developmental disorders	15,542	18,531	21,283	23,912	26,087	27,924	29,697	30,039	32,521	33,925	33,256
Disorders usually diagnosed in infancy, childhood, adolescence	6,906	8,055	9,065	9,963	10,743	11,313	11,814	11,536	11,987	11,961	11,550
Impulse control disorders, NEC	4,763	5,127	5,391	5,478	5,444	5,252	5,057	4,538	4,286	3,921	3,406
Miscellaneous disorders	10,559	11,468	12,032	12,439	12,679	12,702	12,659	11,869	12,121	11,895	11,079
Mood disorders	26,303	28,110	28,842	29,113	28,832	28,006	26,826	24,173	23,602	22,288	19,796
Personality disorders	2,537	2,647	2,682	2,606	2,535	2,394	2,224	1,974	1,872	1,683	1,462
Schizophrenia and other psychotic disorders	3,649	3,823	3,880	3,984	3,990	3,924	3,828	3,567	3,553	3,303	2,921
Screening and history of mental health and substance abuse	44,054	47,723	49,629	50,367	49,790	48,436	47,150	43,172	43,213	41,680	37,825
Substance-related disorders	8,220	8,541	8,522	8,269	7,992	7,603	7,102	6,244	6,100	5,552	4,673
Suicide and intentional self-inflicted injury	2,740	2,864	2,974	3,033	3,035	3,010	2,964	2,810	2,856	2,738	2,411



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Urban Triangle

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	67,163	73,647	78,742	83,165	86,969	90,054	92,415	87,674	96,496	99,884	96,971
Female	49,807	54,413	58,154	61,477	64,241	66,150	67,688	63,459	69,966	71,690	68,767
Male	17,356	19,234	20,588	21,688	22,728	23,904	24,727	24,215	26,530	28,194	28,204
White	50,243	55,202	59,032	62,411	65,342	67,857	70,022	66,131	72,883	75,099	72,295
Minority	14,042	15,338	16,341	17,174	17,873	18,200	18,117	17,198	19,055	20,006	19,417

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	63%	64%	66%	66%	67%	67%	62%	66%	65%	61%
Female	61%	64%	66%	67%	68%	68%	68%	63%	67%	66%	62%
Male	56%	59%	61%	62%	63%	63%	64%	61%	64%	63%	59%
White	62%	65%	67%	68%	69%	69%	69%	64%	68%	67%	63%
Minority	45%	47%	48%	49%	49%	49%	49%	46%	49%	49%	45%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	8,409	9,366	10,253	11,011	11,765	12,412	12,977	12,725	14,045	14,794	14,627
Alcohol-related disorders	7,449	8,314	9,033	9,656	10,150	10,568	10,978	10,544	11,348	11,503	11,041
Anxiety disorders	33,417	37,219	40,461	43,326	45,675	47,743	49,839	48,586	52,641	54,198	52,811
Attention-deficit, conduct, and disruptive behavior disorders	4,466	5,164	5,884	6,683	7,452	8,379	9,347	9,789	11,390	12,481	12,732
Delirium, dementia, and amnesic and other cognitive disorders	17,556	18,321	18,549	18,298	18,020	17,996	17,740	16,717	16,706	16,027	14,747
Developmental disorders	6,568	7,067	7,511	7,840	8,073	8,412	8,750	8,787	9,336	9,755	9,904
Disorders usually diagnosed in infancy, childhood, adolescence	879	1,009	1,098	1,213	1,322	1,455	1,632	1,769	2,029	2,299	2,491
Impulse control disorders, NEC	1,253	1,420	1,569	1,723	1,899	2,076	2,312	2,422	2,726	2,975	3,095
Miscellaneous disorders	13,781	15,580	17,331	19,226	20,815	22,115	23,039	21,292	23,250	23,739	22,516
Mood disorders	38,692	42,900	46,532	49,838	52,464	54,675	56,873	54,943	59,608	61,362	59,977
Personality disorders	5,570	6,165	6,708	7,101	7,405	7,711	8,103	7,814	8,170	8,196	7,843
Schizophrenia and other psychotic disorders	16,590	17,727	18,401	18,676	18,939	19,325	19,516	18,787	19,099	18,857	17,864
Screening and history of mental health and substance abuse	37,558	42,476	46,986	51,077	54,639	56,996	59,002	56,104	62,415	64,656	62,622
Substance-related disorders	11,891	13,518	14,848	16,421	17,633	18,704	19,815	19,156	21,373	22,427	21,799
Suicide and intentional self-inflicted injury	2,861	3,210	3,583	3,933	4,253	4,620	5,129	5,132	5,813	6,104	6,092



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Eastern Kentucky

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	66,650	70,986	73,047	75,315	76,110	76,192	75,796	71,812	73,544	72,422	67,461
Female	33,529	35,162	35,709	36,446	36,489	36,133	35,779	33,591	34,224	33,421	31,012
Male	33,121	35,824	37,338	38,869	39,621	40,059	40,017	38,221	39,320	39,001	36,449
White	61,931	65,764	67,385	69,247	69,730	69,552	68,845	64,994	66,430	65,261	60,784
Minority	1,168	1,235	1,260	1,239	1,210	1,184	1,171	1,116	1,180	1,161	1,103

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	58%	60%	60%	61%	61%	61%	58%	59%	57%	52%
Female	58%	59%	60%	60%	60%	60%	59%	56%	57%	54%	50%
Male	54%	56%	59%	60%	61%	62%	62%	59%	61%	59%	54%
White	55%	57%	58%	59%	59%	59%	59%	56%	57%	55%	51%
Minority	19%	19%	18%	17%	15%	15%	14%	13%	14%	12%	12%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	12,151	12,933	13,506	13,989	14,056	14,036	13,739	12,737	12,640	12,114	10,959
Alcohol-related disorders	4,053	4,015	3,900	3,777	3,603	3,292	2,956	2,509	2,271	1,967	1,597
Anxiety disorders	26,948	28,384	29,174	30,209	30,696	30,708	30,442	28,844	28,910	28,037	26,057
Attention-deficit, conduct, and disruptive behavior disorders	24,602	26,800	28,344	29,814	30,517	30,755	30,525	28,566	28,687	27,717	25,660
Delirium, dementia, and amnestic and other cognitive disorders	916	906	905	897	863	838	816	748	724	681	601
Developmental disorders	17,206	19,242	21,060	22,861	24,378	25,233	25,607	24,981	25,986	25,849	24,591
Disorders usually diagnosed in infancy, childhood, adolescence	4,407	4,929	5,362	5,821	6,196	6,385	6,497	6,197	6,293	6,132	5,794
Impulse control disorders, NEC	2,256	2,367	2,398	2,407	2,345	2,215	2,088	1,873	1,720	1,534	1,281
Miscellaneous disorders	10,473	10,685	10,675	10,673	10,532	10,270	9,981	9,292	9,218	8,832	8,101
Mood disorders	22,158	22,213	21,709	21,159	20,105	18,782	17,424	15,288	14,328	12,984	11,102
Personality disorders	2,989	3,007	2,993	2,987	2,915	2,732	2,480	2,152	1,940	1,733	1,444
Schizophrenia and other psychotic disorders	2,442	2,406	2,391	2,348	2,262	2,159	2,010	1,779	1,659	1,515	1,285
Screening and history of mental health and substance abuse	42,063	44,023	44,493	44,748	43,928	42,388	40,919	37,546	37,385	35,846	32,451
Substance-related disorders	9,644	9,417	8,924	8,500	7,904	7,118	6,401	5,581	5,159	4,660	3,913
Suicide and intentional self-inflicted injury	1,943	1,926	1,847	1,801	1,681	1,579	1,480	1,321	1,224	1,110	919



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Eastern Kentucky

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	83,419	88,902	92,986	96,422	98,671	99,550	100,968	96,094	101,930	102,205	99,214
Female	55,431	58,854	61,416	63,410	64,818	65,478	66,427	62,850	66,672	66,603	64,220
Male	27,988	30,048	31,570	33,012	33,853	34,072	34,541	33,244	35,258	35,602	34,994
White	76,403	81,625	85,546	88,837	91,070	92,003	93,336	88,614	94,346	94,606	91,456
Minority	2,144	2,140	2,018	1,903	1,674	1,412	1,114	958	1,038	1,068	1,055

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	67%	69%	71%	71%	71%	71%	72%	68%	72%	71%	68%
Female	71%	73%	74%	75%	75%	75%	76%	71%	75%	75%	71%
Male	61%	63%	64%	65%	65%	65%	66%	63%	66%	66%	63%
White	67%	69%	70%	71%	71%	71%	71%	68%	71%	71%	68%
Minority	22%	22%	21%	20%	17%	15%	12%	10%	11%	11%	10%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	7,509	8,142	8,713	9,335	9,822	10,206	10,650	10,404	11,126	11,377	11,178
Alcohol-related disorders	9,251	10,024	10,555	11,250	11,691	11,769	12,011	11,537	11,999	11,909	11,452
Anxiety disorders	54,604	58,641	61,915	64,831	66,791	67,839	69,126	66,811	69,822	69,808	67,926
Attention-deficit, conduct, and disruptive behavior disorders	3,085	3,454	3,855	4,357	4,921	5,402	5,950	5,997	6,870	7,458	7,764
Delirium, dementia, and amnesic and other cognitive disorders	15,632	15,915	15,799	15,451	15,056	14,558	14,101	13,117	12,796	11,966	10,857
Developmental disorders	7,176	7,561	7,899	8,331	8,656	8,852	9,063	8,902	9,310	9,527	9,639
Disorders usually diagnosed in infancy, childhood, adolescence	863	945	987	1,049	1,121	1,201	1,263	1,304	1,450	1,573	1,674
Impulse control disorders, NEC	1,405	1,547	1,652	1,790	1,929	2,018	2,071	2,045	2,229	2,312	2,322
Miscellaneous disorders	23,137	25,105	26,740	28,372	29,643	30,236	30,459	29,328	30,245	29,960	28,750
Mood disorders	49,229	53,029	56,164	59,010	60,972	62,060	63,323	60,907	63,937	63,980	62,320
Personality disorders	6,961	7,535	8,011	8,581	9,045	9,084	9,019	8,662	8,856	8,791	8,604
Schizophrenia and other psychotic disorders	15,959	16,756	17,100	17,348	17,360	17,167	17,080	16,261	16,209	15,659	14,767
Screening and history of mental health and substance abuse	50,253	54,779	58,803	62,128	64,515	65,822	67,446	64,746	69,289	69,771	68,037
Substance-related disorders	18,462	20,197	21,685	23,408	24,789	25,783	26,748	26,004	27,847	28,225	27,689
Suicide and intentional self-inflicted injury	2,942	3,237	3,455	3,722	3,980	4,239	4,582	4,604	4,884	4,913	4,864



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Purchase Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	8,478	9,362	10,065	10,521	10,740	10,740	10,588	9,780	10,140	10,039	9,346
Female	4,102	4,503	4,790	4,929	4,934	4,889	4,842	4,417	4,553	4,502	4,155
Male	4,376	4,859	5,275	5,592	5,806	5,851	5,746	5,363	5,587	5,537	5,191
White	6,328	7,005	7,524	7,908	8,063	8,030	7,907	7,301	7,577	7,541	7,050
Minority	1,719	1,817	1,904	1,906	1,874	1,850	1,781	1,607	1,632	1,570	1,404

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	51%	54%	55%	55%	55%	53%	52%	48%	49%	47%	42%
Female	50%	53%	54%	53%	51%	50%	49%	45%	45%	43%	38%
Male	52%	55%	57%	57%	58%	56%	55%	52%	53%	51%	46%
White	50%	53%	54%	54%	53%	51%	50%	46%	47%	45%	40%
Minority	43%	45%	44%	43%	42%	40%	39%	35%	36%	34%	30%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,036	1,154	1,222	1,202	1,200	1,184	1,165	1,072	1,051	1,003	909
Alcohol-related disorders	243	230	229	218	219	194	201	167	153	134	107
Anxiety disorders	1,964	2,073	2,132	2,165	2,190	2,127	2,052	1,832	1,788	1,691	1,500
Attention-deficit, conduct, and disruptive behavior disorders	4,416	4,949	5,287	5,615	5,885	5,894	5,787	5,340	5,419	5,296	4,944
Delirium, dementia, and amnestic and other cognitive disorders	107	114	119	130	125	125	126	126	135	128	121
Developmental disorders	1,732	2,065	2,397	2,693	2,951	3,186	3,341	3,281	3,554	3,655	3,509
Disorders usually diagnosed in infancy, childhood, adolescence	556	637	712	806	872	921	937	881	914	899	857
Impulse control disorders, NEC	576	609	627	630	645	626	593	533	503	452	387
Miscellaneous disorders	1,163	1,242	1,282	1,312	1,340	1,315	1,264	1,150	1,145	1,061	940
Mood disorders	2,915	3,069	3,176	3,166	3,176	3,030	2,905	2,566	2,536	2,387	2,118
Personality disorders	240	239	227	221	208	190	186	167	165	144	122
Schizophrenia and other psychotic disorders	281	275	275	271	274	266	254	223	231	214	188
Screening and history of mental health and substance abuse	4,852	5,277	5,548	5,575	5,366	5,102	4,859	4,347	4,386	4,202	3,740
Substance-related disorders	600	599	585	586	565	510	488	410	391	352	296
Suicide and intentional self-inflicted injury	222	233	233	217	222	203	192	175	171	171	154



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Purchase Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	9,387	10,202	10,760	11,118	11,369	11,599	11,679	10,822	11,672	11,720	11,292
Female	7,115	7,670	8,041	8,311	8,481	8,616	8,637	7,884	8,489	8,429	8,037
Male	2,272	2,532	2,719	2,807	2,888	2,983	3,042	2,938	3,183	3,291	3,255
White	7,486	8,150	8,605	8,939	9,146	9,358	9,428	8,736	9,450	9,495	9,051
Minority	1,543	1,656	1,732	1,736	1,753	1,730	1,695	1,513	1,625	1,614	1,587

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	61%	64%	66%	66%	66%	66%	66%	62%	66%	65%	61%
Female	63%	67%	68%	68%	68%	69%	69%	64%	68%	67%	63%
Male	55%	58%	60%	61%	61%	60%	60%	58%	62%	61%	59%
White	61%	64%	66%	67%	67%	66%	66%	62%	66%	66%	61%
Minority	49%	52%	52%	52%	52%	51%	50%	46%	49%	47%	44%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	597	657	711	765	796	847	889	860	903	941	908
Alcohol-related disorders	784	878	959	1,010	1,066	1,089	1,092	1,030	1,101	1,098	1,059
Anxiety disorders	4,785	5,233	5,548	5,900	6,162	6,366	6,496	6,180	6,574	6,711	6,490
Attention-deficit, conduct, and disruptive behavior disorders	468	520	589	695	781	873	975	981	1,135	1,296	1,342
Delirium, dementia, and amnesic and other cognitive disorders	2,538	2,611	2,621	2,528	2,421	2,360	2,341	2,169	2,217	2,064	1,882
Developmental disorders	793	856	880	906	905	927	982	967	1,018	1,036	1,030
Disorders usually diagnosed in infancy, childhood, adolescence	58	63	73	77	84	95	106	109	129	149	163
Impulse control disorders, NEC	149	161	187	199	223	248	276	269	288	331	341
Miscellaneous disorders	1,762	1,987	2,164	2,317	2,461	2,545	2,650	2,466	2,627	2,673	2,540
Mood disorders	5,405	5,942	6,350	6,737	7,005	7,236	7,369	6,938	7,431	7,511	7,310
Personality disorders	596	670	739	758	793	796	832	818	890	915	881
Schizophrenia and other psychotic disorders	1,966	2,095	2,155	2,211	2,209	2,198	2,209	2,115	2,156	2,114	2,000
Screening and history of mental health and substance abuse	4,514	5,192	5,690	6,015	6,233	6,379	6,407	5,964	6,422	6,533	6,331
Substance-related disorders	1,160	1,327	1,443	1,525	1,622	1,711	1,758	1,657	1,785	1,812	1,734
Suicide and intentional self-inflicted injury	377	423	456	503	556	594	674	690	785	812	792



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Pennyriple Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	11,004	12,058	12,680	13,298	13,524	13,886	13,864	12,734	13,096	13,098	12,333
Female	5,400	5,903	6,077	6,327	6,376	6,487	6,443	5,863	6,019	5,949	5,520
Male	5,604	6,155	6,603	6,971	7,148	7,399	7,421	6,871	7,077	7,149	6,813
White	8,065	8,846	9,265	9,669	9,784	9,960	9,930	9,157	9,415	9,344	8,816
Minority	2,301	2,484	2,579	2,682	2,719	2,818	2,750	2,426	2,494	2,545	2,384

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	55%	57%	58%	57%	58%	57%	53%	54%	51%	46%
Female	54%	55%	56%	57%	56%	56%	55%	51%	51%	48%	43%
Male	53%	55%	58%	60%	59%	60%	60%	55%	56%	54%	49%
White	54%	55%	57%	58%	57%	57%	56%	52%	52%	50%	44%
Minority	41%	41%	43%	44%	43%	44%	43%	38%	39%	38%	35%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,888	2,066	2,156	2,236	2,234	2,244	2,238	2,002	2,017	1,954	1,810
Alcohol-related disorders	312	325	325	301	273	261	217	168	153	142	103
Anxiety disorders	2,602	2,738	2,875	2,962	2,924	2,934	2,896	2,598	2,569	2,417	2,171
Attention-deficit, conduct, and disruptive behavior disorders	5,448	6,023	6,422	6,859	7,005	7,181	7,133	6,559	6,577	6,428	6,055
Delirium, dementia, and amnestic and other cognitive disorders	177	187	177	183	190	192	187	175	179	173	151
Developmental disorders	2,069	2,418	2,736	3,086	3,314	3,542	3,728	3,654	3,845	3,949	3,798
Disorders usually diagnosed in infancy, childhood, adolescence	549	643	719	797	841	918	962	927	947	979	943
Impulse control disorders, NEC	1,092	1,140	1,154	1,144	1,095	1,067	989	861	822	758	657
Miscellaneous disorders	2,282	2,404	2,445	2,475	2,455	2,397	2,289	2,013	1,962	1,875	1,701
Mood disorders	3,818	4,030	4,060	4,098	3,942	3,786	3,561	3,098	2,967	2,741	2,374
Personality disorders	182	188	197	187	175	160	149	131	116	97	72
Schizophrenia and other psychotic disorders	340	340	352	353	346	322	305	279	264	252	223
Screening and history of mental health and substance abuse	6,304	6,784	7,046	7,138	7,097	7,125	6,984	6,260	6,235	6,072	5,605
Substance-related disorders	905	937	922	897	843	781	695	576	568	523	437
Suicide and intentional self-inflicted injury	272	288	283	291	283	258	238	205	192	172	137



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Pennyriple Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	11,006	11,830	12,437	12,821	13,137	13,445	13,853	12,699	13,855	14,079	13,417
Female	8,020	8,544	8,965	9,220	9,440	9,598	9,858	8,924	9,774	9,953	9,356
Male	2,986	3,286	3,472	3,601	3,697	3,847	3,995	3,775	4,081	4,126	4,061
White	8,764	9,369	9,902	10,208	10,459	10,700	11,047	10,025	11,028	11,207	10,604
Minority	1,800	1,984	2,030	2,071	2,082	2,105	2,109	1,967	2,115	2,149	2,066

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	64%	66%	67%	68%	68%	69%	63%	68%	67%	62%
Female	64%	67%	68%	69%	70%	70%	71%	64%	70%	69%	64%
Male	56%	59%	61%	63%	64%	64%	65%	61%	64%	63%	59%
White	63%	66%	68%	68%	69%	69%	70%	63%	69%	68%	63%
Minority	45%	48%	49%	51%	50%	50%	50%	46%	49%	48%	44%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,021	1,123	1,247	1,287	1,343	1,439	1,503	1,420	1,556	1,611	1,514
Alcohol-related disorders	884	960	1,020	1,073	1,116	1,111	1,165	1,076	1,156	1,177	1,113
Anxiety disorders	5,298	5,743	6,198	6,584	6,830	7,061	7,401	6,949	7,470	7,509	7,222
Attention-deficit, conduct, and disruptive behavior disorders	751	846	932	1,034	1,138	1,269	1,402	1,428	1,640	1,790	1,785
Delirium, dementia, and amnesic and other cognitive disorders	3,098	3,174	3,158	3,056	2,919	2,806	2,756	2,554	2,589	2,385	2,174
Developmental disorders	1,305	1,351	1,396	1,376	1,379	1,388	1,444	1,386	1,396	1,384	1,374
Disorders usually diagnosed in infancy, childhood, adolescence	98	109	120	133	141	157	158	169	185	192	203
Impulse control disorders, NEC	243	269	288	318	334	372	430	427	468	502	517
Miscellaneous disorders	2,195	2,418	2,547	2,673	2,812	2,900	3,036	2,800	3,049	3,111	2,989
Mood disorders	5,935	6,519	7,097	7,502	7,832	8,139	8,469	8,003	8,719	8,857	8,470
Personality disorders	512	569	581	607	654	669	674	648	682	682	628
Schizophrenia and other psychotic disorders	2,761	2,906	2,968	2,966	2,922	2,921	2,954	2,838	2,879	2,796	2,632
Screening and history of mental health and substance abuse	5,667	6,286	6,888	7,340	7,695	8,063	8,412	7,612	8,363	8,626	8,181
Substance-related disorders	1,260	1,404	1,572	1,698	1,791	1,917	2,055	1,893	2,122	2,240	2,102
Suicide and intentional self-inflicted injury	383	428	469	511	548	600	655	635	742	811	758



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Green River Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	10,933	11,922	12,620	13,276	13,591	13,778	13,931	13,196	13,619	13,688	12,975
Female	5,403	5,841	6,083	6,327	6,390	6,476	6,472	6,088	6,212	6,171	5,821
Male	5,530	6,081	6,537	6,949	7,201	7,302	7,459	7,108	7,407	7,517	7,154
White	9,016	9,800	10,311	10,824	11,041	11,190	11,307	10,684	10,986	11,039	10,441
Minority	1,205	1,271	1,359	1,422	1,427	1,406	1,364	1,260	1,336	1,335	1,294

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	58%	61%	60%	60%	59%	59%	55%	55%	52%	48%
Female	57%	58%	60%	59%	58%	57%	56%	52%	51%	48%	44%
Male	57%	58%	61%	62%	61%	61%	62%	58%	58%	56%	51%
White	56%	57%	59%	59%	58%	57%	57%	53%	53%	50%	46%
Minority	39%	37%	40%	39%	37%	36%	35%	32%	33%	31%	29%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,298	2,491	2,631	2,739	2,732	2,730	2,707	2,474	2,426	2,334	2,168
Alcohol-related disorders	401	415	384	374	335	323	294	262	249	215	176
Anxiety disorders	3,246	3,490	3,624	3,829	3,850	3,889	3,848	3,570	3,541	3,393	3,120
Attention-deficit, conduct, and disruptive behavior disorders	5,629	6,174	6,614	7,014	7,266	7,357	7,330	6,830	6,878	6,760	6,366
Delirium, dementia, and amnestic and other cognitive disorders	158	167	177	175	188	184	178	157	158	147	132
Developmental disorders	2,126	2,512	2,912	3,253	3,577	3,833	4,130	4,142	4,397	4,564	4,478
Disorders usually diagnosed in infancy, childhood, adolescence	727	811	901	981	1,068	1,128	1,174	1,141	1,196	1,210	1,163
Impulse control disorders, NEC	990	1,060	1,103	1,142	1,161	1,136	1,104	994	938	877	779
Miscellaneous disorders	1,602	1,700	1,726	1,722	1,710	1,716	1,683	1,548	1,569	1,521	1,423
Mood disorders	4,675	4,925	5,056	5,073	4,996	4,879	4,675	4,194	4,054	3,835	3,435
Personality disorders	433	447	425	405	409	380	359	321	293	252	217
Schizophrenia and other psychotic disorders	513	519	530	537	512	499	477	430	397	346	300
Screening and history of mental health and substance abuse	6,765	7,268	7,511	7,503	7,270	7,095	6,889	6,336	6,351	6,178	5,639
Substance-related disorders	1,097	1,131	1,073	1,051	957	895	825	728	683	584	476
Suicide and intentional self-inflicted injury	338	339	328	316	315	295	300	248	237	211	192



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Green River Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	10,113	11,096	11,798	12,471	13,055	13,365	13,675	12,813	14,077	14,298	13,794
Female	7,429	8,157	8,667	9,185	9,560	9,751	9,948	9,219	10,042	10,167	9,727
Male	2,684	2,939	3,131	3,286	3,495	3,614	3,727	3,594	4,035	4,131	4,067
White	8,778	9,642	10,209	10,785	11,302	11,612	11,886	11,079	12,277	12,419	11,852
Minority	935	1,008	1,092	1,147	1,156	1,124	1,094	1,041	1,097	1,131	1,140

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	66%	68%	70%	70%	70%	70%	65%	70%	68%	64%
Female	66%	69%	71%	72%	73%	72%	72%	67%	72%	71%	66%
Male	58%	60%	62%	63%	65%	64%	64%	61%	65%	64%	60%
White	64%	67%	68%	70%	70%	70%	70%	65%	70%	69%	64%
Minority	43%	42%	47%	47%	46%	44%	42%	41%	42%	41%	38%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,103	1,277	1,414	1,507	1,552	1,571	1,634	1,521	1,659	1,712	1,661
Alcohol-related disorders	925	1,015	1,077	1,184	1,280	1,298	1,392	1,345	1,444	1,478	1,444
Anxiety disorders	5,171	5,789	6,238	6,753	7,153	7,447	7,778	7,524	8,184	8,256	8,038
Attention-deficit, conduct, and disruptive behavior disorders	964	1,091	1,231	1,426	1,638	1,763	1,987	2,065	2,374	2,574	2,635
Delirium, dementia, and amnesic and other cognitive disorders	2,374	2,414	2,460	2,398	2,349	2,318	2,244	2,145	2,157	2,076	1,934
Developmental disorders	1,157	1,208	1,263	1,313	1,381	1,400	1,427	1,443	1,487	1,541	1,560
Disorders usually diagnosed in infancy, childhood, adolescence	105	117	130	145	165	173	196	210	230	254	282
Impulse control disorders, NEC	221	250	269	295	327	353	400	397	467	516	520
Miscellaneous disorders	2,259	2,562	2,813	3,060	3,302	3,462	3,515	3,210	3,470	3,507	3,354
Mood disorders	6,058	6,801	7,337	7,939	8,408	8,652	8,988	8,616	9,323	9,475	9,209
Personality disorders	901	977	1,071	1,117	1,169	1,209	1,232	1,202	1,277	1,298	1,233
Schizophrenia and other psychotic disorders	2,397	2,548	2,658	2,715	2,767	2,799	2,858	2,754	2,830	2,810	2,713
Screening and history of mental health and substance abuse	5,373	6,181	6,725	7,385	7,943	8,221	8,525	7,955	8,878	8,982	8,615
Substance-related disorders	1,617	1,844	2,027	2,242	2,440	2,573	2,729	2,643	2,916	2,984	2,838
Suicide and intentional self-inflicted injury	402	455	513	554	633	678	749	773	884	922	904



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Barren River Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	12,632	14,102	15,013	15,585	15,890	16,278	16,397	15,707	16,397	16,738	15,863
Female	6,206	6,878	7,281	7,510	7,608	7,722	7,702	7,274	7,630	7,724	7,272
Male	6,426	7,224	7,732	8,075	8,282	8,556	8,695	8,433	8,767	9,014	8,591
White	10,584	11,824	12,566	12,980	13,190	13,526	13,566	12,996	13,515	13,806	13,096
Minority	1,547	1,701	1,781	1,865	1,867	1,869	1,841	1,736	1,833	1,844	1,763

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	53%	55%	54%	54%	53%	53%	50%	51%	48%	43%
Female	52%	53%	54%	54%	53%	52%	51%	48%	49%	46%	41%
Male	51%	53%	55%	55%	55%	55%	55%	52%	53%	50%	46%
White	52%	53%	55%	54%	53%	53%	52%	49%	50%	48%	43%
Minority	39%	40%	40%	40%	38%	37%	37%	34%	34%	30%	27%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,344	2,605	2,809	2,895	2,901	2,905	2,808	2,585	2,598	2,533	2,324
Alcohol-related disorders	410	424	415	399	382	353	327	264	239	216	161
Anxiety disorders	3,969	4,406	4,764	4,955	5,211	5,412	5,514	5,357	5,546	5,501	5,160
Attention-deficit, conduct, and disruptive behavior disorders	5,841	6,534	7,042	7,377	7,571	7,724	7,727	7,259	7,364	7,304	6,780
Delirium, dementia, and amnestic and other cognitive disorders	168	178	194	192	195	191	192	176	187	187	174
Developmental disorders	2,701	3,192	3,621	3,988	4,339	4,627	4,918	4,931	5,259	5,471	5,346
Disorders usually diagnosed in infancy, childhood, adolescence	1,003	1,162	1,321	1,453	1,548	1,641	1,692	1,639	1,714	1,711	1,659
Impulse control disorders, NEC	931	1,021	1,069	1,069	1,049	985	920	804	767	716	610
Miscellaneous disorders	1,938	2,076	2,149	2,208	2,187	2,176	2,085	1,937	1,888	1,816	1,659
Mood disorders	4,640	4,888	5,021	4,931	4,759	4,506	4,260	3,847	3,667	3,396	2,961
Personality disorders	267	279	275	269	269	246	220	185	179	159	121
Schizophrenia and other psychotic disorders	475	494	506	515	518	486	456	416	403	360	312
Screening and history of mental health and substance abuse	8,452	9,187	9,417	9,272	8,888	8,529	8,166	7,408	7,542	7,375	6,754
Substance-related disorders	1,090	1,138	1,111	1,086	995	927	853	724	692	605	477
Suicide and intentional self-inflicted injury	307	318	320	308	301	268	252	221	202	193	155



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Barren River Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	15,292	16,659	17,599	17,921	18,227	18,564	18,773	17,580	18,631	19,048	18,555
Female	10,717	11,671	12,304	12,555	12,765	13,005	13,112	12,168	12,945	13,143	12,718
Male	4,575	4,988	5,295	5,366	5,462	5,559	5,661	5,412	5,686	5,905	5,837
White	12,884	14,076	14,900	15,177	15,453	15,751	15,920	14,919	15,843	16,176	15,744
Minority	1,657	1,770	1,825	1,830	1,814	1,788	1,773	1,588	1,695	1,761	1,686

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	68%	69%	70%	69%	69%	69%	64%	68%	66%	62%
Female	66%	69%	71%	71%	71%	71%	70%	65%	70%	68%	63%
Male	61%	64%	65%	66%	66%	65%	65%	63%	64%	62%	59%
White	65%	68%	70%	70%	70%	70%	69%	65%	68%	67%	63%
Minority	43%	44%	45%	45%	43%	42%	41%	37%	40%	37%	34%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,292	1,457	1,550	1,640	1,684	1,757	1,835	1,744	1,873	1,927	1,923
Alcohol-related disorders	1,444	1,618	1,727	1,743	1,822	1,832	1,849	1,748	1,815	1,834	1,777
Anxiety disorders	8,492	9,293	10,014	10,371	10,742	11,053	11,287	10,903	11,476	11,706	11,403
Attention-deficit, conduct, and disruptive behavior disorders	783	913	1,009	1,136	1,296	1,402	1,578	1,612	1,852	2,050	2,117
Delirium, dementia, and amnesic and other cognitive disorders	4,220	4,339	4,307	4,135	3,955	3,876	3,735	3,489	3,394	3,155	2,882
Developmental disorders	1,436	1,531	1,587	1,624	1,640	1,655	1,739	1,719	1,779	1,837	1,808
Disorders usually diagnosed in infancy, childhood, adolescence	124	147	153	165	186	207	237	256	293	333	347
Impulse control disorders, NEC	260	286	318	354	382	400	443	442	480	523	544
Miscellaneous disorders	3,273	3,681	3,973	4,213	4,358	4,552	4,753	4,487	4,717	4,814	4,631
Mood disorders	9,430	10,411	11,181	11,623	12,018	12,245	12,476	11,887	12,585	12,831	12,466
Personality disorders	854	916	982	1,015	1,014	1,022	1,036	995	1,024	1,030	1,002
Schizophrenia and other psychotic disorders	3,666	3,887	4,033	4,064	4,031	4,066	4,049	3,915	3,909	3,739	3,492
Screening and history of mental health and substance abuse	8,231	9,261	10,145	10,476	10,794	10,969	11,163	10,524	11,198	11,464	11,233
Substance-related disorders	1,917	2,171	2,346	2,464	2,597	2,645	2,754	2,641	2,792	2,883	2,807
Suicide and intentional self-inflicted injury	483	545	614	644	705	737	810	813	888	924	943



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lincoln Trail Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	10,320	11,619	12,578	13,352	13,819	14,022	14,186	13,524	14,185	14,595	13,806
Female	4,962	5,570	5,916	6,227	6,369	6,396	6,484	6,163	6,438	6,500	6,131
Male	5,358	6,049	6,662	7,125	7,450	7,626	7,702	7,361	7,747	8,095	7,675
White	8,571	9,632	10,395	11,023	11,369	11,505	11,543	10,953	11,513	11,855	11,231
Minority	1,147	1,260	1,335	1,375	1,374	1,385	1,414	1,317	1,371	1,404	1,336

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	49%	51%	53%	54%	54%	54%	54%	51%	51%	48%	44%
Female	49%	51%	52%	52%	52%	51%	51%	47%	47%	44%	40%
Male	50%	52%	55%	56%	57%	57%	57%	54%	55%	52%	48%
White	49%	51%	53%	54%	54%	54%	53%	49%	50%	47%	43%
Minority	33%	34%	33%	33%	32%	31%	32%	29%	30%	27%	26%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,503	2,882	3,203	3,394	3,483	3,524	3,539	3,332	3,475	3,559	3,336
Alcohol-related disorders	428	442	446	452	429	404	360	310	291	257	195
Anxiety disorders	3,098	3,459	3,724	3,881	3,982	3,975	4,042	3,887	3,987	3,989	3,689
Attention-deficit, conduct, and disruptive behavior disorders	5,365	6,086	6,671	7,116	7,518	7,626	7,683	7,328	7,546	7,567	7,123
Delirium, dementia, and amnestic and other cognitive disorders	160	173	183	184	184	184	189	165	171	168	154
Developmental disorders	2,241	2,635	3,094	3,479	3,818	4,045	4,236	4,264	4,594	4,852	4,702
Disorders usually diagnosed in infancy, childhood, adolescence	923	1,072	1,211	1,330	1,427	1,499	1,546	1,525	1,595	1,649	1,578
Impulse control disorders, NEC	1,000	1,083	1,120	1,139	1,128	1,087	1,056	943	903	843	719
Miscellaneous disorders	1,327	1,430	1,494	1,564	1,584	1,570	1,563	1,476	1,496	1,444	1,314
Mood disorders	3,998	4,347	4,465	4,516	4,447	4,204	4,040	3,709	3,614	3,465	3,032
Personality disorders	652	686	687	697	700	645	623	575	535	495	424
Schizophrenia and other psychotic disorders	375	386	414	418	409	388	397	354	343	323	282
Screening and history of mental health and substance abuse	6,555	7,214	7,497	7,644	7,398	7,101	6,845	6,184	6,090	5,935	5,431
Substance-related disorders	1,026	1,068	1,105	1,105	1,068	991	902	763	705	634	501
Suicide and intentional self-inflicted injury	242	258	268	255	258	255	260	249	245	244	215



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lincoln Trail Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	9,705	10,485	11,253	12,157	12,875	13,215	13,468	12,819	14,007	14,333	13,595
Female	7,058	7,630	8,193	8,848	9,385	9,564	9,733	9,182	10,006	10,159	9,596
Male	2,647	2,855	3,060	3,309	3,490	3,651	3,735	3,637	4,001	4,174	3,999
White	8,294	8,944	9,626	10,438	11,093	11,430	11,670	11,126	12,145	12,439	11,720
Minority	922	1,002	1,066	1,123	1,146	1,112	1,084	958	1,085	1,113	1,078

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	59%	60%	63%	64%	65%	65%	61%	65%	64%	60%
Female	58%	61%	63%	65%	67%	67%	68%	63%	67%	66%	62%
Male	51%	52%	54%	57%	58%	59%	59%	56%	60%	59%	55%
White	57%	59%	61%	63%	65%	65%	65%	61%	65%	64%	60%
Minority	35%	37%	37%	39%	40%	38%	37%	34%	37%	36%	34%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	915	1,020	1,119	1,201	1,271	1,372	1,442	1,402	1,531	1,601	1,566
Alcohol-related disorders	883	984	1,091	1,209	1,274	1,325	1,361	1,254	1,330	1,337	1,258
Anxiety disorders	5,034	5,520	6,090	6,656	7,103	7,371	7,650	7,416	8,002	8,167	7,816
Attention-deficit, conduct, and disruptive behavior disorders	551	627	709	860	997	1,064	1,195	1,195	1,426	1,600	1,643
Delirium, dementia, and amnestic and other cognitive disorders	2,341	2,394	2,404	2,399	2,402	2,387	2,388	2,264	2,243	2,146	1,978
Developmental disorders	989	1,039	1,101	1,179	1,227	1,251	1,277	1,288	1,362	1,416	1,466
Disorders usually diagnosed in infancy, childhood, adolescence	181	191	201	230	252	261	282	281	315	339	367
Impulse control disorders, NEC	224	250	293	319	356	384	410	418	477	489	498
Miscellaneous disorders	1,849	2,061	2,277	2,554	2,747	2,883	2,973	2,807	2,984	3,043	2,897
Mood disorders	5,691	6,313	6,875	7,464	7,952	8,290	8,608	8,291	9,022	9,239	8,801
Personality disorders	1,073	1,177	1,279	1,379	1,425	1,446	1,490	1,427	1,524	1,538	1,501
Schizophrenia and other psychotic disorders	2,170	2,310	2,397	2,471	2,512	2,533	2,542	2,429	2,511	2,458	2,288
Screening and history of mental health and substance abuse	4,741	5,330	5,930	6,676	7,309	7,517	7,659	7,170	7,875	8,039	7,594
Substance-related disorders	1,242	1,417	1,580	1,752	1,896	1,985	2,097	1,935	2,100	2,186	2,055
Suicide and intentional self-inflicted injury	282	300	351	389	430	450	515	500	561	567	570



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

KIPDA Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	31,949	35,571	38,318	40,777	42,251	42,731	43,200	41,165	42,440	42,825	40,831
Female	15,252	16,764	17,953	18,938	19,466	19,463	19,496	18,469	18,907	18,906	17,807
Male	16,697	18,807	20,365	21,839	22,785	23,268	23,704	22,696	23,533	23,919	23,024
White	15,999	18,168	19,607	21,174	21,994	22,444	22,730	21,680	22,298	22,649	21,763
Minority	14,087	15,247	16,217	16,809	17,132	16,977	16,903	15,822	16,275	16,142	15,165

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	48%	50%	52%	52%	52%	50%	49%	46%	45%	43%	39%
Female	47%	48%	50%	50%	49%	47%	46%	42%	42%	39%	35%
Male	49%	52%	54%	55%	55%	54%	53%	49%	49%	47%	43%
White	48%	50%	52%	52%	51%	50%	48%	44%	44%	42%	37%
Minority	42%	43%	45%	45%	44%	43%	42%	38%	38%	36%	33%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	9,598	10,793	11,655	12,248	12,543	12,534	12,398	11,426	11,311	10,864	10,026
Alcohol-related disorders	1,237	1,267	1,249	1,227	1,183	1,111	1,024	870	784	690	578
Anxiety disorders	8,764	9,596	10,073	10,549	10,761	10,605	10,457	9,754	9,602	9,268	8,382
Attention-deficit, conduct, and disruptive behavior disorders	15,845	17,923	19,579	21,023	21,969	22,357	22,496	21,241	21,363	21,091	19,873
Delirium, dementia, and amnestic and other cognitive disorders	577	612	641	675	670	656	660	631	639	618	582
Developmental disorders	6,580	7,895	9,055	10,235	11,154	11,797	12,475	12,682	13,439	14,018	13,880
Disorders usually diagnosed in infancy, childhood, adolescence	2,814	3,279	3,642	4,016	4,346	4,586	4,835	4,756	4,924	4,890	4,753
Impulse control disorders, NEC	2,914	3,173	3,324	3,402	3,391	3,273	3,166	2,865	2,741	2,518	2,169
Miscellaneous disorders	3,425	3,634	3,777	3,934	3,973	3,974	3,928	3,674	3,665	3,581	3,331
Mood disorders	11,757	12,619	12,971	13,187	13,128	12,704	12,223	11,044	10,675	10,007	8,869
Personality disorders	909	961	1,003	1,013	1,003	977	939	869	856	778	664
Schizophrenia and other psychotic disorders	1,764	1,860	1,897	1,946	1,959	1,931	1,905	1,787	1,795	1,661	1,445
Screening and history of mental health and substance abuse	19,143	20,684	21,644	22,040	21,950	21,169	20,537	18,795	18,558	17,766	16,136
Substance-related disorders	3,388	3,519	3,506	3,466	3,349	3,205	2,972	2,607	2,527	2,325	1,990
Suicide and intentional self-inflicted injury	1,104	1,157	1,219	1,267	1,273	1,252	1,237	1,164	1,183	1,127	993



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

KIPDA Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	26,694	29,118	31,451	33,409	35,376	36,716	37,511	35,884	39,388	40,707	39,641
Female	20,056	21,724	23,441	24,942	26,372	27,126	27,604	26,052	28,581	29,216	28,123
Male	6,638	7,394	8,010	8,467	9,004	9,590	9,907	9,832	10,807	11,491	11,518
White	16,320	17,772	19,173	20,351	21,597	22,560	23,213	22,008	24,130	24,745	23,751
Minority	9,208	10,056	10,869	11,550	12,204	12,512	12,546	12,065	13,324	13,918	13,629

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	58%	60%	62%	63%	63%	63%	59%	63%	62%	58%
Female	56%	59%	61%	63%	64%	64%	64%	59%	63%	63%	59%
Male	51%	55%	57%	59%	60%	60%	61%	59%	61%	60%	57%
White	57%	60%	62%	65%	65%	66%	65%	61%	65%	64%	59%
Minority	46%	49%	50%	51%	52%	52%	53%	49%	52%	52%	48%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	3,126	3,481	3,789	3,984	4,308	4,545	4,768	4,673	5,176	5,476	5,491
Alcohol-related disorders	3,007	3,367	3,669	3,932	4,176	4,380	4,530	4,410	4,685	4,768	4,579
Anxiety disorders	10,972	12,204	13,361	14,392	15,329	16,081	16,878	16,704	18,037	18,687	18,183
Attention-deficit, conduct, and disruptive behavior disorders	1,788	2,049	2,354	2,607	2,897	3,255	3,712	3,975	4,652	5,040	5,177
Delirium, dementia, and amnesic and other cognitive disorders	7,057	7,387	7,557	7,481	7,367	7,463	7,477	7,198	7,222	6,959	6,388
Developmental disorders	2,504	2,688	2,881	2,982	3,073	3,189	3,300	3,338	3,563	3,731	3,812
Disorders usually diagnosed in infancy, childhood, adolescence	358	402	445	495	550	609	678	742	840	961	1,062
Impulse control disorders, NEC	612	686	781	857	953	1,025	1,166	1,222	1,392	1,532	1,629
Miscellaneous disorders	3,983	4,512	5,085	5,806	6,500	6,982	7,207	6,642	7,352	7,443	7,014
Mood disorders	14,587	16,047	17,511	18,706	19,891	20,733	21,522	21,004	22,753	23,519	23,099
Personality disorders	1,567	1,727	1,859	1,896	1,985	2,075	2,184	2,147	2,242	2,278	2,192
Schizophrenia and other psychotic disorders	6,858	7,333	7,657	7,806	7,987	8,154	8,257	8,011	8,134	8,100	7,724
Screening and history of mental health and substance abuse	14,417	16,294	18,176	19,886	21,751	22,734	23,272	22,220	24,689	25,450	24,641
Substance-related disorders	4,378	5,004	5,513	6,077	6,537	6,877	7,271	7,084	7,768	8,131	7,861
Suicide and intentional self-inflicted injury	996	1,127	1,283	1,411	1,538	1,685	1,859	1,880	2,178	2,286	2,280



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Northern KY Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	13,232	15,065	16,220	17,230	17,784	18,617	19,197	18,430	19,804	20,385	19,516
Female	6,407	7,204	7,688	8,082	8,244	8,592	8,820	8,368	9,018	9,160	8,711
Male	6,825	7,861	8,532	9,148	9,540	10,025	10,377	10,062	10,786	11,225	10,805
White	11,249	12,802	13,723	14,545	14,985	15,718	16,143	15,441	16,572	17,065	16,349
Minority	1,378	1,526	1,646	1,701	1,696	1,701	1,742	1,698	1,820	1,880	1,812

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	56%	58%	58%	57%	56%	56%	52%	53%	49%	45%
Female	54%	55%	57%	56%	55%	54%	53%	49%	50%	46%	42%
Male	55%	57%	59%	59%	59%	59%	59%	55%	56%	52%	47%
White	54%	56%	58%	58%	57%	56%	56%	52%	52%	49%	44%
Minority	39%	38%	38%	36%	34%	33%	33%	31%	32%	30%	27%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,794	3,145	3,324	3,506	3,534	3,606	3,623	3,371	3,446	3,369	3,074
Alcohol-related disorders	654	711	709	687	701	659	606	532	516	476	395
Anxiety disorders	4,147	4,559	4,819	4,963	4,971	5,029	4,934	4,539	4,602	4,462	4,100
Attention-deficit, conduct, and disruptive behavior disorders	7,076	8,113	8,978	9,772	10,390	11,012	11,406	10,971	11,494	11,611	11,141
Delirium, dementia, and amnestic and other cognitive disorders	288	298	332	337	342	350	357	344	369	348	309
Developmental disorders	3,223	3,807	4,325	4,831	5,290	5,724	6,151	6,102	6,765	7,074	6,906
Disorders usually diagnosed in infancy, childhood, adolescence	1,458	1,705	1,939	2,138	2,273	2,391	2,474	2,410	2,500	2,482	2,365
Impulse control disorders, NEC	665	709	763	771	759	756	733	656	611	584	526
Miscellaneous disorders	2,521	2,790	2,916	3,026	3,085	3,070	3,063	2,821	2,912	2,859	2,658
Mood disorders	5,012	5,446	5,636	5,692	5,592	5,521	5,325	4,757	4,719	4,484	4,022
Personality disorders	693	715	698	656	612	562	495	424	377	329	286
Schizophrenia and other psychotic disorders	578	606	620	643	631	631	614	574	576	563	511
Screening and history of mental health and substance abuse	8,548	9,441	9,857	10,034	9,820	9,783	9,659	8,892	9,120	8,950	8,160
Substance-related disorders	1,452	1,531	1,553	1,484	1,477	1,409	1,310	1,168	1,152	1,055	877
Suicide and intentional self-inflicted injury	544	571	568	580	598	619	600	598	626	598	528



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Northern KY Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	12,442	13,784	14,914	15,681	16,411	17,091	17,872	16,567	18,490	19,171	18,382
Female	9,293	10,252	11,095	11,662	12,177	12,675	13,220	12,164	13,591	13,995	13,256
Male	3,149	3,532	3,819	4,019	4,234	4,416	4,652	4,403	4,899	5,176	5,126
White	11,117	12,306	13,318	14,013	14,722	15,345	16,086	14,895	16,642	17,231	16,475
Minority	859	999	1,072	1,119	1,109	1,122	1,115	993	1,134	1,194	1,091

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	69%	70%	71%	71%	72%	65%	70%	69%	64%
Female	66%	69%	71%	71%	72%	73%	73%	66%	71%	70%	65%
Male	59%	61%	64%	65%	66%	66%	67%	63%	67%	66%	62%
White	65%	68%	70%	71%	72%	72%	72%	66%	71%	70%	65%
Minority	39%	42%	43%	42%	41%	41%	40%	35%	39%	37%	32%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,523	1,687	1,865	1,969	2,072	2,200	2,328	2,243	2,422	2,565	2,495
Alcohol-related disorders	1,588	1,792	1,953	2,068	2,173	2,295	2,373	2,217	2,424	2,441	2,308
Anxiety disorders	6,992	7,772	8,502	9,050	9,536	10,047	10,587	10,117	11,078	11,412	11,017
Attention-deficit, conduct, and disruptive behavior disorders	996	1,159	1,353	1,580	1,798	2,007	2,232	2,262	2,615	2,854	2,820
Delirium, dementia, and amnesic and other cognitive disorders	3,025	3,282	3,372	3,391	3,363	3,368	3,341	3,158	3,215	3,038	2,800
Developmental disorders	1,133	1,225	1,341	1,388	1,427	1,510	1,603	1,633	1,747	1,819	1,840
Disorders usually diagnosed in infancy, childhood, adolescence	160	186	201	227	243	273	313	326	378	430	465
Impulse control disorders, NEC	222	262	283	307	330	369	402	407	469	507	517
Miscellaneous disorders	2,792	3,153	3,641	4,085	4,393	4,660	4,897	4,362	4,761	4,853	4,475
Mood disorders	7,765	8,666	9,493	10,171	10,739	11,272	11,890	11,293	12,427	12,803	12,308
Personality disorders	1,294	1,463	1,593	1,745	1,856	1,971	2,151	1,999	2,091	2,039	1,882
Schizophrenia and other psychotic disorders	2,910	3,198	3,348	3,446	3,492	3,642	3,666	3,547	3,721	3,645	3,435
Screening and history of mental health and substance abuse	7,544	8,491	9,539	10,309	10,900	11,513	12,183	11,298	12,765	13,291	12,726
Substance-related disorders	2,261	2,598	2,892	3,168	3,364	3,656	3,891	3,704	4,141	4,369	4,217
Suicide and intentional self-inflicted injury	672	755	861	928	982	1,080	1,239	1,212	1,377	1,475	1,431



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Buffalo Trace Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,424	3,671	3,957	4,064	4,169	4,164	4,265	3,987	4,143	4,202	3,855
Female	1,741	1,834	1,947	1,994	2,019	1,988	2,030	1,883	1,935	1,953	1,777
Male	1,683	1,837	2,010	2,070	2,150	2,176	2,235	2,104	2,208	2,249	2,078
White	3,112	3,318	3,576	3,660	3,743	3,724	3,799	3,537	3,663	3,724	3,411
Minority	159	166	168	174	172	170	173	169	188	182	169

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	55%	58%	57%	57%	56%	56%	52%	52%	50%	45%
Female	57%	56%	58%	56%	56%	55%	54%	50%	50%	48%	43%
Male	53%	53%	57%	57%	58%	57%	57%	53%	54%	52%	48%
White	54%	54%	57%	56%	56%	55%	54%	50%	51%	49%	44%
Minority	32%	30%	30%	30%	29%	27%	25%	25%	27%	24%	22%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	850	888	971	981	1,016	1,022	1,057	1,049	1,092	1,062	973
Alcohol-related disorders	193	189	186	172	173	151	140	107	104	88	65
Anxiety disorders	1,084	1,138	1,231	1,247	1,247	1,208	1,214	1,106	1,116	1,083	981
Attention-deficit, conduct, and disruptive behavior disorders	1,441	1,568	1,715	1,778	1,829	1,847	1,874	1,694	1,748	1,726	1,567
Delirium, dementia, and amnestic and other cognitive disorders	60	59	59	61	59	56	56	45	42	40	34
Developmental disorders	951	1,071	1,231	1,322	1,467	1,529	1,602	1,526	1,641	1,661	1,573
Disorders usually diagnosed in infancy, childhood, adolescence	289	316	364	391	422	421	426	387	411	409	382
Impulse control disorders, NEC	115	125	132	133	124	120	119	96	83	70	61
Miscellaneous disorders	579	600	633	643	648	634	629	571	585	569	510
Mood disorders	1,181	1,217	1,289	1,244	1,192	1,121	1,086	929	914	849	734
Personality disorders	309	327	337	335	339	334	302	248	225	200	170
Schizophrenia and other psychotic disorders	206	197	200	201	205	207	203	178	166	165	149
Screening and history of mental health and substance abuse	2,439	2,573	2,740	2,758	2,758	2,660	2,632	2,379	2,390	2,332	2,111
Substance-related disorders	386	398	391	375	359	319	289	227	204	170	131
Suicide and intentional self-inflicted injury	86	86	93	90	79	79	78	70	73	65	54



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Buffalo Trace Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,783	4,050	4,251	4,462	4,637	4,702	4,917	4,598	5,034	5,104	4,898
Female	2,634	2,792	2,940	3,071	3,192	3,250	3,417	3,131	3,445	3,509	3,303
Male	1,149	1,258	1,311	1,391	1,445	1,452	1,500	1,467	1,589	1,595	1,595
White	3,423	3,665	3,857	4,035	4,215	4,291	4,472	4,181	4,584	4,654	4,430
Minority	153	168	158	178	165	143	154	138	158	156	153

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	65%	66%	67%	69%	67%	68%	63%	68%	67%	63%
Female	66%	68%	69%	71%	72%	71%	71%	65%	71%	70%	65%
Male	57%	59%	59%	60%	62%	60%	61%	58%	61%	60%	58%
White	63%	65%	66%	67%	68%	67%	67%	62%	67%	67%	63%
Minority	27%	28%	27%	29%	28%	25%	25%	23%	26%	23%	21%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	332	377	413	461	499	520	571	551	612	664	651
Alcohol-related disorders	358	396	423	470	479	502	525	516	537	550	534
Anxiety disorders	2,132	2,305	2,486	2,640	2,722	2,798	2,940	2,789	2,957	2,982	2,869
Attention-deficit, conduct, and disruptive behavior disorders	197	219	257	302	338	380	402	400	460	474	491
Delirium, dementia, and amnestic and other cognitive disorders	879	887	887	855	859	820	797	759	764	700	660
Developmental disorders	402	432	447	464	489	519	534	508	523	538	546
Disorders usually diagnosed in infancy, childhood, adolescence	53	59	66	76	83	88	89	88	94	101	104
Impulse control disorders, NEC	61	68	76	88	95	93	100	101	124	133	122
Miscellaneous disorders	731	788	867	943	1,026	1,077	1,115	1,046	1,125	1,147	1,104
Mood disorders	1,994	2,158	2,311	2,467	2,576	2,647	2,773	2,603	2,818	2,871	2,789
Personality disorders	406	449	499	546	575	605	621	579	602	616	600
Schizophrenia and other psychotic disorders	810	855	886	903	924	922	937	889	878	855	816
Screening and history of mental health and substance abuse	2,254	2,515	2,724	2,938	3,136	3,199	3,325	3,109	3,464	3,545	3,403
Substance-related disorders	527	581	646	716	780	845	892	839	953	941	930
Suicide and intentional self-inflicted injury	104	119	125	135	151	164	183	174	184	203	201



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Gateway Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,623	6,052	6,350	6,498	6,542	6,617	6,779	6,451	6,655	6,598	6,168
Female	2,904	3,086	3,162	3,208	3,165	3,179	3,237	3,030	3,090	3,026	2,810
Male	2,719	2,966	3,188	3,290	3,377	3,438	3,542	3,421	3,565	3,572	3,358
White	5,217	5,597	5,851	5,967	5,973	6,042	6,149	5,859	6,029	5,966	5,576
Minority	108	127	141	129	134	128	121	104	109	111	102

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	59%	61%	63%	64%	63%	63%	63%	59%	59%	57%	52%
Female	62%	63%	64%	64%	63%	62%	62%	57%	57%	54%	49%
Male	56%	60%	62%	63%	63%	64%	64%	60%	60%	59%	54%
White	58%	60%	62%	62%	62%	61%	61%	57%	57%	55%	50%
Minority	19%	21%	22%	20%	19%	18%	16%	14%	14%	13%	12%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,130	1,202	1,295	1,346	1,355	1,389	1,398	1,257	1,243	1,223	1,122
Alcohol-related disorders	266	256	251	244	228	212	196	153	134	108	86
Anxiety disorders	2,244	2,408	2,528	2,617	2,683	2,701	2,790	2,661	2,702	2,644	2,492
Attention-deficit, conduct, and disruptive behavior disorders	1,976	2,145	2,291	2,434	2,479	2,492	2,605	2,399	2,426	2,360	2,167
Delirium, dementia, and amnestic and other cognitive disorders	69	68	69	63	71	68	70	69	69	62	62
Developmental disorders	1,416	1,589	1,762	1,920	2,043	2,128	2,240	2,210	2,282	2,309	2,225
Disorders usually diagnosed in infancy, childhood, adolescence	478	524	575	656	691	710	753	700	714	690	660
Impulse control disorders, NEC	183	195	209	220	215	204	206	168	162	145	126
Miscellaneous disorders	785	804	820	811	799	785	797	750	722	719	660
Mood disorders	2,173	2,222	2,215	2,201	2,145	2,037	1,998	1,755	1,703	1,560	1,352
Personality disorders	197	188	177	178	166	145	147	120	116	103	83
Schizophrenia and other psychotic disorders	220	205	220	219	218	207	211	186	175	165	141
Screening and history of mental health and substance abuse	4,289	4,524	4,689	4,693	4,592	4,494	4,503	4,114	4,139	3,988	3,642
Substance-related disorders	659	634	633	599	560	506	495	414	367	317	249
Suicide and intentional self-inflicted injury	199	196	194	195	195	186	185	158	151	140	111



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Gateway Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	6,251	6,692	7,013	7,289	7,478	7,675	7,863	7,510	8,029	8,017	7,802
Female	4,262	4,517	4,744	4,938	5,073	5,198	5,370	5,050	5,415	5,352	5,185
Male	1,989	2,175	2,269	2,351	2,405	2,477	2,493	2,460	2,614	2,665	2,617
White	5,658	6,083	6,391	6,643	6,831	7,026	7,209	6,886	7,394	7,398	7,159
Minority	212	214	204	198	181	165	129	101	115	108	109

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	67%	69%	69%	71%	71%	71%	72%	68%	71%	70%	67%
Female	70%	72%	73%	75%	75%	75%	76%	71%	75%	73%	70%
Male	61%	62%	63%	64%	65%	65%	65%	64%	65%	64%	61%
White	66%	68%	69%	70%	71%	71%	72%	68%	71%	70%	66%
Minority	26%	26%	25%	23%	21%	19%	16%	12%	14%	13%	12%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	499	551	607	648	700	748	798	803	858	817	821
Alcohol-related disorders	681	742	790	823	886	879	941	917	946	919	881
Anxiety disorders	3,618	3,931	4,175	4,428	4,644	4,780	4,976	4,781	5,075	5,040	4,933
Attention-deficit, conduct, and disruptive behavior disorders	243	267	300	337	378	419	466	484	553	590	611
Delirium, dementia, and amnesic and other cognitive disorders	1,459	1,465	1,444	1,408	1,362	1,313	1,254	1,164	1,140	1,058	913
Developmental disorders	647	696	740	773	819	834	855	826	853	883	887
Disorders usually diagnosed in infancy, childhood, adolescence	111	124	130	125	141	146	164	159	176	184	188
Impulse control disorders, NEC	107	120	129	133	149	170	166	166	172	187	180
Miscellaneous disorders	1,427	1,556	1,659	1,752	1,846	1,926	2,018	1,892	1,984	1,983	1,931
Mood disorders	3,600	3,915	4,171	4,438	4,668	4,812	5,024	4,780	5,088	5,053	4,928
Personality disorders	682	755	820	859	904	901	918	861	872	851	848
Schizophrenia and other psychotic disorders	1,293	1,332	1,381	1,399	1,394	1,383	1,432	1,351	1,334	1,293	1,215
Screening and history of mental health and substance abuse	3,840	4,245	4,636	4,903	5,073	5,281	5,496	5,318	5,804	5,830	5,713
Substance-related disorders	1,010	1,140	1,233	1,343	1,452	1,545	1,657	1,611	1,738	1,763	1,763
Suicide and intentional self-inflicted injury	212	239	245	266	291	318	361	362	386	383	399



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

FIVCO Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	8,557	9,398	9,639	10,078	10,306	10,329	10,331	9,660	9,968	9,783	9,046
Female	4,177	4,506	4,634	4,776	4,868	4,752	4,746	4,425	4,568	4,434	4,075
Male	4,380	4,892	5,005	5,302	5,438	5,577	5,585	5,235	5,400	5,349	4,971
White	7,966	8,737	8,911	9,272	9,437	9,429	9,367	8,753	8,994	8,810	8,144
Minority	189	197	186	179	178	169	169	150	173	167	161

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	54%	55%	56%	56%	56%	56%	53%	55%	52%	48%
Female	53%	54%	55%	55%	55%	54%	54%	51%	52%	49%	44%
Male	51%	54%	56%	57%	57%	58%	58%	55%	57%	55%	51%
White	51%	53%	54%	55%	55%	54%	54%	51%	53%	51%	46%
Minority	24%	22%	20%	17%	16%	15%	14%	13%	14%	13%	12%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,517	1,661	1,744	1,842	1,891	1,903	1,899	1,753	1,772	1,721	1,523
Alcohol-related disorders	418	393	375	361	334	295	264	223	190	170	127
Anxiety disorders	2,842	3,006	3,037	3,103	3,084	3,040	2,980	2,781	2,778	2,679	2,417
Attention-deficit, conduct, and disruptive behavior disorders	3,801	4,212	4,428	4,747	4,941	5,076	5,105	4,721	4,794	4,625	4,270
Delirium, dementia, and amnestic and other cognitive disorders	124	124	125	127	117	117	116	107	106	95	86
Developmental disorders	2,102	2,422	2,687	2,969	3,202	3,373	3,462	3,390	3,632	3,668	3,491
Disorders usually diagnosed in infancy, childhood, adolescence	695	792	873	955	1,025	1,049	1,096	1,041	1,068	1,038	974
Impulse control disorders, NEC	286	305	302	311	297	279	266	242	220	194	158
Miscellaneous disorders	935	982	995	1,024	1,032	1,018	1,016	950	966	913	830
Mood disorders	3,318	3,464	3,431	3,428	3,307	3,071	2,892	2,550	2,399	2,179	1,877
Personality disorders	303	303	300	278	262	234	214	176	161	145	118
Schizophrenia and other psychotic disorders	335	333	333	324	318	309	286	246	234	211	173
Screening and history of mental health and substance abuse	6,367	6,949	7,068	7,272	7,321	7,169	7,028	6,375	6,390	6,096	5,538
Substance-related disorders	1,119	1,131	1,033	970	891	788	706	618	587	536	456
Suicide and intentional self-inflicted injury	266	267	263	259	245	227	220	210	198	172	146



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

FIVCO Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	9,946	10,916	11,572	12,167	12,529	12,744	13,005	11,991	12,758	12,707	12,184
Female	6,917	7,553	8,015	8,378	8,592	8,774	8,928	8,142	8,631	8,567	8,158
Male	3,029	3,363	3,557	3,789	3,937	3,970	4,077	3,849	4,127	4,140	4,026
White	9,083	10,006	10,644	11,217	11,572	11,772	12,035	11,058	11,773	11,715	11,166
Minority	278	280	264	258	218	179	124	108	134	132	131

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	66%	67%	68%	67%	67%	68%	63%	68%	67%	63%
Female	66%	69%	70%	71%	71%	71%	72%	66%	71%	70%	66%
Male	57%	60%	60%	62%	61%	61%	62%	58%	62%	61%	59%
White	63%	65%	66%	67%	67%	67%	68%	62%	67%	66%	63%
Minority	23%	22%	21%	21%	17%	14%	10%	9%	10%	9%	9%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	679	748	817	894	948	1,015	1,078	1,014	1,097	1,121	1,101
Alcohol-related disorders	1,199	1,339	1,397	1,527	1,601	1,626	1,654	1,544	1,629	1,592	1,511
Anxiety disorders	5,781	6,461	6,924	7,321	7,622	7,840	8,071	7,591	7,949	7,919	7,611
Attention-deficit, conduct, and disruptive behavior disorders	520	602	667	742	804	865	1,003	972	1,085	1,137	1,170
Delirium, dementia, and amnestic and other cognitive disorders	2,187	2,255	2,280	2,282	2,250	2,184	2,152	1,973	1,912	1,779	1,617
Developmental disorders	1,120	1,186	1,254	1,333	1,372	1,397	1,403	1,340	1,395	1,394	1,406
Disorders usually diagnosed in infancy, childhood, adolescence	116	126	136	144	156	172	185	205	226	253	284
Impulse control disorders, NEC	182	208	225	248	270	284	306	309	329	343	337
Miscellaneous disorders	1,938	2,158	2,281	2,407	2,488	2,569	2,628	2,441	2,544	2,476	2,331
Mood disorders	5,990	6,702	7,230	7,630	7,989	8,263	8,503	7,890	8,306	8,294	7,934
Personality disorders	1,372	1,492	1,579	1,672	1,742	1,725	1,706	1,599	1,598	1,583	1,506
Schizophrenia and other psychotic disorders	2,222	2,392	2,487	2,573	2,597	2,556	2,556	2,408	2,419	2,320	2,140
Screening and history of mental health and substance abuse	5,931	6,684	7,279	7,832	8,238	8,407	8,677	8,043	8,658	8,712	8,407
Substance-related disorders	1,815	2,096	2,288	2,519	2,711	2,907	3,085	2,914	3,172	3,161	3,024
Suicide and intentional self-inflicted injury	340	389	416	455	483	526	583	585	634	625	612



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Big Sandy Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	14,331	15,016	15,287	15,661	15,665	15,534	15,369	14,462	14,677	14,482	13,485
Female	7,180	7,391	7,426	7,550	7,510	7,338	7,170	6,718	6,765	6,634	6,154
Male	7,151	7,625	7,861	8,111	8,155	8,196	8,199	7,744	7,912	7,848	7,331
White	13,331	13,919	14,094	14,379	14,317	14,146	13,918	13,040	13,217	12,989	12,107
Minority	142	139	143	134	130	113	103	102	98	94	93

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	56%	58%	59%	59%	59%	59%	56%	57%	56%	52%
Female	57%	58%	59%	59%	59%	58%	58%	54%	55%	54%	50%
Male	53%	55%	58%	59%	59%	60%	61%	58%	60%	59%	54%
White	53%	55%	57%	57%	57%	57%	57%	54%	55%	54%	50%
Minority	11%	11%	10%	9%	8%	7%	6%	6%	6%	5%	5%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,068	2,223	2,266	2,343	2,333	2,310	2,256	2,023	1,968	1,892	1,704
Alcohol-related disorders	593	555	529	489	436	388	358	284	250	214	168
Anxiety disorders	4,914	5,019	5,023	5,169	5,249	5,178	5,170	4,830	4,811	4,651	4,302
Attention-deficit, conduct, and disruptive behavior disorders	4,867	5,323	5,668	5,936	6,055	6,071	6,081	5,644	5,655	5,526	5,150
Delirium, dementia, and amnestic and other cognitive disorders	155	149	153	153	142	141	131	126	120	120	110
Developmental disorders	2,949	3,332	3,656	3,958	4,203	4,395	4,536	4,470	4,681	4,706	4,497
Disorders usually diagnosed in infancy, childhood, adolescence	786	876	939	1,014	1,080	1,139	1,183	1,141	1,155	1,142	1,082
Impulse control disorders, NEC	286	298	298	285	279	268	250	223	201	181	139
Miscellaneous disorders	2,492	2,505	2,480	2,508	2,496	2,453	2,407	2,226	2,198	2,093	1,930
Mood disorders	4,377	4,308	4,103	3,969	3,747	3,465	3,229	2,783	2,573	2,381	2,028
Personality disorders	209	212	209	204	189	182	163	132	119	106	79
Schizophrenia and other psychotic disorders	393	395	394	392	364	353	333	268	255	243	207
Screening and history of mental health and substance abuse	9,542	9,875	9,998	10,087	9,820	9,433	9,072	8,307	8,255	7,925	7,153
Substance-related disorders	1,718	1,602	1,458	1,366	1,235	1,085	967	841	797	741	643
Suicide and intentional self-inflicted injury	356	349	344	339	312	296	265	229	204	192	163



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Big Sandy Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	18,073	19,104	19,989	20,673	21,174	21,382	21,743	20,609	21,996	22,070	21,643
Female	11,932	12,601	13,101	13,526	13,809	14,002	14,249	13,445	14,359	14,318	13,995
Male	6,141	6,503	6,888	7,147	7,365	7,380	7,494	7,164	7,637	7,752	7,648
White	16,651	17,637	18,470	19,127	19,594	19,808	20,127	18,990	20,350	20,445	19,980
Minority	312	300	286	242	199	150	89	67	84	85	95

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	70%	72%	73%	73%	73%	74%	70%	74%	73%	71%
Female	71%	74%	76%	77%	77%	77%	78%	73%	77%	77%	74%
Male	61%	64%	66%	67%	67%	66%	67%	64%	67%	67%	65%
White	67%	69%	71%	72%	72%	72%	73%	69%	73%	73%	70%
Minority	17%	16%	15%	13%	10%	8%	5%	3%	4%	4%	4%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,478	1,561	1,673	1,784	1,872	1,952	2,009	1,953	2,085	2,115	2,082
Alcohol-related disorders	1,846	2,010	2,102	2,215	2,268	2,256	2,346	2,268	2,357	2,341	2,206
Anxiety disorders	12,151	12,941	13,597	14,159	14,560	14,790	15,084	14,465	15,146	15,108	14,711
Attention-deficit, conduct, and disruptive behavior disorders	481	540	589	644	719	814	912	924	1,100	1,250	1,339
Delirium, dementia, and amnesic and other cognitive disorders	2,705	2,747	2,728	2,692	2,650	2,581	2,526	2,338	2,299	2,189	1,977
Developmental disorders	1,104	1,164	1,182	1,239	1,256	1,277	1,325	1,313	1,366	1,398	1,419
Disorders usually diagnosed in infancy, childhood, adolescence	155	168	172	188	201	212	214	217	248	266	267
Impulse control disorders, NEC	191	210	225	246	256	277	293	287	305	313	321
Miscellaneous disorders	5,709	6,199	6,564	6,979	7,251	7,393	7,460	7,083	7,241	7,104	6,750
Mood disorders	11,173	11,976	12,647	13,172	13,527	13,773	14,070	13,476	14,145	14,108	13,754
Personality disorders	810	880	914	970	993	995	1,020	1,006	1,023	1,003	1,000
Schizophrenia and other psychotic disorders	2,800	2,926	2,968	2,974	2,985	2,953	2,931	2,738	2,737	2,657	2,518
Screening and history of mental health and substance abuse	11,908	12,801	13,730	14,492	14,948	15,187	15,650	15,081	16,141	16,250	16,005
Substance-related disorders	4,074	4,402	4,688	5,010	5,290	5,473	5,764	5,670	6,099	6,228	6,162
Suicide and intentional self-inflicted injury	622	674	715	764	819	861	930	933	959	962	955



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

KY River Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	12,240	12,644	12,716	12,913	12,829	12,798	12,727	12,011	12,308	12,101	11,233
Female	6,121	6,227	6,178	6,193	6,040	5,998	5,959	5,536	5,652	5,493	5,101
Male	6,119	6,417	6,538	6,720	6,789	6,800	6,768	6,475	6,656	6,608	6,132
White	11,479	11,840	11,868	12,031	11,901	11,796	11,631	10,932	11,184	10,982	10,200
Minority	166	170	174	166	163	172	168	160	168	159	152

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	59%	61%	62%	62%	63%	64%	61%	63%	60%	55%
Female	60%	61%	62%	62%	62%	62%	63%	59%	61%	57%	53%
Male	55%	58%	60%	61%	62%	63%	65%	62%	65%	62%	57%
White	56%	58%	60%	61%	61%	61%	62%	59%	61%	58%	53%
Minority	18%	17%	17%	16%	15%	15%	14%	12%	13%	11%	11%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,610	2,739	2,837	2,922	2,904	2,928	2,885	2,736	2,737	2,631	2,413
Alcohol-related disorders	844	836	770	761	700	639	574	500	466	421	335
Anxiety disorders	4,847	4,936	4,923	4,998	4,914	4,873	4,785	4,465	4,406	4,220	3,850
Attention-deficit, conduct, and disruptive behavior disorders	4,073	4,287	4,433	4,626	4,659	4,680	4,635	4,361	4,332	4,158	3,801
Delirium, dementia, and amnestic and other cognitive disorders	176	170	167	171	171	164	174	158	150	137	113
Developmental disorders	3,332	3,647	3,915	4,232	4,436	4,559	4,645	4,566	4,723	4,676	4,446
Disorders usually diagnosed in infancy, childhood, adolescence	705	770	818	861	884	906	918	884	913	886	824
Impulse control disorders, NEC	542	568	578	583	565	528	491	448	401	358	294
Miscellaneous disorders	1,877	1,897	1,880	1,852	1,779	1,760	1,710	1,572	1,575	1,534	1,442
Mood disorders	3,945	3,827	3,655	3,488	3,210	3,005	2,758	2,408	2,240	2,043	1,695
Personality disorders	555	547	542	541	519	481	433	387	341	314	249
Schizophrenia and other psychotic disorders	446	434	412	387	368	336	319	285	251	226	185
Screening and history of mental health and substance abuse	8,180	8,369	8,230	8,202	8,009	7,642	7,377	6,760	6,682	6,421	5,783
Substance-related disorders	1,927	1,857	1,703	1,599	1,448	1,332	1,239	1,093	1,042	966	831
Suicide and intentional self-inflicted injury	343	331	292	278	248	218	218	180	153	153	108



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

KY River Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	17,117	18,044	18,766	19,350	19,623	19,570	19,702	18,598	19,566	19,547	18,823
Female	10,895	11,485	11,919	12,182	12,365	12,410	12,466	11,718	12,325	12,301	11,755
Male	6,222	6,559	6,847	7,168	7,258	7,160	7,236	6,880	7,241	7,246	7,068
White	15,855	16,749	17,444	18,030	18,300	18,273	18,391	17,311	18,308	18,299	17,535
Minority	359	348	323	279	248	208	151	135	139	136	135

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	70%	72%	73%	73%	73%	74%	71%	75%	74%	71%
Female	72%	74%	76%	77%	77%	77%	78%	74%	79%	78%	74%
Male	62%	64%	66%	67%	67%	67%	68%	65%	69%	68%	66%
White	68%	70%	72%	72%	72%	73%	74%	70%	74%	74%	71%
Minority	21%	20%	19%	17%	15%	13%	9%	8%	9%	8%	8%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,657	1,780	1,873	2,027	2,108	2,165	2,258	2,164	2,322	2,380	2,347
Alcohol-related disorders	2,132	2,269	2,427	2,572	2,650	2,658	2,669	2,559	2,597	2,588	2,512
Anxiety disorders	11,376	12,110	12,716	13,321	13,627	13,687	13,883	13,318	13,836	13,785	13,310
Attention-deficit, conduct, and disruptive behavior disorders	469	514	579	672	771	852	956	947	1,125	1,230	1,290
Delirium, dementia, and amnesic and other cognitive disorders	3,144	3,198	3,157	3,016	2,911	2,766	2,669	2,453	2,396	2,242	2,060
Developmental disorders	1,547	1,606	1,654	1,745	1,799	1,822	1,918	1,848	1,946	1,968	1,967
Disorders usually diagnosed in infancy, childhood, adolescence	134	146	154	169	182	193	203	214	242	259	283
Impulse control disorders, NEC	364	387	406	443	467	479	493	475	522	542	546
Miscellaneous disorders	4,947	5,388	5,754	6,100	6,352	6,405	6,389	6,088	6,160	6,037	5,762
Mood disorders	10,004	10,624	11,182	11,718	12,029	12,120	12,331	11,816	12,301	12,251	11,928
Personality disorders	1,272	1,338	1,430	1,557	1,661	1,637	1,606	1,499	1,529	1,514	1,482
Schizophrenia and other psychotic disorders	3,527	3,680	3,730	3,765	3,752	3,662	3,624	3,435	3,426	3,339	3,170
Screening and history of mental health and substance abuse	10,738	11,563	12,315	12,873	13,289	13,469	13,670	12,916	13,646	13,612	13,111
Substance-related disorders	4,495	4,864	5,203	5,566	5,802	5,969	6,139	5,895	6,231	6,279	6,152
Suicide and intentional self-inflicted injury	668	710	766	818	866	913	967	941	1,007	1,003	996



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Cumberland Valley Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	22,478	24,210	25,134	26,132	26,603	26,751	26,639	25,318	25,971	25,439	23,734
Female	11,407	12,118	12,380	12,741	12,890	12,879	12,781	12,039	12,310	11,966	11,132
Male	11,071	12,092	12,754	13,391	13,713	13,872	13,858	13,279	13,661	13,473	12,602
White	20,828	22,357	23,118	23,968	24,363	24,416	24,222	22,941	23,509	22,961	21,399
Minority	404	436	448	457	433	432	437	432	445	450	426

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	59%	61%	62%	63%	64%	63%	60%	61%	59%	55%
Female	59%	61%	62%	63%	63%	63%	63%	59%	59%	57%	53%
Male	55%	57%	60%	62%	63%	64%	64%	62%	62%	61%	56%
White	56%	58%	60%	61%	62%	62%	62%	59%	59%	57%	53%
Minority	19%	19%	18%	17%	16%	15%	15%	15%	15%	14%	13%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	3,976	4,221	4,401	4,562	4,559	4,485	4,321	3,943	3,870	3,641	3,241
Alcohol-related disorders	1,739	1,787	1,789	1,757	1,732	1,607	1,443	1,245	1,131	969	819
Anxiety disorders	11,019	11,879	12,442	13,089	13,522	13,709	13,646	13,027	13,165	12,841	12,041
Attention-deficit, conduct, and disruptive behavior disorders	8,445	9,267	9,828	10,312	10,556	10,590	10,400	9,788	9,804	9,424	8,735
Delirium, dementia, and amnestic and other cognitive disorders	332	336	333	322	303	292	276	244	239	229	198
Developmental disorders	6,458	7,184	7,823	8,476	9,028	9,249	9,278	8,857	9,093	8,890	8,387
Disorders usually diagnosed in infancy, childhood, adolescence	1,454	1,652	1,793	1,945	2,095	2,160	2,159	2,049	2,051	1,984	1,880
Impulse control disorders, NEC	844	876	880	878	865	816	776	697	658	597	506
Miscellaneous disorders	3,806	3,898	3,873	3,843	3,778	3,621	3,486	3,237	3,191	3,029	2,746
Mood disorders	7,165	7,176	7,031	6,841	6,504	6,084	5,580	4,893	4,541	4,027	3,433
Personality disorders	1,416	1,430	1,429	1,455	1,440	1,356	1,239	1,093	984	872	748
Schizophrenia and other psychotic disorders	842	842	834	829	789	747	679	617	585	514	433
Screening and history of mental health and substance abuse	11,248	11,737	11,798	11,763	11,428	10,991	10,523	9,663	9,659	9,209	8,264
Substance-related disorders	3,835	3,797	3,708	3,599	3,411	3,088	2,741	2,395	2,181	1,947	1,612
Suicide and intentional self-inflicted injury	693	697	663	643	602	573	529	475	449	396	339



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Cumberland Valley Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	28,250	30,096	31,423	32,488	33,232	33,478	34,263	32,844	34,637	34,885	33,952
Female	18,791	19,906	20,718	21,321	21,787	21,845	22,340	21,402	22,563	22,645	21,878
Male	9,459	10,190	10,705	11,167	11,445	11,633	11,923	11,442	12,074	12,240	12,074
White	25,734	27,485	28,764	29,792	30,560	30,834	31,579	30,241	32,020	32,213	31,269
Minority	830	830	783	748	663	567	469	410	410	452	432

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	70%	71%	72%	72%	72%	73%	69%	72%	72%	69%
Female	72%	74%	75%	76%	75%	75%	76%	72%	76%	75%	72%
Male	63%	64%	65%	66%	66%	66%	67%	64%	67%	66%	64%
White	68%	70%	71%	71%	71%	71%	72%	69%	72%	71%	68%
Minority	25%	24%	23%	22%	20%	17%	14%	13%	13%	13%	12%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,864	3,125	3,334	3,521	3,695	3,806	4,021	3,930	4,172	4,301	4,198
Alcohol-related disorders	3,036	3,268	3,421	3,643	3,808	3,848	3,969	3,746	3,947	3,937	3,827
Anxiety disorders	19,547	20,893	22,040	22,965	23,617	23,945	24,604	23,916	24,926	25,067	24,557
Attention-deficit, conduct, and disruptive behavior disorders	1,175	1,312	1,463	1,660	1,911	2,072	2,271	2,283	2,561	2,801	2,884
Delirium, dementia, and amnesic and other cognitive disorders	5,259	5,363	5,304	5,199	5,024	4,894	4,788	4,443	4,302	4,042	3,661
Developmental disorders	2,356	2,477	2,624	2,777	2,921	3,003	3,114	3,081	3,240	3,384	3,440
Disorders usually diagnosed in infancy, childhood, adolescence	294	322	329	347	358	390	417	421	466	522	551
Impulse control disorders, NEC	500	554	591	632	693	715	741	710	780	808	826
Miscellaneous disorders	8,386	9,016	9,624	10,193	10,680	10,867	11,061	10,798	11,222	11,267	10,901
Mood disorders	16,469	17,654	18,644	19,587	20,185	20,446	21,032	20,388	21,347	21,502	21,050
Personality disorders	2,420	2,621	2,772	2,977	3,170	3,222	3,246	3,131	3,244	3,251	3,189
Schizophrenia and other psychotic disorders	5,307	5,571	5,651	5,736	5,708	5,691	5,735	5,459	5,436	5,242	4,948
Screening and history of mental health and substance abuse	15,583	16,971	18,146	19,097	19,833	20,280	21,070	20,330	21,649	21,923	21,471
Substance-related disorders	6,542	7,114	7,641	8,256	8,755	9,044	9,428	9,101	9,689	9,897	9,692
Suicide and intentional self-inflicted injury	996	1,106	1,190	1,284	1,370	1,457	1,615	1,622	1,722	1,755	1,720



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lake Cumberland Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	14,534	15,796	16,430	17,335	17,616	17,734	17,793	16,869	17,288	16,959	15,751
Female	7,293	7,855	8,084	8,432	8,523	8,512	8,495	7,981	8,197	8,022	7,347
Male	7,241	7,941	8,346	8,903	9,093	9,222	9,298	8,888	9,091	8,937	8,404
White	13,440	14,568	15,100	15,890	16,078	16,155	16,152	15,246	15,595	15,302	14,168
Minority	435	462	474	490	486	471	443	440	462	431	412

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	59%	61%	61%	61%	61%	61%	57%	58%	54%	49%
Female	58%	59%	61%	61%	61%	60%	59%	55%	55%	52%	47%
Male	56%	58%	60%	62%	62%	62%	62%	59%	60%	57%	52%
White	56%	58%	60%	60%	60%	60%	60%	56%	56%	53%	48%
Minority	27%	27%	26%	26%	24%	22%	21%	20%	20%	18%	17%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	3,021	3,212	3,276	3,386	3,376	3,324	3,242	2,931	2,893	2,729	2,406
Alcohol-related disorders	702	693	677	644	567	496	466	380	323	249	183
Anxiety disorders	7,020	7,579	7,832	8,258	8,428	8,513	8,537	8,061	8,080	7,754	7,158
Attention-deficit, conduct, and disruptive behavior disorders	6,690	7,225	7,537	7,894	7,997	7,998	7,890	7,227	7,183	6,909	6,349
Delirium, dementia, and amnestic and other cognitive disorders	226	235	231	230	228	227	215	203	188	170	154
Developmental disorders	2,918	3,328	3,684	4,103	4,387	4,631	4,894	4,855	5,213	5,274	5,080
Disorders usually diagnosed in infancy, childhood, adolescence	1,045	1,170	1,276	1,364	1,431	1,434	1,453	1,369	1,390	1,350	1,287
Impulse control disorders, NEC	911	936	963	971	916	876	829	696	651	583	498
Miscellaneous disorders	2,162	2,263	2,255	2,201	2,157	2,058	1,971	1,753	1,710	1,570	1,378
Mood disorders	5,789	5,915	5,804	5,725	5,455	5,147	4,813	4,140	3,905	3,568	3,075
Personality disorders	458	432	424	404	375	346	313	263	234	212	164
Schizophrenia and other psychotic disorders	584	575	576	564	548	518	497	430	416	378	316
Screening and history of mental health and substance abuse	7,833	8,260	8,395	8,495	8,256	7,897	7,581	6,822	6,746	6,368	5,750
Substance-related disorders	1,880	1,828	1,750	1,654	1,492	1,339	1,198	959	855	750	604
Suicide and intentional self-inflicted injury	511	501	499	474	438	419	401	346	340	313	259



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lake Cumberland Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	19,365	20,483	21,164	21,708	22,137	22,301	22,725	21,625	22,899	23,181	22,209
Female	13,043	13,741	14,103	14,426	14,686	14,833	15,140	14,395	15,270	15,308	14,523
Male	6,322	6,742	7,061	7,282	7,451	7,468	7,585	7,230	7,629	7,873	7,686
White	17,538	18,581	19,253	19,785	20,219	20,426	20,828	19,791	21,041	21,330	20,313
Minority	760	763	738	707	652	569	499	455	497	483	497

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	70%	72%	72%	71%	71%	72%	68%	70%	70%	66%
Female	71%	73%	74%	74%	74%	74%	75%	70%	73%	72%	69%
Male	62%	65%	67%	67%	67%	67%	67%	64%	66%	65%	62%
White	67%	70%	71%	71%	71%	71%	72%	67%	70%	70%	66%
Minority	30%	30%	29%	28%	26%	23%	20%	19%	20%	19%	18%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,925	2,101	2,256	2,420	2,513	2,607	2,696	2,653	2,920	2,996	2,909
Alcohol-related disorders	1,856	2,036	2,120	2,185	2,295	2,270	2,293	2,166	2,250	2,304	2,209
Anxiety disorders	12,309	13,188	13,877	14,463	14,972	15,280	15,762	15,232	16,020	16,295	15,812
Attention-deficit, conduct, and disruptive behavior disorders	820	912	1,032	1,147	1,277	1,391	1,593	1,671	1,933	2,139	2,179
Delirium, dementia, and amnesic and other cognitive disorders	4,471	4,549	4,488	4,313	4,127	3,944	3,785	3,551	3,409	3,176	2,888
Developmental disorders	2,066	2,099	2,172	2,218	2,281	2,330	2,379	2,333	2,393	2,447	2,430
Disorders usually diagnosed in infancy, childhood, adolescence	278	295	321	330	340	356	384	375	413	463	468
Impulse control disorders, NEC	380	416	464	505	544	567	631	623	680	731	748
Miscellaneous disorders	4,439	4,789	5,077	5,368	5,569	5,726	5,872	5,644	5,922	5,952	5,656
Mood disorders	12,132	12,970	13,570	14,112	14,562	14,751	15,167	14,598	15,318	15,502	14,949
Personality disorders	1,627	1,707	1,780	1,825	1,893	1,879	1,893	1,791	1,849	1,834	1,790
Schizophrenia and other psychotic disorders	4,430	4,600	4,701	4,644	4,596	4,458	4,455	4,274	4,229	4,085	3,838
Screening and history of mental health and substance abuse	10,049	10,894	11,544	12,091	12,570	12,755	13,169	12,593	13,455	13,700	13,074
Substance-related disorders	3,234	3,526	3,714	3,952	4,172	4,271	4,451	4,298	4,704	4,905	4,808
Suicide and intentional self-inflicted injury	776	837	897	956	1,009	1,066	1,134	1,117	1,218	1,260	1,245



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Bluegrass Area Development District

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	26,973	30,256	32,251	34,142	35,411	36,195	36,898	35,489	37,749	38,315	36,328
Female	13,483	14,838	15,583	16,213	16,666	16,831	17,046	16,095	17,097	17,259	16,212
Male	13,490	15,418	16,668	17,929	18,745	19,364	19,852	19,394	20,652	21,056	20,116
White	20,925	23,486	24,905	26,357	27,496	28,079	28,509	27,382	29,173	29,666	28,218
Minority	4,641	5,161	5,521	5,769	5,752	5,778	5,857	5,571	5,855	5,871	5,502

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	56%	58%	58%	57%	57%	56%	52%	52%	49%	44%
Female	54%	55%	57%	56%	55%	54%	53%	49%	49%	46%	41%
Male	53%	56%	59%	59%	59%	59%	58%	55%	56%	53%	48%
White	54%	56%	58%	58%	57%	57%	55%	51%	52%	49%	44%
Minority	40%	41%	42%	42%	41%	40%	40%	38%	38%	35%	32%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	7,589	8,537	9,120	9,620	9,863	9,859	9,799	9,064	9,243	8,923	8,182
Alcohol-related disorders	1,216	1,280	1,291	1,261	1,205	1,176	1,114	953	913	819	677
Anxiety disorders	9,843	10,906	11,716	12,474	12,991	13,273	13,549	12,980	13,487	13,424	12,671
Attention-deficit, conduct, and disruptive behavior disorders	12,796	14,354	15,424	16,425	17,002	17,222	17,399	16,419	16,794	16,550	15,503
Delirium, dementia, and amnestic and other cognitive disorders	488	538	560	558	596	613	616	596	598	563	504
Developmental disorders	5,739	6,830	7,909	8,854	9,644	10,403	11,123	11,275	12,359	12,886	12,494
Disorders usually diagnosed in infancy, childhood, adolescence	2,634	3,072	3,487	3,813	4,124	4,336	4,533	4,382	4,593	4,629	4,446
Impulse control disorders, NEC	1,184	1,246	1,306	1,309	1,294	1,223	1,172	1,021	956	844	718
Miscellaneous disorders	4,613	5,045	5,344	5,481	5,621	5,658	5,686	5,380	5,573	5,482	5,099
Mood disorders	9,534	10,046	10,242	10,245	10,112	9,781	9,322	8,390	8,260	7,860	6,924
Personality disorders	935	972	981	940	920	855	793	684	648	586	518
Schizophrenia and other psychotic disorders	1,307	1,357	1,364	1,400	1,400	1,362	1,316	1,214	1,196	1,102	970
Screening and history of mental health and substance abuse	16,363	17,599	18,136	18,306	18,021	17,484	17,027	15,513	15,611	15,063	13,562
Substance-related disorders	3,380	3,491	3,466	3,323	3,166	2,989	2,827	2,474	2,431	2,191	1,813
Suicide and intentional self-inflicted injury	1,092	1,137	1,189	1,190	1,164	1,139	1,134	1,054	1,058	1,030	894



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Bluegrass Area Development District

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	28,027	30,746	32,391	34,080	35,183	36,248	37,172	35,248	38,685	40,105	38,996
Female	20,458	22,438	23,630	24,877	25,693	26,349	26,953	25,255	27,841	28,549	27,418
Male	7,569	8,308	8,761	9,203	9,490	9,899	10,219	9,993	10,844	11,556	11,578
White	22,806	25,125	26,553	28,052	29,024	29,953	30,837	29,250	32,169	33,202	32,110
Minority	3,975	4,283	4,402	4,505	4,560	4,566	4,469	4,142	4,604	4,912	4,701

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	66%	67%	68%	68%	68%	69%	64%	69%	67%	63%
Female	65%	69%	69%	70%	70%	70%	70%	65%	70%	69%	64%
Male	59%	61%	63%	64%	65%	65%	66%	63%	66%	64%	61%
White	65%	68%	69%	70%	70%	70%	71%	66%	70%	68%	64%
Minority	45%	46%	46%	45%	45%	44%	44%	41%	45%	45%	41%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	3,760	4,198	4,605	5,059	5,385	5,667	5,917	5,813	6,470	6,771	6,655
Alcohol-related disorders	2,854	3,155	3,414	3,657	3,802	3,894	4,101	3,923	4,247	4,309	4,164
Anxiety disorders	15,453	17,244	18,609	19,886	20,811	21,616	22,478	21,787	23,577	24,162	23,649
Attention-deficit, conduct, and disruptive behavior disorders	1,682	1,957	2,178	2,496	2,757	3,117	3,432	3,558	4,141	4,611	4,749
Delirium, dementia, and amnesic and other cognitive disorders	7,474	7,653	7,624	7,426	7,290	7,165	6,950	6,369	6,285	6,062	5,573
Developmental disorders	2,931	3,155	3,290	3,470	3,573	3,713	3,875	3,818	4,040	4,230	4,265
Disorders usually diagnosed in infancy, childhood, adolescence	361	421	452	491	529	573	647	702	812	912	969
Impulse control disorders, NEC	419	472	505	559	616	682	759	794	873	946	956
Miscellaneous disorders	7,006	7,915	8,608	9,335	9,923	10,473	10,974	10,294	11,163	11,473	11,041
Mood disorders	16,340	18,188	19,540	20,965	21,835	22,671	23,571	22,666	24,483	25,112	24,609
Personality disorders	2,709	2,975	3,258	3,460	3,564	3,665	3,802	3,671	3,854	3,895	3,783
Schizophrenia and other psychotic disorders	6,822	7,197	7,401	7,424	7,460	7,530	7,646	7,238	7,265	7,146	6,731
Screening and history of mental health and substance abuse	15,597	17,691	19,280	20,885	21,988	22,750	23,654	22,605	25,017	25,983	25,295
Substance-related disorders	5,252	5,916	6,446	7,178	7,733	8,172	8,689	8,377	9,491	9,954	9,738
Suicide and intentional self-inflicted injury	1,193	1,328	1,440	1,594	1,734	1,856	2,057	2,044	2,270	2,357	2,396



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Adair, Edmonson, Garrard, Green, Lincoln, & Russell

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,951	6,572	6,829	7,162	7,368	7,506	7,567	7,178	7,482	7,481	6,963
Female	2,974	3,268	3,333	3,441	3,545	3,569	3,576	3,369	3,519	3,466	3,166
Male	2,977	3,304	3,496	3,721	3,823	3,937	3,991	3,809	3,963	4,015	3,797
White	5,523	6,075	6,310	6,580	6,750	6,850	6,888	6,523	6,798	6,783	6,311
Minority	172	191	194	216	212	222	196	192	200	197	192

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	58%	60%	60%	61%	61%	60%	57%	57%	53%	49%
Female	57%	59%	60%	60%	60%	60%	59%	55%	55%	51%	46%
Male	54%	57%	60%	61%	61%	62%	62%	58%	59%	56%	51%
White	55%	57%	59%	59%	60%	60%	59%	56%	56%	52%	47%
Minority	26%	27%	27%	27%	26%	25%	22%	21%	22%	19%	19%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,515	1,645	1,707	1,784	1,798	1,743	1,731	1,582	1,594	1,561	1,406
Alcohol-related disorders	228	226	221	205	198	177	164	141	127	113	87
Anxiety disorders	2,649	2,898	3,001	3,190	3,301	3,379	3,414	3,199	3,219	3,133	2,874
Attention-deficit, conduct, and disruptive behavior disorders	2,645	2,867	3,036	3,198	3,285	3,314	3,339	3,119	3,145	3,085	2,866
Delirium, dementia, and amnestic and other cognitive disorders	110	116	116	110	116	116	101	100	93	81	73
Developmental disorders	1,155	1,358	1,530	1,720	1,874	2,003	2,129	2,096	2,279	2,339	2,267
Disorders usually diagnosed in infancy, childhood, adolescence	462	509	574	623	662	688	702	696	739	753	717
Impulse control disorders, NEC	385	390	415	416	404	386	366	309	303	273	234
Miscellaneous disorders	1,067	1,162	1,165	1,139	1,149	1,129	1,138	1,022	1,013	979	865
Mood disorders	2,316	2,359	2,311	2,247	2,187	2,077	1,989	1,732	1,655	1,565	1,341
Personality disorders	215	203	209	197	192	177	166	134	124	110	97
Schizophrenia and other psychotic disorders	242	242	242	243	247	240	237	212	206	183	158
Screening and history of mental health and substance abuse	3,419	3,584	3,649	3,670	3,516	3,389	3,263	2,971	3,017	2,944	2,682
Substance-related disorders	670	651	607	590	549	516	455	382	346	317	268
Suicide and intentional self-inflicted injury	176	177	185	180	175	176	166	154	143	135	110



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Adair, Edmonson, Garrard, Green, Lincoln, & Russell

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	7,527	8,096	8,434	8,634	8,716	8,849	9,095	8,486	9,133	9,321	8,906
Female	5,174	5,587	5,773	5,916	5,961	6,068	6,258	5,761	6,264	6,309	5,993
Male	2,353	2,509	2,661	2,718	2,755	2,781	2,837	2,725	2,869	3,012	2,913
White	6,821	7,358	7,677	7,894	7,974	8,108	8,334	7,782	8,409	8,600	8,185
Minority	335	349	328	302	292	256	237	214	237	228	213

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	66%	69%	71%	72%	71%	71%	73%	67%	70%	69%	65%
Female	69%	72%	74%	75%	74%	75%	76%	70%	74%	73%	68%
Male	61%	63%	66%	66%	65%	65%	66%	63%	64%	63%	60%
White	66%	69%	71%	72%	71%	71%	73%	67%	70%	69%	65%
Minority	32%	34%	33%	30%	29%	26%	24%	22%	24%	23%	20%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	912	1,034	1,122	1,207	1,280	1,344	1,411	1,351	1,483	1,522	1,503
Alcohol-related disorders	653	710	766	783	807	822	852	785	816	838	799
Anxiety disorders	4,670	5,082	5,392	5,607	5,793	5,946	6,203	5,876	6,287	6,408	6,189
Attention-deficit, conduct, and disruptive behavior disorders	316	361	411	460	492	540	639	654	774	856	877
Delirium, dementia, and amnesic and other cognitive disorders	1,969	1,988	1,988	1,923	1,828	1,764	1,734	1,586	1,564	1,474	1,324
Developmental disorders	691	709	740	762	777	785	821	815	830	857	821
Disorders usually diagnosed in infancy, childhood, adolescence	50	54	56	61	66	66	83	89	99	113	121
Impulse control disorders, NEC	99	104	118	126	131	147	175	173	194	223	228
Miscellaneous disorders	1,786	1,986	2,129	2,263	2,325	2,411	2,510	2,320	2,494	2,485	2,381
Mood disorders	4,656	5,118	5,391	5,601	5,738	5,901	6,140	5,788	6,176	6,298	6,068
Personality disorders	774	848	883	923	952	956	1,003	935	965	962	933
Schizophrenia and other psychotic disorders	2,035	2,093	2,131	2,078	2,027	2,009	2,011	1,882	1,888	1,831	1,700
Screening and history of mental health and substance abuse	3,694	4,097	4,398	4,604	4,706	4,852	5,090	4,758	5,174	5,290	5,069
Substance-related disorders	1,132	1,259	1,354	1,424	1,504	1,573	1,684	1,580	1,763	1,822	1,781
Suicide and intentional self-inflicted injury	272	302	317	342	356	378	422	417	452	467	462



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Allen, Barren, Butler, Logan, & Simpson

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,144	5,819	6,263	6,584	6,698	6,959	7,078	6,746	6,960	7,152	6,723
Female	2,578	2,901	3,088	3,232	3,251	3,368	3,391	3,201	3,293	3,320	3,105
Male	2,566	2,918	3,175	3,352	3,447	3,591	3,687	3,545	3,667	3,832	3,618
White	4,502	5,107	5,521	5,758	5,865	6,100	6,162	5,886	6,050	6,224	5,850
Minority	455	500	503	549	512	506	497	463	478	476	465

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	51%	53%	54%	53%	52%	52%	52%	49%	50%	47%	43%
Female	52%	53%	54%	53%	51%	50%	50%	47%	48%	45%	40%
Male	51%	52%	54%	54%	53%	53%	54%	51%	52%	50%	45%
White	51%	52%	54%	53%	52%	51%	51%	49%	49%	47%	42%
Minority	36%	38%	37%	38%	34%	32%	32%	29%	29%	27%	26%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	992	1,124	1,184	1,226	1,239	1,261	1,233	1,125	1,151	1,135	1,016
Alcohol-related disorders	182	191	185	176	168	147	135	115	111	100	71
Anxiety disorders	1,724	1,894	2,073	2,150	2,259	2,378	2,452	2,330	2,392	2,356	2,185
Attention-deficit, conduct, and disruptive behavior disorders	2,403	2,708	2,917	3,077	3,189	3,307	3,329	3,100	3,141	3,158	2,895
Delirium, dementia, and amnestic and other cognitive disorders	64	67	77	75	82	79	81	77	81	80	76
Developmental disorders	984	1,172	1,352	1,518	1,659	1,826	1,995	1,956	2,112	2,228	2,172
Disorders usually diagnosed in infancy, childhood, adolescence	402	456	521	579	634	681	712	664	691	692	678
Impulse control disorders, NEC	334	362	386	380	373	348	339	288	273	256	213
Miscellaneous disorders	832	898	935	971	973	980	965	891	855	821	734
Mood disorders	1,879	1,987	2,085	2,073	2,000	1,908	1,814	1,618	1,536	1,426	1,217
Personality disorders	102	108	111	104	108	99	93	81	72	58	38
Schizophrenia and other psychotic disorders	205	208	216	225	218	201	191	165	163	142	123
Screening and history of mental health and substance abuse	3,315	3,696	3,797	3,783	3,583	3,454	3,332	2,994	3,015	2,952	2,650
Substance-related disorders	487	505	495	480	442	418	381	325	319	266	202
Suicide and intentional self-inflicted injury	150	154	161	142	143	128	122	108	96	86	63



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Allen, Barren, Butler, Logan, & Simpson

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	6,487	7,068	7,483	7,635	7,846	8,002	8,226	7,687	8,083	8,231	7,999
Female	4,588	4,989	5,267	5,389	5,522	5,656	5,785	5,347	5,633	5,730	5,535
Male	1,899	2,079	2,216	2,246	2,324	2,346	2,441	2,340	2,450	2,501	2,464
White	5,574	6,107	6,495	6,622	6,835	6,977	7,167	6,697	7,059	7,183	6,976
Minority	597	628	625	629	617	608	604	538	579	602	559

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	68%	68%	68%	68%	69%	65%	68%	66%	62%
Female	66%	69%	70%	70%	70%	70%	71%	66%	70%	68%	64%
Male	60%	63%	63%	64%	65%	64%	65%	62%	63%	61%	58%
White	64%	67%	68%	68%	68%	68%	69%	65%	68%	66%	62%
Minority	41%	42%	42%	42%	40%	39%	39%	36%	39%	36%	33%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	482	552	598	647	674	716	792	757	804	823	835
Alcohol-related disorders	556	623	654	668	708	710	733	695	709	733	707
Anxiety disorders	3,587	3,951	4,269	4,440	4,632	4,778	4,985	4,790	4,999	5,107	4,987
Attention-deficit, conduct, and disruptive behavior disorders	286	332	375	423	490	532	635	631	707	797	841
Delirium, dementia, and amnesic and other cognitive disorders	1,962	2,021	2,001	1,894	1,803	1,767	1,684	1,559	1,529	1,427	1,288
Developmental disorders	641	686	705	732	734	753	792	778	799	826	803
Disorders usually diagnosed in infancy, childhood, adolescence	37	47	48	52	61	73	84	97	109	127	133
Impulse control disorders, NEC	83	94	104	126	135	148	170	160	177	198	204
Miscellaneous disorders	1,360	1,555	1,718	1,820	1,914	2,007	2,159	2,016	2,108	2,137	2,072
Mood disorders	3,859	4,265	4,606	4,821	5,015	5,126	5,335	5,077	5,356	5,426	5,236
Personality disorders	286	316	349	373	382	396	412	389	392	415	398
Schizophrenia and other psychotic disorders	1,598	1,695	1,768	1,786	1,769	1,784	1,795	1,723	1,701	1,640	1,526
Screening and history of mental health and substance abuse	3,216	3,643	4,012	4,170	4,396	4,477	4,655	4,365	4,591	4,709	4,610
Substance-related disorders	745	866	955	1,021	1,078	1,075	1,137	1,089	1,142	1,192	1,158
Suicide and intentional self-inflicted injury	178	206	240	255	285	301	342	331	362	380	400



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Anderson, Clark, Jessamin, Madison, Washington, & Woodford

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	8,265	9,251	9,859	10,529	10,993	11,238	11,686	11,262	12,076	12,231	11,591
Female	4,147	4,551	4,764	5,015	5,212	5,264	5,428	5,149	5,516	5,514	5,198
Male	4,118	4,700	5,095	5,514	5,781	5,974	6,258	6,113	6,560	6,717	6,393
White	7,181	8,029	8,535	9,104	9,502	9,697	10,025	9,623	10,311	10,483	9,927
Minority	696	771	814	856	870	871	902	874	943	914	867

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	56%	59%	59%	59%	59%	59%	55%	56%	52%	48%
Female	55%	56%	58%	58%	57%	56%	56%	52%	53%	49%	44%
Male	54%	57%	60%	61%	60%	61%	62%	59%	59%	56%	51%
White	54%	56%	59%	59%	58%	58%	58%	54%	54%	51%	47%
Minority	36%	38%	38%	37%	36%	35%	36%	34%	36%	31%	29%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,365	2,653	2,840	2,997	3,058	3,047	3,108	2,871	2,895	2,766	2,524
Alcohol-related disorders	380	382	398	384	370	351	348	299	286	251	220
Anxiety disorders	3,268	3,593	3,872	4,141	4,279	4,352	4,579	4,425	4,619	4,620	4,341
Attention-deficit, conduct, and disruptive behavior disorders	3,769	4,204	4,520	4,826	5,026	5,070	5,219	4,907	5,058	4,964	4,634
Delirium, dementia, and amnestic and other cognitive disorders	146	152	159	155	166	177	195	191	193	178	159
Developmental disorders	2,014	2,390	2,760	3,092	3,405	3,668	3,977	3,990	4,388	4,557	4,424
Disorders usually diagnosed in infancy, childhood, adolescence	838	1,005	1,152	1,269	1,402	1,479	1,580	1,507	1,600	1,635	1,581
Impulse control disorders, NEC	342	360	378	382	377	359	349	295	259	240	198
Miscellaneous disorders	1,416	1,563	1,685	1,765	1,811	1,831	1,889	1,825	1,894	1,879	1,756
Mood disorders	2,876	3,000	3,096	3,078	3,007	2,904	2,845	2,575	2,554	2,428	2,123
Personality disorders	282	299	310	298	285	265	248	216	215	190	167
Schizophrenia and other psychotic disorders	403	411	428	444	451	450	441	411	398	370	328
Screening and history of mental health and substance abuse	4,857	5,196	5,372	5,462	5,381	5,231	5,154	4,703	4,770	4,530	4,077
Substance-related disorders	1,023	1,030	1,035	1,022	960	880	866	772	750	684	564
Suicide and intentional self-inflicted injury	317	322	338	333	320	333	339	327	332	328	299



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Anderson, Clark, Jessamin, Madison, Washington, & Woodford

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	7,894	8,763	9,198	9,761	10,205	10,443	11,030	10,508	11,634	12,126	11,799
Female	5,837	6,437	6,799	7,246	7,581	7,729	8,126	7,626	8,463	8,721	8,383
Male	2,057	2,326	2,399	2,515	2,624	2,714	2,904	2,882	3,171	3,405	3,416
White	6,853	7,601	8,023	8,550	8,940	9,163	9,710	9,275	10,313	10,740	10,389
Minority	676	763	755	762	783	775	767	676	739	802	777

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	67%	67%	69%	69%	69%	71%	66%	70%	69%	65%
Female	65%	69%	69%	71%	71%	71%	72%	67%	71%	70%	66%
Male	58%	62%	62%	63%	64%	66%	67%	64%	67%	66%	63%
White	64%	67%	68%	70%	70%	70%	71%	66%	71%	69%	66%
Minority	39%	41%	40%	40%	40%	39%	39%	35%	37%	37%	34%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,334	1,485	1,643	1,826	1,918	2,015	2,159	2,091	2,357	2,479	2,420
Alcohol-related disorders	711	801	877	960	1,003	1,022	1,117	1,050	1,145	1,141	1,096
Anxiety disorders	4,477	5,075	5,478	5,936	6,225	6,466	6,902	6,660	7,290	7,518	7,408
Attention-deficit, conduct, and disruptive behavior disorders	465	549	606	703	786	904	1,035	1,040	1,249	1,415	1,457
Delirium, dementia, and amnestic and other cognitive disorders	2,094	2,165	2,139	2,069	2,075	2,053	2,039	1,901	1,867	1,787	1,655
Developmental disorders	837	933	950	996	1,045	1,078	1,156	1,152	1,241	1,316	1,333
Disorders usually diagnosed in infancy, childhood, adolescence	120	137	149	164	174	185	211	222	274	313	316
Impulse control disorders, NEC	122	143	151	163	171	180	210	214	253	275	285
Miscellaneous disorders	2,009	2,265	2,507	2,709	2,943	3,079	3,339	3,132	3,415	3,545	3,420
Mood disorders	4,728	5,321	5,700	6,150	6,468	6,683	7,154	6,809	7,459	7,733	7,595
Personality disorders	667	728	813	870	918	950	1,036	971	1,026	1,057	1,011
Schizophrenia and other psychotic disorders	1,796	1,911	1,973	1,991	2,019	2,048	2,159	2,018	2,059	2,039	1,921
Screening and history of mental health and substance abuse	4,278	4,907	5,343	5,900	6,259	6,472	7,001	6,712	7,500	7,838	7,677
Substance-related disorders	1,431	1,633	1,809	2,039	2,195	2,313	2,553	2,436	2,790	2,926	2,854
Suicide and intentional self-inflicted injury	271	312	367	401	457	469	541	514	575	611	621



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Ballard, Fulton, Graves, & Hickman

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,860	3,117	3,339	3,572	3,628	3,645	3,717	3,392	3,505	3,426	3,213
Female	1,351	1,454	1,539	1,629	1,627	1,609	1,670	1,494	1,532	1,502	1,381
Male	1,509	1,663	1,800	1,943	2,001	2,036	2,047	1,898	1,973	1,924	1,832
White	2,091	2,254	2,422	2,616	2,649	2,676	2,721	2,470	2,551	2,487	2,363
Minority	632	679	694	704	686	663	655	583	585	561	490

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	51%	53%	54%	54%	53%	52%	52%	48%	48%	46%	42%
Female	49%	51%	51%	51%	49%	48%	49%	44%	44%	42%	37%
Male	52%	55%	56%	57%	57%	56%	55%	52%	53%	50%	46%
White	49%	51%	52%	52%	51%	49%	49%	45%	45%	42%	39%
Minority	45%	47%	46%	45%	43%	42%	41%	37%	37%	35%	31%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	323	361	389	379	384	381	391	352	348	336	306
Alcohol-related disorders	80	78	73	70	67	56	57	49	47	44	33
Anxiety disorders	613	641	654	676	684	644	636	550	539	517	462
Attention-deficit, conduct, and disruptive behavior disorders	1,485	1,644	1,732	1,896	1,951	1,966	1,973	1,806	1,848	1,783	1,678
Delirium, dementia, and amnestic and other cognitive disorders	35	41	39	38	45	44	48	49	49	48	48
Developmental disorders	695	833	955	1,076	1,197	1,300	1,397	1,378	1,474	1,503	1,432
Disorders usually diagnosed in infancy, childhood, adolescence	196	227	255	286	307	324	330	310	324	320	297
Impulse control disorders, NEC	177	195	196	202	200	196	188	170	173	156	129
Miscellaneous disorders	367	396	416	446	459	435	441	407	420	386	357
Mood disorders	951	981	1,019	1,052	1,030	979	951	830	825	754	659
Personality disorders	61	61	60	66	63	61	58	46	41	43	35
Schizophrenia and other psychotic disorders	82	91	82	87	84	75	69	64	67	68	62
Screening and history of mental health and substance abuse	1,563	1,681	1,736	1,797	1,706	1,616	1,585	1,373	1,391	1,314	1,161
Substance-related disorders	180	180	172	188	182	170	158	130	141	119	93
Suicide and intentional self-inflicted injury	62	71	71	71	68	64	62	54	58	54	46



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Ballard, Fulton, Graves, & Hickman

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,406	3,697	3,940	4,056	4,178	4,226	4,273	3,955	4,210	4,189	4,064
Female	2,516	2,719	2,864	2,970	3,044	3,075	3,072	2,793	2,988	2,948	2,848
Male	890	978	1,076	1,086	1,134	1,151	1,201	1,162	1,222	1,241	1,216
White	2,684	2,917	3,129	3,228	3,320	3,382	3,419	3,171	3,395	3,372	3,257
Minority	602	653	678	690	709	679	668	597	623	623	605

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	61%	64%	66%	66%	66%	66%	66%	62%	65%	65%	62%
Female	63%	67%	68%	69%	69%	68%	68%	63%	67%	67%	63%
Male	54%	57%	60%	60%	60%	60%	61%	59%	61%	61%	58%
White	61%	64%	66%	67%	67%	66%	66%	62%	66%	65%	62%
Minority	50%	53%	54%	54%	53%	52%	51%	46%	48%	49%	46%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	242	267	289	321	330	347	371	356	367	370	361
Alcohol-related disorders	294	326	364	368	384	391	396	377	400	391	364
Anxiety disorders	1,740	1,893	2,018	2,139	2,229	2,292	2,351	2,217	2,331	2,363	2,287
Attention-deficit, conduct, and disruptive behavior disorders	201	219	248	272	312	332	385	389	425	468	474
Delirium, dementia, and amnesic and other cognitive disorders	1,002	1,031	1,045	994	975	947	934	864	873	790	723
Developmental disorders	337	366	376	382	384	389	420	408	426	416	420
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33	36	39	41	49
Impulse control disorders, NEC	65	71	83	82	91	97	109	100	104	112	118
Miscellaneous disorders	627	707	774	814	865	884	927	889	913	935	900
Mood disorders	2,061	2,244	2,397	2,505	2,619	2,710	2,766	2,580	2,730	2,688	2,652
Personality disorders	188	192	219	220	230	229	257	247	273	265	253
Schizophrenia and other psychotic disorders	801	837	871	893	900	894	909	859	871	831	791
Screening and history of mental health and substance abuse	1,600	1,819	2,039	2,162	2,241	2,252	2,304	2,159	2,305	2,380	2,334
Substance-related disorders	431	493	553	549	591	619	646	620	642	642	599
Suicide and intentional self-inflicted injury	147	164	174	181	199	208	230	235	255	264	252



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Bath, Elliott, Meniffee, & Morgan

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,433	3,686	3,790	3,910	3,911	3,939	3,962	3,736	3,830	3,770	3,490
Female	1,778	1,902	1,897	1,950	1,919	1,933	1,937	1,784	1,819	1,765	1,643
Male	1,655	1,784	1,893	1,960	1,992	2,006	2,025	1,952	2,011	2,005	1,847
White	3,211	3,439	3,524	3,629	3,623	3,646	3,639	3,443	3,526	3,460	3,202
Minority	38	48	48	43	43	39	31	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	62%	63%	64%	64%	64%	64%	60%	61%	59%	54%
Female	64%	64%	65%	65%	64%	64%	64%	60%	60%	57%	53%
Male	57%	59%	62%	63%	63%	64%	64%	61%	61%	61%	55%
White	59%	60%	62%	62%	62%	63%	62%	59%	59%	58%	52%
Minority	14%	16%	16%	14%	13%	12%	9%	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	605	637	665	689	685	704	708	642	626	617	562
Alcohol-related disorders	163	158	153	148	140	136	131	101	83	65	51
Anxiety disorders	1,349	1,438	1,465	1,538	1,545	1,570	1,605	1,497	1,510	1,447	1,349
Attention-deficit, conduct, and disruptive behavior disorders	1,075	1,162	1,222	1,314	1,329	1,327	1,391	1,248	1,269	1,235	1,113
Delirium, dementia, and amnestic and other cognitive disorders	42	39	41	37	39	39	40	39	42	38	36
Developmental disorders	905	1,013	1,112	1,224	1,300	1,340	1,378	1,328	1,379	1,381	1,312
Disorders usually diagnosed in infancy, childhood, adolescence	241	259	280	324	336	346	367	320	337	331	316
Impulse control disorders, NEC	98	102	107	116	110	99	96	73	69	57	51
Miscellaneous disorders	479	481	475	477	449	427	428	392	378	371	329
Mood disorders	1,217	1,231	1,199	1,182	1,132	1,066	1,034	898	868	785	665
Personality disorders	96	90	85	84	75	67	70	54	54	43	35
Schizophrenia and other psychotic disorders	122	115	123	116	110	104	105	91	82	78	61
Screening and history of mental health and substance abuse	2,567	2,716	2,747	2,773	2,708	2,660	2,621	2,387	2,385	2,282	2,075
Substance-related disorders	363	357	343	323	298	272	257	211	181	161	131
Suicide and intentional self-inflicted injury	114	110	107	103	101	101	94	79	72	62	47



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Bath, Elliott, Meniffee, & Morgan

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,795	4,073	4,299	4,489	4,572	4,703	4,784	4,535	4,771	4,818	4,668
Female	2,514	2,682	2,832	2,950	3,004	3,095	3,169	2,966	3,116	3,120	3,020
Male	1,281	1,391	1,467	1,539	1,568	1,608	1,615	1,569	1,655	1,698	1,648
White	3,453	3,725	3,949	4,140	4,220	4,344	4,415	4,189	4,432	4,480	4,310
Minority	103	105	96	86	69	65	46	31	33	34	41

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	67%	68%	69%	70%	70%	72%	68%	71%	71%	67%
Female	69%	71%	72%	74%	74%	75%	77%	72%	75%	75%	72%
Male	58%	59%	60%	61%	62%	62%	63%	62%	64%	64%	60%
White	64%	66%	67%	68%	69%	69%	71%	67%	70%	70%	67%
Minority	23%	23%	21%	19%	15%	14%	10%	7%	8%	8%	9%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	250	276	303	331	342	374	398	401	421	415	402
Alcohol-related disorders	359	394	424	460	496	502	538	530	533	529	504
Anxiety disorders	2,197	2,377	2,573	2,720	2,851	2,947	3,054	2,899	3,043	3,053	2,952
Attention-deficit, conduct, and disruptive behavior disorders	135	148	166	188	201	226	246	253	296	296	305
Delirium, dementia, and amnesic and other cognitive disorders	877	892	882	859	834	806	763	706	682	635	566
Developmental disorders	301	321	345	357	373	376	378	365	397	405	409
Disorders usually diagnosed in infancy, childhood, adolescence	42	45	50	47	61	59	63	61	69	67	76
Impulse control disorders, NEC	N.D.	33	38	41	48	56	54	58	63	66	60
Miscellaneous disorders	824	916	978	1,036	1,093	1,121	1,163	1,083	1,117	1,124	1,102
Mood disorders	2,109	2,299	2,494	2,676	2,788	2,878	2,986	2,809	2,957	2,975	2,877
Personality disorders	401	420	457	490	514	512	507	478	473	480	457
Schizophrenia and other psychotic disorders	744	788	821	846	839	817	832	780	769	756	701
Screening and history of mental health and substance abuse	2,324	2,538	2,793	3,007	3,081	3,221	3,355	3,210	3,432	3,478	3,383
Substance-related disorders	529	593	671	754	811	874	950	903	961	983	970
Suicide and intentional self-inflicted injury	112	127	131	146	158	175	199	202	212	216	220



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Bell & Harlan

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	6,172	6,480	6,666	6,855	6,825	6,786	6,719	6,369	6,429	6,227	5,803
Female	3,199	3,282	3,291	3,335	3,300	3,293	3,249	3,079	3,078	2,931	2,727
Male	2,973	3,198	3,375	3,520	3,525	3,493	3,470	3,290	3,351	3,296	3,076
White	5,696	5,946	6,075	6,218	6,171	6,107	6,015	5,675	5,708	5,501	5,109
Minority	207	216	227	226	204	209	205	205	201	206	204

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	55%	57%	59%	59%	60%	61%	58%	59%	57%	53%
Female	56%	57%	58%	59%	59%	60%	60%	57%	57%	55%	51%
Male	51%	53%	56%	59%	59%	60%	61%	59%	60%	59%	55%
White	53%	54%	56%	58%	58%	58%	59%	56%	57%	55%	52%
Minority	30%	29%	29%	27%	23%	24%	23%	23%	22%	21%	21%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,008	1,057	1,110	1,136	1,100	1,097	1,076	1,002	995	931	829
Alcohol-related disorders	440	427	423	409	388	364	311	264	229	185	155
Anxiety disorders	2,820	2,936	3,028	3,162	3,181	3,198	3,129	2,974	2,956	2,872	2,638
Attention-deficit, conduct, and disruptive behavior disorders	2,150	2,343	2,463	2,589	2,599	2,573	2,568	2,385	2,371	2,268	2,122
Delirium, dementia, and amnestic and other cognitive disorders	68	65	67	67	61	59	54	44	44	39	33
Developmental disorders	2,012	2,185	2,378	2,576	2,714	2,740	2,757	2,628	2,642	2,532	2,388
Disorders usually diagnosed in infancy, childhood, adolescence	309	343	372	415	437	457	462	446	436	424	398
Impulse control disorders, NEC	178	188	189	188	175	170	150	135	125	107	93
Miscellaneous disorders	813	815	799	774	740	688	658	621	614	579	542
Mood disorders	1,739	1,717	1,686	1,639	1,524	1,421	1,297	1,121	1,036	909	781
Personality disorders	402	403	403	409	400	374	329	283	259	215	181
Schizophrenia and other psychotic disorders	245	237	240	231	215	202	184	163	149	136	108
Screening and history of mental health and substance abuse	2,864	2,923	2,938	2,866	2,688	2,553	2,373	2,168	2,138	2,001	1,805
Substance-related disorders	1,108	1,069	1,046	1,016	937	862	755	663	589	518	440
Suicide and intentional self-inflicted injury	219	212	198	190	185	178	160	133	125	105	83



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Bell & Harlan

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	8,067	8,557	8,805	9,041	9,266	9,249	9,339	8,779	9,046	9,048	8,799
Female	5,348	5,666	5,837	5,962	6,120	6,106	6,165	5,794	5,969	5,949	5,777
Male	2,719	2,891	2,968	3,079	3,146	3,143	3,174	2,985	3,077	3,099	3,022
White	7,293	7,754	7,985	8,212	8,448	8,459	8,544	8,037	8,306	8,286	8,041
Minority	354	357	354	339	317	273	242	212	213	223	210

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	66%	68%	68%	69%	69%	69%	70%	67%	69%	69%	66%
Female	69%	72%	73%	73%	73%	73%	74%	70%	73%	73%	70%
Male	60%	62%	61%	62%	62%	62%	63%	61%	63%	62%	60%
White	65%	68%	68%	68%	68%	68%	69%	66%	69%	69%	66%
Minority	32%	32%	32%	31%	29%	26%	24%	21%	22%	21%	20%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	737	795	842	870	916	930	963	907	913	942	918
Alcohol-related disorders	877	955	963	1,010	1,050	1,043	1,072	986	1,043	1,029	985
Anxiety disorders	5,538	5,890	6,135	6,320	6,476	6,495	6,621	6,306	6,452	6,467	6,348
Attention-deficit, conduct, and disruptive behavior disorders	317	348	380	428	493	523	598	584	632	695	708
Delirium, dementia, and amnesic and other cognitive disorders	1,401	1,426	1,416	1,374	1,314	1,285	1,251	1,165	1,129	1,075	968
Developmental disorders	669	708	758	821	877	884	900	883	915	963	953
Disorders usually diagnosed in infancy, childhood, adolescence	75	85	85	83	85	89	96	89	99	114	107
Impulse control disorders, NEC	93	105	114	123	140	140	159	142	161	168	167
Miscellaneous disorders	2,016	2,147	2,266	2,375	2,498	2,508	2,522	2,381	2,402	2,389	2,319
Mood disorders	4,411	4,720	4,985	5,231	5,390	5,394	5,499	5,203	5,352	5,377	5,264
Personality disorders	796	851	914	968	1,040	1,033	1,039	971	976	986	956
Schizophrenia and other psychotic disorders	1,407	1,477	1,492	1,512	1,509	1,474	1,490	1,402	1,390	1,349	1,290
Screening and history of mental health and substance abuse	4,162	4,533	4,798	4,999	5,207	5,283	5,404	5,089	5,262	5,259	5,177
Substance-related disorders	1,728	1,881	1,980	2,136	2,266	2,324	2,395	2,211	2,330	2,358	2,305
Suicide and intentional self-inflicted injury	330	352	379	402	431	458	495	498	515	531	538



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Boone & Campbell

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,548	5,138	5,541	5,877	6,168	6,566	6,931	6,587	7,206	7,459	7,146
Female	2,207	2,434	2,598	2,756	2,853	3,018	3,190	2,989	3,264	3,291	3,153
Male	2,341	2,704	2,943	3,121	3,315	3,548	3,741	3,598	3,942	4,168	3,993
White	3,937	4,427	4,750	5,033	5,288	5,621	5,898	5,596	6,109	6,321	6,059
Minority	369	416	450	462	456	460	482	486	528	549	539

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	55%	57%	57%	55%	55%	56%	51%	52%	48%	43%
Female	53%	54%	56%	55%	53%	53%	53%	48%	49%	44%	40%
Male	54%	56%	58%	58%	57%	58%	58%	54%	55%	52%	46%
White	53%	55%	57%	57%	56%	55%	55%	51%	52%	48%	42%
Minority	32%	32%	33%	30%	29%	28%	28%	26%	28%	24%	23%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	904	1,005	1,054	1,111	1,118	1,186	1,215	1,118	1,176	1,170	1,052
Alcohol-related disorders	234	251	252	241	248	229	207	176	175	155	126
Anxiety disorders	1,285	1,388	1,488	1,518	1,522	1,559	1,576	1,407	1,444	1,398	1,272
Attention-deficit, conduct, and disruptive behavior disorders	2,424	2,775	3,090	3,364	3,631	3,900	4,156	3,929	4,195	4,236	4,015
Delirium, dementia, and amnestic and other cognitive disorders	108	114	129	126	129	139	137	119	135	133	112
Developmental disorders	1,269	1,500	1,693	1,868	2,040	2,219	2,441	2,391	2,677	2,822	2,738
Disorders usually diagnosed in infancy, childhood, adolescence	487	581	672	738	780	834	900	878	925	937	883
Impulse control disorders, NEC	225	238	263	266	245	246	265	226	213	210	185
Miscellaneous disorders	868	946	984	1,006	1,029	1,017	1,042	947	980	977	900
Mood disorders	1,700	1,836	1,913	1,912	1,888	1,877	1,851	1,603	1,630	1,547	1,375
Personality disorders	210	224	221	200	186	169	161	126	117	102	81
Schizophrenia and other psychotic disorders	200	216	228	226	224	216	215	197	200	191	166
Screening and history of mental health and substance abuse	2,784	3,021	3,149	3,168	3,110	3,112	3,125	2,813	2,953	2,900	2,633
Substance-related disorders	463	492	500	466	465	448	412	373	387	355	299
Suicide and intentional self-inflicted injury	177	193	194	189	194	202	199	185	197	182	167



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Boone & Campbell

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,503	4,940	5,400	5,697	5,946	6,242	6,648	5,985	6,664	6,969	6,692
Female	3,417	3,745	4,096	4,313	4,525	4,709	4,989	4,505	5,004	5,214	4,929
Male	1,086	1,195	1,304	1,384	1,421	1,533	1,659	1,480	1,660	1,755	1,763
White	4,083	4,475	4,932	5,193	5,432	5,709	6,104	5,492	6,111	6,373	6,117
Minority	248	294	280	308	305	308	293	248	292	320	282

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	66%	68%	69%	70%	70%	72%	64%	69%	68%	63%
Female	65%	68%	70%	71%	72%	72%	73%	65%	71%	69%	64%
Male	60%	60%	63%	64%	64%	64%	68%	61%	66%	65%	61%
White	64%	67%	69%	70%	71%	71%	73%	65%	70%	69%	65%
Minority	35%	38%	35%	36%	35%	35%	33%	27%	32%	30%	25%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	555	619	677	725	766	816	898	826	908	953	894
Alcohol-related disorders	601	680	753	795	833	887	935	828	913	948	886
Anxiety disorders	2,543	2,814	3,092	3,254	3,416	3,653	3,911	3,611	3,980	4,136	3,964
Attention-deficit, conduct, and disruptive behavior disorders	367	429	526	612	681	760	867	848	1,000	1,083	1,069
Delirium, dementia, and amnestic and other cognitive disorders	1,162	1,232	1,261	1,265	1,250	1,239	1,242	1,136	1,174	1,136	1,052
Developmental disorders	410	452	502	524	545	590	638	634	673	724	721
Disorders usually diagnosed in infancy, childhood, adolescence	63	73	79	92	99	115	136	139	159	181	188
Impulse control disorders, NEC	81	97	108	117	127	142	162	171	193	210	218
Miscellaneous disorders	958	1,094	1,280	1,422	1,540	1,670	1,791	1,542	1,692	1,717	1,563
Mood disorders	2,914	3,235	3,587	3,800	4,016	4,238	4,545	4,193	4,591	4,745	4,582
Personality disorders	452	503	538	585	613	664	726	681	712	709	663
Schizophrenia and other psychotic disorders	1,055	1,155	1,240	1,276	1,285	1,341	1,376	1,283	1,356	1,331	1,230
Screening and history of mental health and substance abuse	2,611	2,902	3,304	3,577	3,779	4,057	4,374	3,955	4,430	4,653	4,468
Substance-related disorders	834	951	1,087	1,180	1,247	1,341	1,481	1,365	1,524	1,614	1,524
Suicide and intentional self-inflicted injury	238	268	308	323	346	389	452	440	498	530	516



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Bourbon, Bracken, Harrison, Nicholas, Pendleton, & Robertson

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,436	3,803	4,130	4,228	4,348	4,451	4,579	4,374	4,657	4,734	4,434
Female	1,668	1,824	1,951	1,978	2,016	2,039	2,090	1,982	2,071	2,099	1,960
Male	1,768	1,979	2,179	2,250	2,332	2,412	2,489	2,392	2,586	2,635	2,474
White	3,044	3,359	3,640	3,725	3,813	3,890	3,984	3,812	4,062	4,144	3,872
Minority	227	239	256	254	254	263	265	250	267	249	233

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	59%	61%	60%	59%	60%	59%	55%	56%	53%	48%
Female	58%	59%	61%	59%	58%	58%	58%	52%	53%	50%	45%
Male	56%	58%	61%	61%	60%	61%	61%	57%	59%	55%	51%
White	56%	57%	60%	59%	58%	58%	57%	53%	55%	51%	46%
Minority	38%	37%	38%	37%	34%	33%	34%	31%	32%	28%	27%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	805	915	984	998	1,033	1,020	1,035	945	984	929	812
Alcohol-related disorders	188	196	205	195	190	191	190	170	167	141	113
Anxiety disorders	1,169	1,292	1,428	1,467	1,536	1,566	1,637	1,575	1,645	1,610	1,516
Attention-deficit, conduct, and disruptive behavior disorders	1,512	1,699	1,893	1,947	2,037	2,071	2,133	1,986	2,093	2,079	1,930
Delirium, dementia, and amnestic and other cognitive disorders	47	57	60	56	60	60	64	60	64	65	63
Developmental disorders	680	803	958	1,033	1,159	1,227	1,319	1,308	1,408	1,480	1,437
Disorders usually diagnosed in infancy, childhood, adolescence	254	310	351	376	412	406	426	397	406	414	377
Impulse control disorders, NEC	151	166	187	188	178	172	169	152	152	126	103
Miscellaneous disorders	616	667	706	711	730	718	700	669	686	666	610
Mood disorders	1,217	1,299	1,377	1,338	1,311	1,248	1,201	1,062	1,077	1,019	851
Personality disorders	155	154	156	148	134	131	128	116	103	92	74
Schizophrenia and other psychotic disorders	153	169	170	157	156	143	142	132	121	121	113
Screening and history of mental health and substance abuse	2,279	2,460	2,582	2,533	2,492	2,445	2,416	2,198	2,250	2,190	1,950
Substance-related disorders	481	498	506	473	456	461	438	398	405	340	260
Suicide and intentional self-inflicted injury	131	134	135	137	129	120	127	122	130	124	104



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Bourbon, Bracken, Harrison, Nicholas, Pendleton, & Robertson

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,410	3,791	3,978	4,153	4,317	4,426	4,654	4,335	4,827	5,031	4,884
Female	2,475	2,750	2,900	3,048	3,145	3,197	3,365	3,096	3,455	3,593	3,452
Male	935	1,041	1,078	1,105	1,172	1,229	1,289	1,239	1,372	1,438	1,432
White	3,060	3,432	3,600	3,753	3,932	4,026	4,245	3,963	4,429	4,608	4,455
Minority	197	195	204	206	189	190	180	158	176	194	184

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	68%	68%	69%	69%	69%	71%	64%	69%	68%	64%
Female	67%	71%	71%	72%	73%	72%	74%	65%	72%	72%	67%
Male	58%	62%	61%	61%	61%	62%	65%	60%	62%	62%	58%
White	64%	68%	68%	69%	70%	70%	71%	64%	69%	69%	64%
Minority	37%	36%	34%	34%	31%	30%	30%	25%	28%	30%	28%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	281	316	348	384	422	448	497	469	529	590	601
Alcohol-related disorders	301	338	367	418	430	457	498	473	525	525	501
Anxiety disorders	1,763	2,005	2,130	2,257	2,401	2,471	2,634	2,509	2,750	2,855	2,783
Attention-deficit, conduct, and disruptive behavior disorders	214	236	270	299	335	389	427	433	484	526	524
Delirium, dementia, and amnesic and other cognitive disorders	919	955	951	936	945	952	955	867	871	828	777
Developmental disorders	332	361	363	366	382	421	452	412	443	463	453
Disorders usually diagnosed in infancy, childhood, adolescence	38	43	43	46	48	56	65	63	80	87	94
Impulse control disorders, NEC	47	50	54	55	59	70	80	75	84	91	92
Miscellaneous disorders	656	732	805	887	954	1,026	1,124	1,024	1,144	1,194	1,165
Mood disorders	1,867	2,087	2,224	2,382	2,509	2,634	2,844	2,690	2,971	3,077	3,023
Personality disorders	321	361	400	421	464	482	506	465	505	500	498
Schizophrenia and other psychotic disorders	794	841	866	875	883	906	941	892	903	870	836
Screening and history of mental health and substance abuse	1,907	2,196	2,409	2,609	2,748	2,814	2,988	2,798	3,167	3,336	3,210
Substance-related disorders	505	582	656	721	809	898	982	936	1,095	1,137	1,125
Suicide and intentional self-inflicted injury	157	177	192	208	235	248	283	261	304	315	315



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Boyd

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,772	3,026	3,102	3,230	3,324	3,367	3,409	3,132	3,283	3,225	3,032
Female	1,334	1,413	1,455	1,517	1,536	1,520	1,533	1,427	1,507	1,463	1,376
Male	1,438	1,613	1,647	1,713	1,788	1,847	1,876	1,705	1,776	1,762	1,656
White	2,514	2,748	2,811	2,902	2,971	3,004	3,016	2,776	2,894	2,844	2,678
Minority	134	134	123	121	122	113	114	101	115	112	108

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	56%	58%	58%	58%	57%	59%	55%	56%	54%	49%
Female	54%	55%	56%	56%	55%	53%	55%	51%	52%	49%	45%
Male	54%	58%	59%	61%	61%	61%	63%	58%	60%	58%	53%
White	53%	55%	57%	57%	56%	56%	57%	53%	54%	52%	48%
Minority	35%	33%	30%	27%	25%	23%	24%	21%	23%	20%	19%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	559	604	642	675	697	714	732	679	695	677	613
Alcohol-related disorders	164	150	147	147	135	121	111	89	77	74	59
Anxiety disorders	957	992	1,006	1,029	1,026	1,030	1,015	909	915	883	804
Attention-deficit, conduct, and disruptive behavior disorders	1,486	1,630	1,696	1,805	1,892	1,948	1,964	1,809	1,846	1,738	1,627
Delirium, dementia, and amnestic and other cognitive disorders	37	33	36	34	33	31	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	622	727	831	943	1,058	1,142	1,210	1,173	1,274	1,289	1,237
Disorders usually diagnosed in infancy, childhood, adolescence	268	307	345	372	407	419	444	419	420	400	381
Impulse control disorders, NEC	125	134	137	141	131	125	126	109	97	82	63
Miscellaneous disorders	285	301	299	309	314	318	342	335	343	323	306
Mood disorders	1,217	1,277	1,269	1,282	1,247	1,175	1,136	987	932	825	746
Personality disorders	125	124	122	112	109	100	94	72	67	63	50
Schizophrenia and other psychotic disorders	130	129	126	123	129	133	124	98	89	82	72
Screening and history of mental health and substance abuse	2,062	2,224	2,265	2,303	2,335	2,311	2,286	2,034	2,077	1,982	1,835
Substance-related disorders	421	411	386	352	320	302	272	226	220	204	182
Suicide and intentional self-inflicted injury	108	111	107	102	100	98	98	85	79	72	65



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Boyd

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,120	3,405	3,654	3,819	3,979	4,083	4,326	3,985	4,245	4,258	4,151
Female	2,234	2,437	2,630	2,713	2,823	2,901	3,043	2,771	2,962	2,964	2,878
Male	886	968	1,024	1,106	1,156	1,182	1,283	1,214	1,283	1,294	1,273
White	2,822	3,085	3,326	3,472	3,628	3,724	3,973	3,649	3,884	3,893	3,770
Minority	133	141	134	139	121	100	81	77	97	91	91

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	67%	68%	69%	68%	68%	70%	65%	70%	68%	65%
Female	67%	69%	71%	72%	71%	71%	73%	68%	72%	71%	68%
Male	59%	60%	62%	63%	62%	63%	65%	60%	65%	62%	61%
White	64%	66%	68%	69%	68%	68%	70%	65%	69%	68%	65%
Minority	32%	31%	30%	31%	26%	21%	18%	16%	20%	17%	15%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	257	288	317	344	375	401	436	403	437	446	445
Alcohol-related disorders	459	509	510	571	594	623	653	599	617	604	584
Anxiety disorders	1,918	2,138	2,289	2,421	2,525	2,621	2,767	2,611	2,715	2,696	2,635
Attention-deficit, conduct, and disruptive behavior disorders	193	223	265	290	314	345	414	403	446	484	498
Delirium, dementia, and amnesic and other cognitive disorders	564	586	625	642	639	625	676	589	586	571	538
Developmental disorders	347	370	403	437	457	459	487	454	489	507	526
Disorders usually diagnosed in infancy, childhood, adolescence	45	46	50	55	60	68	72	83	90	104	118
Impulse control disorders, NEC	65	68	79	92	101	103	116	122	130	147	154
Miscellaneous disorders	590	648	694	741	762	784	838	771	810	788	754
Mood disorders	2,065	2,291	2,502	2,617	2,783	2,893	3,051	2,837	2,943	2,946	2,854
Personality disorders	520	563	607	646	675	669	684	641	638	653	641
Schizophrenia and other psychotic disorders	727	788	854	905	935	925	955	898	910	863	819
Screening and history of mental health and substance abuse	1,917	2,101	2,317	2,457	2,607	2,681	2,861	2,673	2,857	2,900	2,880
Substance-related disorders	707	804	861	951	1,013	1,073	1,169	1,117	1,207	1,227	1,181
Suicide and intentional self-inflicted injury	119	135	152	168	194	209	243	239	261	278	270



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Boyle, Breckinridge, Hancock, Larue, Marion, & Mercer

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,824	5,405	5,806	6,229	6,454	6,543	6,630	6,225	6,554	6,617	6,294
Female	2,362	2,665	2,825	2,981	2,981	3,035	3,094	2,879	3,016	3,021	2,861
Male	2,462	2,740	2,981	3,248	3,473	3,508	3,536	3,346	3,538	3,596	3,433
White	4,102	4,583	4,917	5,291	5,483	5,553	5,548	5,202	5,500	5,550	5,308
Minority	463	508	524	543	550	551	579	519	531	533	498

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	55%	57%	58%	58%	58%	58%	54%	54%	50%	46%
Female	53%	55%	57%	57%	56%	56%	55%	51%	50%	46%	42%
Male	53%	55%	58%	59%	60%	61%	60%	57%	58%	54%	49%
White	52%	54%	56%	57%	57%	57%	56%	52%	52%	48%	44%
Minority	38%	39%	39%	38%	37%	36%	37%	33%	32%	30%	29%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,425	1,614	1,749	1,837	1,857	1,860	1,841	1,703	1,755	1,721	1,615
Alcohol-related disorders	187	181	176	187	188	162	145	114	111	95	72
Anxiety disorders	1,696	1,903	2,044	2,214	2,284	2,352	2,372	2,232	2,346	2,298	2,162
Attention-deficit, conduct, and disruptive behavior disorders	2,300	2,590	2,799	3,040	3,196	3,198	3,222	3,008	3,083	3,036	2,886
Delirium, dementia, and amnestic and other cognitive disorders	69	71	70	68	68	68	67	58	62	65	58
Developmental disorders	1,233	1,437	1,655	1,862	2,008	2,107	2,202	2,174	2,282	2,342	2,262
Disorders usually diagnosed in infancy, childhood, adolescence	395	435	498	555	597	621	652	639	662	666	634
Impulse control disorders, NEC	306	334	341	344	338	321	324	289	271	237	218
Miscellaneous disorders	740	808	862	902	903	892	904	838	861	839	757
Mood disorders	1,714	1,815	1,835	1,882	1,822	1,744	1,683	1,482	1,429	1,362	1,207
Personality disorders	206	226	224	217	215	196	188	163	159	142	120
Schizophrenia and other psychotic disorders	176	178	176	176	171	164	158	137	136	122	112
Screening and history of mental health and substance abuse	2,797	3,045	3,145	3,242	3,160	3,021	2,916	2,621	2,575	2,473	2,281
Substance-related disorders	445	454	454	461	433	382	347	276	260	221	181
Suicide and intentional self-inflicted injury	156	163	161	155	155	146	154	138	141	128	111



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Boyle, Breckinridge, Hancock, Larue, Marion, & Mercer

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,337	5,761	6,158	6,635	6,868	6,956	7,125	6,763	7,275	7,383	7,082
Female	3,809	4,108	4,391	4,737	4,918	4,923	5,029	4,756	5,103	5,164	4,914
Male	1,528	1,653	1,767	1,898	1,950	2,033	2,096	2,007	2,172	2,219	2,168
White	4,630	5,002	5,376	5,831	6,028	6,122	6,294	5,954	6,396	6,489	6,170
Minority	487	518	525	546	564	540	503	472	534	552	541

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	62%	64%	66%	67%	66%	68%	63%	66%	66%	62%
Female	63%	65%	67%	69%	70%	69%	70%	65%	69%	68%	65%
Male	55%	57%	59%	61%	61%	61%	62%	59%	62%	61%	57%
White	60%	63%	65%	67%	67%	67%	68%	63%	66%	66%	62%
Minority	41%	42%	42%	43%	44%	41%	39%	37%	41%	42%	39%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	620	683	781	842	907	940	1,001	921	1,041	1,107	1,060
Alcohol-related disorders	481	539	575	627	672	679	697	651	689	682	653
Anxiety disorders	2,906	3,208	3,534	3,846	4,015	4,126	4,296	4,100	4,380	4,458	4,265
Attention-deficit, conduct, and disruptive behavior disorders	302	343	389	447	507	557	627	609	711	770	775
Delirium, dementia, and amnestic and other cognitive disorders	1,454	1,470	1,471	1,476	1,438	1,401	1,375	1,277	1,251	1,210	1,089
Developmental disorders	576	588	617	657	674	682	716	705	750	779	797
Disorders usually diagnosed in infancy, childhood, adolescence	63	65	67	77	89	101	111	121	124	134	132
Impulse control disorders, NEC	105	110	131	139	155	165	180	175	203	212	212
Miscellaneous disorders	1,124	1,233	1,391	1,546	1,617	1,654	1,719	1,592	1,679	1,732	1,650
Mood disorders	3,160	3,480	3,800	4,094	4,265	4,390	4,588	4,373	4,657	4,689	4,492
Personality disorders	642	689	776	837	848	849	886	802	843	853	831
Schizophrenia and other psychotic disorders	1,370	1,449	1,506	1,551	1,551	1,554	1,555	1,454	1,464	1,411	1,322
Screening and history of mental health and substance abuse	2,527	2,836	3,164	3,538	3,767	3,796	3,889	3,641	3,972	4,009	3,832
Substance-related disorders	721	813	909	1,010	1,082	1,135	1,176	1,060	1,187	1,230	1,161
Suicide and intentional self-inflicted injury	209	224	250	276	298	318	350	331	372	401	400



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Breathitt, Owsley, Perry, & Wolfe

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	6,318	6,533	6,558	6,678	6,660	6,649	6,656	6,286	6,457	6,351	5,860
Female	3,231	3,285	3,248	3,259	3,184	3,154	3,153	2,926	2,995	2,920	2,681
Male	3,087	3,248	3,310	3,419	3,476	3,495	3,503	3,360	3,462	3,431	3,179
White	5,867	6,054	6,042	6,146	6,103	6,053	5,988	5,642	5,786	5,684	5,246
Minority	107	116	120	112	109	113	113	109	113	107	95

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	59%	61%	63%	63%	64%	64%	65%	62%	63%	61%	56%
Female	63%	63%	64%	64%	63%	63%	63%	60%	61%	59%	54%
Male	57%	59%	62%	63%	64%	65%	66%	63%	65%	63%	57%
White	58%	60%	61%	62%	62%	62%	63%	60%	61%	59%	54%
Minority	20%	20%	19%	17%	16%	16%	15%	15%	15%	13%	12%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,253	1,324	1,393	1,447	1,462	1,501	1,503	1,419	1,437	1,381	1,268
Alcohol-related disorders	516	508	467	472	437	406	372	327	312	281	222
Anxiety disorders	2,623	2,696	2,682	2,747	2,714	2,678	2,639	2,458	2,418	2,318	2,092
Attention-deficit, conduct, and disruptive behavior disorders	2,219	2,335	2,412	2,547	2,562	2,578	2,577	2,447	2,440	2,344	2,141
Delirium, dementia, and amnestic and other cognitive disorders	96	91	93	97	94	87	97	101	96	83	68
Developmental disorders	1,709	1,878	2,028	2,186	2,279	2,342	2,417	2,381	2,469	2,441	2,323
Disorders usually diagnosed in infancy, childhood, adolescence	416	448	491	515	535	542	552	529	549	536	493
Impulse control disorders, NEC	348	366	379	389	374	354	336	302	281	255	210
Miscellaneous disorders	901	918	914	893	868	850	831	762	772	755	703
Mood disorders	2,075	2,022	1,939	1,872	1,739	1,651	1,523	1,339	1,279	1,173	970
Personality disorders	282	279	282	290	275	261	240	218	199	190	152
Schizophrenia and other psychotic disorders	250	244	229	212	198	183	172	154	146	136	107
Screening and history of mental health and substance abuse	4,273	4,368	4,278	4,299	4,220	4,072	3,972	3,636	3,627	3,502	3,159
Substance-related disorders	1,023	1,005	922	884	805	765	715	636	616	574	490
Suicide and intentional self-inflicted injury	184	184	166	155	138	125	122	101	93	97	62



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Breathitt, Owsley, Perry, & Wolfe

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	9,102	9,614	10,017	10,237	10,292	10,243	10,407	9,815	10,375	10,385	9,990
Female	5,742	6,046	6,288	6,367	6,392	6,405	6,529	6,094	6,455	6,470	6,167
Male	3,360	3,568	3,729	3,870	3,900	3,838	3,878	3,721	3,920	3,915	3,823
White	8,412	8,891	9,279	9,497	9,561	9,525	9,690	9,106	9,679	9,698	9,290
Minority	212	219	203	177	156	137	95	90	91	88	89

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	70%	72%	74%	75%	74%	74%	76%	72%	76%	75%	72%
Female	74%	76%	79%	79%	79%	79%	81%	76%	80%	80%	76%
Male	63%	65%	68%	69%	68%	68%	70%	67%	70%	69%	66%
White	69%	71%	74%	74%	74%	74%	76%	72%	76%	75%	71%
Minority	23%	24%	22%	19%	17%	15%	11%	10%	11%	10%	10%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	827	895	946	1,010	1,036	1,060	1,129	1,065	1,134	1,186	1,180
Alcohol-related disorders	1,208	1,284	1,379	1,455	1,482	1,500	1,514	1,430	1,468	1,463	1,422
Anxiety disorders	6,137	6,514	6,884	7,126	7,241	7,288	7,441	7,116	7,446	7,449	7,188
Attention-deficit, conduct, and disruptive behavior disorders	264	298	337	383	434	465	546	531	637	691	723
Delirium, dementia, and amnesic and other cognitive disorders	1,756	1,767	1,746	1,660	1,605	1,496	1,461	1,356	1,330	1,248	1,137
Developmental disorders	869	915	952	1,006	1,025	1,040	1,115	1,063	1,133	1,137	1,131
Disorders usually diagnosed in infancy, childhood, adolescence	87	95	100	111	115	121	127	133	153	160	172
Impulse control disorders, NEC	215	226	239	261	273	283	296	282	310	329	325
Miscellaneous disorders	2,523	2,742	2,947	3,101	3,187	3,215	3,250	3,074	3,121	3,067	2,911
Mood disorders	5,324	5,653	5,981	6,211	6,318	6,366	6,557	6,256	6,539	6,542	6,366
Personality disorders	604	640	691	736	776	765	774	708	722	721	710
Schizophrenia and other psychotic disorders	1,795	1,872	1,913	1,929	1,924	1,880	1,891	1,768	1,785	1,763	1,654
Screening and history of mental health and substance abuse	5,730	6,194	6,604	6,837	6,985	7,081	7,241	6,814	7,256	7,255	6,964
Substance-related disorders	2,432	2,620	2,829	3,005	3,100	3,196	3,316	3,138	3,298	3,346	3,275
Suicide and intentional self-inflicted injury	356	378	421	433	462	477	527	499	541	538	539



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Bullitt, Meade, Nelson, Oldham, Spencer, Shelby, & Scott

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	7,852	8,977	9,782	10,627	11,129	11,458	11,729	11,285	12,037	12,392	11,801
Female	3,835	4,351	4,697	5,087	5,254	5,336	5,423	5,164	5,476	5,583	5,251
Male	4,017	4,626	5,085	5,540	5,875	6,122	6,306	6,121	6,561	6,809	6,550
White	6,826	7,801	8,467	9,211	9,598	9,900	10,072	9,650	10,299	10,629	10,125
Minority	639	732	805	832	876	867	879	854	912	914	874

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	49%	51%	53%	54%	53%	53%	52%	48%	49%	46%	41%
Female	48%	50%	52%	52%	51%	50%	49%	45%	46%	42%	38%
Male	49%	51%	54%	55%	55%	55%	55%	51%	53%	49%	45%
White	49%	50%	52%	53%	52%	52%	51%	47%	48%	45%	40%
Minority	30%	32%	35%	34%	34%	32%	32%	30%	30%	27%	25%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,107	2,394	2,676	2,891	2,978	3,034	3,034	2,820	2,943	2,936	2,733
Alcohol-related disorders	331	354	365	399	372	358	322	268	257	228	191
Anxiety disorders	2,469	2,827	3,012	3,249	3,397	3,474	3,565	3,372	3,498	3,471	3,238
Attention-deficit, conduct, and disruptive behavior disorders	3,990	4,516	4,963	5,480	5,805	5,906	5,997	5,672	5,903	5,979	5,646
Delirium, dementia, and amnestic and other cognitive disorders	130	140	142	155	159	163	181	173	174	176	167
Developmental disorders	1,435	1,765	2,100	2,393	2,657	2,896	3,130	3,212	3,487	3,698	3,606
Disorders usually diagnosed in infancy, childhood, adolescence	790	938	1,072	1,186	1,307	1,390	1,503	1,435	1,512	1,532	1,489
Impulse control disorders, NEC	576	615	652	670	674	649	632	567	563	515	441
Miscellaneous disorders	1,025	1,155	1,214	1,306	1,368	1,372	1,434	1,372	1,429	1,425	1,319
Mood disorders	3,048	3,338	3,479	3,624	3,620	3,516	3,370	3,041	3,056	2,944	2,644
Personality disorders	270	294	299	308	305	294	268	247	240	230	211
Schizophrenia and other psychotic disorders	331	359	368	397	392	399	402	362	376	354	305
Screening and history of mental health and substance abuse	4,519	5,012	5,259	5,417	5,353	5,182	5,028	4,565	4,605	4,456	4,065
Substance-related disorders	814	875	884	921	870	841	774	660	633	607	511
Suicide and intentional self-inflicted injury	272	303	322	327	318	327	324	303	317	319	287



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Bullitt, Meade, Nelson, Oldham, Spencer, Shelby, & Scott

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	6,355	6,926	7,622	8,199	8,889	9,336	9,872	9,216	10,307	10,582	10,134
Female	4,724	5,141	5,668	6,141	6,652	6,996	7,356	6,796	7,619	7,759	7,342
Male	1,631	1,785	1,954	2,058	2,237	2,340	2,516	2,420	2,688	2,823	2,792
White	5,517	6,009	6,651	7,156	7,758	8,171	8,657	8,148	9,093	9,389	8,929
Minority	557	605	646	692	757	778	775	633	760	725	696

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	59%	61%	64%	65%	66%	67%	61%	66%	65%	60%
Female	59%	61%	63%	66%	67%	68%	68%	62%	68%	67%	61%
Male	53%	54%	56%	59%	61%	60%	62%	59%	63%	61%	57%
White	58%	60%	62%	65%	66%	66%	67%	62%	67%	66%	60%
Minority	36%	35%	37%	39%	42%	42%	42%	34%	40%	38%	35%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	619	704	811	841	921	1,034	1,124	1,077	1,169	1,240	1,226
Alcohol-related disorders	590	620	715	772	848	891	1,002	912	971	985	954
Anxiety disorders	3,148	3,510	3,938	4,243	4,635	4,918	5,346	5,156	5,587	5,732	5,520
Attention-deficit, conduct, and disruptive behavior disorders	416	481	552	625	707	780	930	949	1,136	1,262	1,326
Delirium, dementia, and amnesic and other cognitive disorders	1,663	1,706	1,735	1,703	1,677	1,670	1,703	1,598	1,592	1,564	1,471
Developmental disorders	658	713	783	791	820	882	950	924	996	1,030	1,062
Disorders usually diagnosed in infancy, childhood, adolescence	126	135	150	157	177	195	208	215	241	272	305
Impulse control disorders, NEC	142	163	194	207	237	242	298	304	336	379	376
Miscellaneous disorders	1,076	1,235	1,390	1,559	1,776	1,946	2,052	1,872	2,088	2,157	2,035
Mood disorders	3,610	4,052	4,535	4,872	5,261	5,552	5,969	5,716	6,239	6,424	6,215
Personality disorders	525	566	625	652	693	710	772	749	766	767	741
Schizophrenia and other psychotic disorders	1,522	1,616	1,702	1,707	1,735	1,739	1,822	1,728	1,740	1,734	1,629
Screening and history of mental health and substance abuse	3,378	3,847	4,357	4,885	5,509	5,842	6,258	5,756	6,589	6,700	6,338
Substance-related disorders	908	1,045	1,177	1,322	1,454	1,567	1,731	1,620	1,828	1,921	1,868
Suicide and intentional self-inflicted injury	229	249	302	331	372	401	470	466	528	558	566



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Calloway, Carlisle, Marshall, & Trigg

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,809	3,183	3,450	3,646	3,770	3,846	3,837	3,574	3,756	3,815	3,546
Female	1,377	1,554	1,664	1,708	1,749	1,782	1,782	1,644	1,704	1,728	1,599
Male	1,432	1,629	1,786	1,938	2,021	2,064	2,055	1,930	2,052	2,087	1,947
White	2,500	2,825	3,052	3,221	3,318	3,357	3,332	3,124	3,271	3,319	3,083
Minority	180	197	208	214	215	226	226	189	207	213	196

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	48%	51%	53%	53%	53%	52%	51%	47%	48%	46%	41%
Female	48%	52%	53%	51%	51%	50%	49%	45%	45%	42%	38%
Male	48%	51%	53%	55%	54%	54%	53%	50%	52%	50%	45%
White	48%	51%	52%	52%	52%	51%	50%	46%	47%	45%	40%
Minority	30%	31%	31%	31%	29%	28%	27%	23%	26%	24%	23%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	465	519	556	570	569	555	543	498	493	471	440
Alcohol-related disorders	77	70	73	76	77	71	66	52	42	41	N.D.
Anxiety disorders	764	827	845	883	884	888	853	780	781	721	624
Attention-deficit, conduct, and disruptive behavior disorders	1,425	1,649	1,753	1,891	1,985	2,013	2,008	1,875	1,922	1,923	1,803
Delirium, dementia, and amnestic and other cognitive disorders	34	41	45	48	40	41	42	38	43	41	39
Developmental disorders	425	513	612	708	790	879	947	958	1,062	1,134	1,084
Disorders usually diagnosed in infancy, childhood, adolescence	185	211	249	273	295	325	331	310	324	326	310
Impulse control disorders, NEC	221	234	250	259	250	247	236	201	188	177	146
Miscellaneous disorders	468	517	539	557	570	580	558	503	488	458	403
Mood disorders	975	1,072	1,110	1,108	1,108	1,062	1,019	904	901	851	745
Personality disorders	96	101	96	91	83	75	79	73	75	66	55
Schizophrenia and other psychotic disorders	94	93	94	90	90	93	92	76	76	74	61
Screening and history of mental health and substance abuse	1,430	1,574	1,699	1,741	1,742	1,724	1,672	1,525	1,566	1,564	1,395
Substance-related disorders	199	204	202	203	210	184	173	150	124	135	108
Suicide and intentional self-inflicted injury	68	70	70	66	71	64	61	57	57	60	48



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Calloway, Carlisle, Marshall, & Trigg

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,093	3,356	3,476	3,621	3,756	3,874	4,020	3,677	4,093	4,211	4,025
Female	2,353	2,528	2,621	2,712	2,806	2,865	2,966	2,695	2,989	3,029	2,858
Male	740	828	855	909	950	1,009	1,054	982	1,104	1,182	1,167
White	2,795	3,023	3,139	3,289	3,412	3,525	3,639	3,326	3,703	3,806	3,610
Minority	191	213	206	197	200	188	201	171	198	196	191

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	59%	63%	64%	63%	64%	64%	65%	59%	64%	64%	59%
Female	61%	65%	66%	66%	66%	66%	68%	62%	66%	65%	60%
Male	54%	57%	56%	58%	58%	58%	58%	54%	59%	59%	56%
White	60%	63%	64%	64%	64%	64%	65%	59%	64%	64%	59%
Minority	34%	38%	36%	34%	34%	31%	33%	29%	32%	29%	27%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	187	204	216	235	254	286	312	303	330	361	336
Alcohol-related disorders	160	184	187	192	233	245	273	242	272	282	279
Anxiety disorders	1,563	1,700	1,790	1,902	2,040	2,141	2,266	2,100	2,330	2,444	2,360
Attention-deficit, conduct, and disruptive behavior disorders	143	159	183	229	272	302	345	348	421	466	480
Delirium, dementia, and amnesic and other cognitive disorders	906	924	927	919	856	820	823	741	776	740	663
Developmental disorders	243	264	274	287	293	302	327	319	338	354	354
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	31	37	39	40	50	63	66
Impulse control disorders, NEC	45	51	55	65	80	83	94	100	121	132	133
Miscellaneous disorders	536	604	665	725	780	803	870	774	862	875	816
Mood disorders	1,711	1,880	2,009	2,183	2,324	2,386	2,523	2,337	2,576	2,706	2,595
Personality disorders	164	210	224	234	249	246	259	249	268	284	258
Schizophrenia and other psychotic disorders	666	731	754	764	755	743	771	746	765	748	680
Screening and history of mental health and substance abuse	1,195	1,415	1,497	1,619	1,740	1,824	1,921	1,764	1,963	2,031	1,943
Substance-related disorders	296	341	372	401	439	475	522	464	524	551	520
Suicide and intentional self-inflicted injury	92	99	113	130	145	157	202	206	231	258	254



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Carroll, Franklin, Gallatin, Grant, Henry, Owen, & Trimble

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,250	6,045	6,503	6,928	7,161	7,308	7,414	7,097	7,512	7,644	7,204
Female	2,546	2,902	3,120	3,264	3,368	3,405	3,391	3,229	3,401	3,403	3,206
Male	2,704	3,143	3,383	3,664	3,793	3,903	4,023	3,868	4,111	4,241	3,998
White	4,647	5,357	5,734	6,077	6,254	6,376	6,427	6,158	6,497	6,624	6,242
Minority	349	375	413	432	448	448	440	394	427	434	412

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	54%	57%	57%	57%	55%	55%	50%	51%	48%	43%
Female	53%	54%	56%	56%	55%	53%	52%	47%	47%	45%	40%
Male	52%	55%	57%	59%	58%	57%	57%	53%	54%	51%	46%
White	52%	54%	56%	57%	56%	54%	53%	49%	49%	47%	42%
Minority	31%	30%	31%	30%	30%	29%	28%	25%	26%	24%	23%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,144	1,324	1,392	1,464	1,503	1,463	1,491	1,382	1,412	1,397	1,271
Alcohol-related disorders	199	220	217	211	199	192	169	140	132	130	105
Anxiety disorders	1,859	2,100	2,228	2,395	2,425	2,389	2,361	2,183	2,223	2,183	1,999
Attention-deficit, conduct, and disruptive behavior disorders	2,646	3,085	3,361	3,602	3,802	3,957	4,019	3,818	3,972	3,973	3,776
Delirium, dementia, and amnesic and other cognitive disorders	86	93	100	98	97	95	101	104	113	108	93
Developmental disorders	880	1,081	1,258	1,444	1,551	1,682	1,840	1,841	2,015	2,093	2,036
Disorders usually diagnosed in infancy, childhood, adolescence	481	562	622	709	742	778	817	782	820	812	775
Impulse control disorders, NEC	289	317	324	326	330	319	309	256	250	236	213
Miscellaneous disorders	804	918	948	972	981	965	966	855	880	863	796
Mood disorders	2,061	2,232	2,334	2,364	2,332	2,288	2,190	1,957	1,965	1,855	1,647
Personality disorders	184	194	192	177	155	146	138	122	111	106	94
Schizophrenia and other psychotic disorders	230	240	240	246	242	245	250	213	214	190	170
Screening and history of mental health and substance abuse	3,240	3,551	3,709	3,780	3,705	3,587	3,445	3,079	3,089	2,968	2,633
Substance-related disorders	539	557	574	535	516	485	461	389	391	347	275
Suicide and intentional self-inflicted injury	197	218	221	212	216	217	219	201	211	208	174



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Carroll, Franklin, Gallatin, Grant, Henry, Owen, & Trimble

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,231	5,719	6,194	6,428	6,700	7,060	7,367	6,888	7,724	8,063	7,685
Female	3,696	4,010	4,364	4,571	4,773	5,026	5,242	4,851	5,488	5,622	5,278
Male	1,535	1,709	1,830	1,857	1,927	2,034	2,125	2,037	2,236	2,441	2,407
White	4,733	5,204	5,623	5,825	6,096	6,401	6,696	6,255	7,037	7,327	6,956
Minority	305	309	337	360	354	381	363	329	372	399	360

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	66%	68%	69%	70%	71%	72%	66%	71%	70%	65%
Female	65%	68%	70%	71%	72%	72%	73%	66%	72%	71%	66%
Male	59%	63%	64%	65%	67%	67%	69%	64%	67%	67%	63%
White	63%	67%	69%	70%	71%	71%	73%	66%	71%	71%	65%
Minority	36%	35%	36%	36%	35%	38%	36%	32%	36%	35%	30%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	525	560	637	667	710	768	824	781	898	963	953
Alcohol-related disorders	537	610	676	713	725	768	828	781	853	855	808
Anxiety disorders	2,828	3,140	3,527	3,748	3,961	4,202	4,530	4,354	4,837	4,960	4,807
Attention-deficit, conduct, and disruptive behavior disorders	380	447	530	570	633	717	808	808	954	1,083	1,069
Delirium, dementia, and amnesic and other cognitive disorders	1,341	1,428	1,445	1,389	1,359	1,362	1,359	1,262	1,254	1,226	1,102
Developmental disorders	618	652	711	744	769	803	853	836	876	921	912
Disorders usually diagnosed in infancy, childhood, adolescence	76	86	95	101	107	115	130	136	160	174	199
Impulse control disorders, NEC	133	157	157	174	194	220	247	245	270	284	292
Miscellaneous disorders	1,088	1,218	1,414	1,598	1,729	1,861	1,967	1,738	1,930	1,971	1,828
Mood disorders	3,019	3,360	3,759	3,976	4,163	4,411	4,720	4,490	4,967	5,111	4,914
Personality disorders	420	476	537	589	597	627	664	640	675	671	640
Schizophrenia and other psychotic disorders	1,212	1,326	1,385	1,394	1,393	1,438	1,479	1,405	1,417	1,426	1,335
Screening and history of mental health and substance abuse	3,058	3,428	3,846	4,172	4,427	4,698	4,974	4,636	5,314	5,538	5,291
Substance-related disorders	892	1,023	1,160	1,277	1,361	1,442	1,588	1,491	1,694	1,818	1,785
Suicide and intentional self-inflicted injury	219	250	283	305	321	352	405	384	453	481	483



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Carter, Knott, Lawrence, & Martin

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	6,508	6,925	7,036	7,304	7,283	7,217	7,227	6,781	7,014	6,832	6,320
Female	3,152	3,289	3,340	3,423	3,416	3,313	3,307	3,075	3,183	3,058	2,825
Male	3,356	3,636	3,696	3,881	3,867	3,904	3,920	3,706	3,831	3,774	3,495
White	6,090	6,469	6,539	6,769	6,730	6,648	6,600	6,193	6,396	6,211	5,739
Minority	59	58	62	65	68	64	55	50	53	52	55

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	56%	58%	59%	59%	59%	59%	56%	59%	56%	51%
Female	56%	56%	58%	58%	58%	57%	57%	54%	56%	53%	48%
Male	53%	56%	58%	60%	59%	60%	61%	58%	61%	59%	54%
White	53%	55%	56%	58%	57%	57%	57%	54%	57%	55%	50%
Minority	12%	11%	11%	10%	11%	10%	8%	7%	8%	7%	7%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,254	1,350	1,387	1,454	1,454	1,439	1,424	1,302	1,311	1,262	1,142
Alcohol-related disorders	309	291	261	252	227	194	181	155	144	127	104
Anxiety disorders	2,146	2,198	2,200	2,251	2,211	2,130	2,141	1,980	1,989	1,897	1,718
Attention-deficit, conduct, and disruptive behavior disorders	2,354	2,584	2,721	2,902	2,946	2,981	3,030	2,754	2,791	2,709	2,509
Delirium, dementia, and amnestic and other cognitive disorders	88	84	82	88	82	84	86	75	72	67	58
Developmental disorders	1,833	2,080	2,232	2,436	2,560	2,660	2,735	2,656	2,804	2,799	2,678
Disorders usually diagnosed in infancy, childhood, adolescence	403	457	482	532	542	557	589	567	595	575	549
Impulse control disorders, NEC	160	174	165	169	165	155	145	130	117	110	86
Miscellaneous disorders	828	835	834	852	824	807	797	726	719	702	636
Mood disorders	2,213	2,211	2,153	2,121	1,972	1,790	1,670	1,471	1,397	1,266	1,080
Personality disorders	198	200	192	195	186	164	154	123	107	98	78
Schizophrenia and other psychotic disorders	206	205	202	204	191	180	173	145	132	115	97
Screening and history of mental health and substance abuse	4,606	4,886	4,908	5,037	4,914	4,686	4,623	4,206	4,209	3,972	3,581
Substance-related disorders	857	833	747	721	641	547	511	444	423	392	330
Suicide and intentional self-inflicted injury	162	153	147	153	139	123	120	107	93	82	68



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Carter, Knott, Lawrence, & Martin

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	8,011	8,640	8,960	9,324	9,545	9,698	9,852	8,997	9,614	9,568	9,217
Female	5,291	5,705	5,873	6,098	6,219	6,331	6,439	5,795	6,163	6,132	5,851
Male	2,720	2,935	3,087	3,226	3,326	3,367	3,413	3,202	3,451	3,436	3,366
White	7,374	7,984	8,292	8,653	8,856	9,021	9,145	8,322	8,926	8,893	8,518
Minority	138	125	116	100	90	59	38	N.D.	34	34	38

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	66%	69%	69%	71%	70%	71%	72%	66%	71%	70%	67%
Female	70%	73%	73%	75%	74%	75%	76%	69%	74%	74%	70%
Male	59%	62%	63%	64%	64%	64%	64%	61%	65%	64%	62%
White	65%	68%	69%	70%	70%	70%	71%	65%	70%	70%	67%
Minority	16%	14%	13%	12%	10%	7%	4%	N.D.	4%	4%	4%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	673	729	768	855	900	935	989	914	993	981	968
Alcohol-related disorders	911	993	1,065	1,126	1,172	1,173	1,190	1,114	1,189	1,191	1,117
Anxiety disorders	5,020	5,451	5,699	5,971	6,125	6,246	6,434	5,983	6,270	6,274	6,071
Attention-deficit, conduct, and disruptive behavior disorders	226	263	289	318	359	411	487	473	544	572	620
Delirium, dementia, and amnestic and other cognitive disorders	1,470	1,503	1,484	1,472	1,443	1,388	1,335	1,212	1,168	1,088	981
Developmental disorders	731	762	789	826	857	878	901	850	891	901	924
Disorders usually diagnosed in infancy, childhood, adolescence	57	67	72	73	84	88	97	101	118	139	153
Impulse control disorders, NEC	154	178	194	200	204	223	243	228	257	251	241
Miscellaneous disorders	1,933	2,157	2,246	2,334	2,434	2,499	2,520	2,325	2,396	2,337	2,215
Mood disorders	4,839	5,247	5,493	5,746	5,926	6,094	6,275	5,769	6,127	6,114	5,878
Personality disorders	714	770	803	860	889	873	897	838	873	855	828
Schizophrenia and other psychotic disorders	1,616	1,698	1,725	1,740	1,718	1,694	1,692	1,561	1,565	1,518	1,428
Screening and history of mental health and substance abuse	4,989	5,491	5,879	6,215	6,487	6,635	6,830	6,270	6,833	6,807	6,559
Substance-related disorders	1,663	1,851	1,994	2,136	2,277	2,374	2,501	2,329	2,552	2,550	2,473
Suicide and intentional self-inflicted injury	271	295	303	325	337	357	387	373	406	385	382



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Casey, Clinton, Cumberland, Hart, & McCreary

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,533	6,029	6,242	6,587	6,723	6,778	6,782	6,432	6,477	6,271	5,826
Female	2,752	2,988	3,049	3,226	3,248	3,231	3,200	3,045	3,067	2,949	2,692
Male	2,781	3,041	3,193	3,361	3,475	3,547	3,582	3,387	3,410	3,322	3,134
White	5,161	5,592	5,770	6,095	6,221	6,269	6,233	5,895	5,927	5,742	5,316
Minority	140	162	165	167	157	151	142	137	141	124	125

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	57%	59%	61%	61%	61%	61%	58%	58%	55%	50%
Female	57%	59%	60%	62%	61%	61%	60%	57%	57%	53%	47%
Male	54%	56%	58%	61%	61%	62%	62%	59%	60%	57%	53%
White	55%	57%	59%	60%	61%	61%	60%	57%	57%	54%	49%
Minority	23%	24%	24%	25%	23%	22%	21%	19%	19%	16%	17%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,106	1,167	1,179	1,222	1,196	1,183	1,157	1,049	1,033	957	831
Alcohol-related disorders	227	227	223	217	195	177	168	133	106	73	55
Anxiety disorders	2,529	2,730	2,790	2,969	3,056	3,084	3,097	2,947	2,930	2,796	2,587
Attention-deficit, conduct, and disruptive behavior disorders	2,273	2,498	2,586	2,687	2,759	2,790	2,791	2,551	2,503	2,383	2,202
Delirium, dementia, and amnestic and other cognitive disorders	68	72	71	70	70	71	74	66	61	59	54
Developmental disorders	1,088	1,259	1,422	1,588	1,714	1,790	1,926	1,887	1,991	2,000	1,927
Disorders usually diagnosed in infancy, childhood, adolescence	356	389	426	463	495	502	513	494	500	483	468
Impulse control disorders, NEC	266	278	288	285	274	269	262	222	207	181	155
Miscellaneous disorders	848	892	888	901	890	862	810	743	724	660	593
Mood disorders	2,021	2,078	2,020	2,007	1,922	1,820	1,702	1,454	1,363	1,197	1,035
Personality disorders	139	128	126	124	121	109	103	90	85	74	58
Schizophrenia and other psychotic disorders	171	174	183	181	182	174	164	142	141	123	100
Screening and history of mental health and substance abuse	2,884	3,077	3,164	3,263	3,199	3,094	2,964	2,637	2,553	2,397	2,143
Substance-related disorders	629	641	609	578	531	468	420	338	308	260	205
Suicide and intentional self-inflicted injury	156	157	161	157	151	134	130	102	93	83	72



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Casey, Clinton, Cumberland, Hart, & McCreary

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	7,468	7,907	8,117	8,350	8,492	8,528	8,727	8,178	8,515	8,641	8,326
Female	5,016	5,336	5,449	5,552	5,656	5,671	5,805	5,404	5,635	5,674	5,456
Male	2,452	2,571	2,668	2,798	2,836	2,857	2,922	2,774	2,880	2,967	2,870
White	6,696	7,113	7,335	7,544	7,677	7,738	7,922	7,427	7,765	7,894	7,558
Minority	324	313	302	308	291	262	223	188	188	190	195

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	67%	70%	71%	72%	72%	71%	72%	68%	70%	70%	67%
Female	70%	73%	75%	75%	75%	75%	76%	70%	73%	74%	70%
Male	61%	63%	64%	66%	65%	65%	66%	63%	64%	64%	62%
White	67%	69%	70%	71%	71%	71%	72%	67%	69%	70%	67%
Minority	30%	29%	29%	29%	28%	25%	22%	19%	19%	19%	19%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	789	851	901	950	998	1,035	1,077	1,005	1,095	1,122	1,086
Alcohol-related disorders	724	776	809	835	843	839	867	807	826	831	810
Anxiety disorders	4,651	5,003	5,220	5,443	5,649	5,750	5,976	5,659	5,873	5,983	5,815
Attention-deficit, conduct, and disruptive behavior disorders	250	278	309	339	386	415	479	488	558	639	670
Delirium, dementia, and amnestic and other cognitive disorders	1,803	1,841	1,797	1,742	1,682	1,586	1,531	1,425	1,334	1,231	1,123
Developmental disorders	663	686	703	715	732	754	774	753	769	792	771
Disorders usually diagnosed in infancy, childhood, adolescence	51	59	61	70	74	82	89	92	104	116	124
Impulse control disorders, NEC	85	89	100	109	123	121	138	138	141	164	173
Miscellaneous disorders	1,778	1,937	2,012	2,116	2,189	2,237	2,348	2,207	2,272	2,323	2,205
Mood disorders	4,612	4,942	5,153	5,359	5,539	5,570	5,758	5,414	5,594	5,679	5,495
Personality disorders	475	497	524	534	538	521	527	479	477	479	469
Schizophrenia and other psychotic disorders	1,662	1,741	1,750	1,752	1,750	1,710	1,735	1,649	1,588	1,551	1,440
Screening and history of mental health and substance abuse	3,741	4,017	4,262	4,462	4,629	4,646	4,786	4,522	4,746	4,846	4,632
Substance-related disorders	1,237	1,324	1,383	1,492	1,524	1,555	1,630	1,562	1,655	1,770	1,747
Suicide and intentional self-inflicted injury	272	289	304	325	335	352	379	364	385	413	417



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Christian, Hopkins, & McLean

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	6,679	7,289	7,669	8,102	8,425	8,728	8,671	7,889	8,184	8,248	7,796
Female	3,283	3,545	3,660	3,870	4,000	4,089	4,033	3,628	3,753	3,752	3,515
Male	3,396	3,744	4,009	4,232	4,425	4,639	4,638	4,261	4,431	4,496	4,281
White	4,280	4,680	4,903	5,176	5,397	5,557	5,507	5,027	5,243	5,252	4,985
Minority	1,977	2,130	2,211	2,306	2,355	2,452	2,387	2,103	2,164	2,207	2,077

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	56%	58%	59%	59%	60%	60%	55%	56%	53%	48%
Female	54%	56%	57%	58%	58%	58%	57%	52%	53%	50%	45%
Male	54%	56%	59%	60%	61%	62%	62%	57%	58%	56%	51%
White	55%	56%	58%	59%	59%	60%	58%	54%	55%	52%	47%
Minority	44%	45%	45%	46%	46%	47%	47%	41%	42%	41%	38%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,071	1,195	1,230	1,284	1,305	1,305	1,255	1,106	1,128	1,106	1,037
Alcohol-related disorders	173	189	182	167	153	147	115	87	81	82	62
Anxiety disorders	1,473	1,540	1,622	1,679	1,679	1,697	1,642	1,457	1,453	1,366	1,216
Attention-deficit, conduct, and disruptive behavior disorders	3,427	3,772	4,049	4,292	4,408	4,533	4,460	4,045	4,074	4,012	3,786
Delirium, dementia, and amnestic and other cognitive disorders	106	115	104	115	116	126	119	107	108	106	90
Developmental disorders	1,234	1,436	1,628	1,823	1,975	2,122	2,238	2,152	2,289	2,357	2,244
Disorders usually diagnosed in infancy, childhood, adolescence	348	412	458	511	536	576	602	577	575	578	558
Impulse control disorders, NEC	757	789	795	780	750	725	659	573	540	498	433
Miscellaneous disorders	1,690	1,772	1,787	1,828	1,821	1,770	1,686	1,472	1,426	1,369	1,249
Mood disorders	2,235	2,389	2,400	2,438	2,349	2,267	2,111	1,835	1,775	1,649	1,441
Personality disorders	114	116	116	106	100	89	85	77	64	56	41
Schizophrenia and other psychotic disorders	201	204	216	219	221	196	185	166	156	147	126
Screening and history of mental health and substance abuse	3,841	4,158	4,312	4,461	4,618	4,759	4,698	4,200	4,257	4,179	3,914
Substance-related disorders	581	604	581	557	515	474	407	325	326	300	240
Suicide and intentional self-inflicted injury	166	173	167	167	169	153	140	114	109	102	82



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Christian, Hopkins, & McLean

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	6,410	6,861	7,142	7,372	7,561	7,800	8,078	7,402	7,995	8,174	7,776
Female	4,699	4,986	5,203	5,361	5,487	5,632	5,799	5,224	5,678	5,809	5,460
Male	1,711	1,875	1,939	2,011	2,074	2,168	2,279	2,178	2,317	2,365	2,316
White	4,743	5,045	5,268	5,428	5,572	5,752	5,966	5,401	5,854	6,006	5,663
Minority	1,418	1,548	1,592	1,639	1,657	1,690	1,721	1,606	1,742	1,768	1,700

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	65%	66%	68%	68%	68%	70%	64%	68%	67%	62%
Female	64%	67%	68%	70%	70%	70%	71%	64%	70%	69%	63%
Male	57%	60%	62%	64%	64%	65%	67%	62%	65%	63%	59%
White	64%	67%	68%	69%	70%	70%	71%	64%	69%	68%	63%
Minority	48%	51%	52%	54%	54%	54%	55%	51%	54%	52%	48%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	647	722	781	815	845	912	946	880	960	999	930
Alcohol-related disorders	554	584	609	634	670	667	713	666	709	736	693
Anxiety disorders	3,018	3,281	3,509	3,681	3,786	3,914	4,143	3,872	4,090	4,088	3,947
Attention-deficit, conduct, and disruptive behavior disorders	524	585	623	683	759	832	924	951	1,050	1,173	1,176
Delirium, dementia, and amnestic and other cognitive disorders	1,878	1,931	1,916	1,854	1,779	1,721	1,701	1,581	1,589	1,489	1,373
Developmental disorders	927	953	977	960	962	941	985	957	939	943	926
Disorders usually diagnosed in infancy, childhood, adolescence	77	83	86	96	96	109	109	117	128	131	126
Impulse control disorders, NEC	162	176	184	210	230	260	313	307	330	354	363
Miscellaneous disorders	1,375	1,488	1,565	1,612	1,679	1,728	1,821	1,661	1,799	1,838	1,776
Mood disorders	3,399	3,740	4,043	4,253	4,437	4,670	4,869	4,610	4,956	5,023	4,826
Personality disorders	388	414	427	446	468	472	473	460	477	476	444
Schizophrenia and other psychotic disorders	1,823	1,904	1,909	1,903	1,862	1,837	1,860	1,787	1,789	1,762	1,661
Screening and history of mental health and substance abuse	3,366	3,713	4,018	4,288	4,507	4,755	5,022	4,551	5,002	5,172	4,886
Substance-related disorders	770	854	957	1,034	1,081	1,157	1,275	1,176	1,315	1,406	1,306
Suicide and intentional self-inflicted injury	251	277	284	314	326	358	390	386	456	486	449



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Clay, Leslie, & Letcher

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,981	6,159	6,230	6,288	6,211	6,142	6,043	5,694	5,764	5,591	5,123
Female	2,936	2,972	2,965	2,955	2,890	2,848	2,817	2,615	2,647	2,533	2,319
Male	3,045	3,187	3,265	3,333	3,321	3,294	3,226	3,079	3,117	3,058	2,804
White	5,560	5,700	5,743	5,776	5,693	5,601	5,460	5,125	5,178	5,006	4,590
Minority	63	64	64	66	58	54	54	43	47	48	47

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	58%	59%	59%	59%	60%	60%	57%	59%	56%	51%
Female	59%	60%	61%	60%	59%	60%	60%	56%	57%	54%	49%
Male	53%	56%	58%	59%	59%	60%	60%	59%	61%	59%	53%
White	54%	56%	58%	58%	58%	58%	58%	55%	57%	54%	49%
Minority	12%	12%	11%	11%	9%	8%	8%	6%	7%	6%	7%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,014	1,043	1,064	1,072	1,050	1,012	990	919	905	849	738
Alcohol-related disorders	331	329	320	296	294	262	234	202	178	166	127
Anxiety disorders	2,455	2,477	2,479	2,452	2,402	2,358	2,296	2,131	2,081	1,973	1,753
Attention-deficit, conduct, and disruptive behavior disorders	1,756	1,868	1,962	2,013	2,026	2,035	2,021	1,910	1,905	1,822	1,634
Delirium, dementia, and amnestic and other cognitive disorders	88	85	88	84	82	84	84	71	63	62	55
Developmental disorders	1,817	1,952	2,058	2,196	2,263	2,285	2,299	2,214	2,241	2,198	2,023
Disorders usually diagnosed in infancy, childhood, adolescence	302	331	354	370	383	387	388	371	374	349	317
Impulse control disorders, NEC	169	174	176	171	165	143	140	129	111	92	72
Miscellaneous disorders	1,216	1,205	1,203	1,177	1,127	1,093	1,045	956	944	891	814
Mood disorders	1,875	1,825	1,736	1,624	1,499	1,352	1,231	1,049	935	833	669
Personality disorders	262	258	257	243	235	214	195	179	159	146	106
Schizophrenia and other psychotic disorders	189	190	191	181	178	160	152	143	126	109	86
Screening and history of mental health and substance abuse	3,404	3,443	3,412	3,323	3,189	2,983	2,857	2,605	2,556	2,418	2,137
Substance-related disorders	970	906	848	770	686	614	555	482	454	401	320
Suicide and intentional self-inflicted injury	136	133	114	114	103	97	95	82	70	64	48



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Clay, Leslie, & Letcher

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	8,484	8,913	9,283	9,683	9,922	9,926	10,072	9,511	9,924	9,871	9,625
Female	5,357	5,641	5,876	6,105	6,255	6,245	6,289	5,947	6,199	6,150	5,948
Male	3,127	3,272	3,407	3,578	3,667	3,681	3,783	3,564	3,725	3,721	3,677
White	7,799	8,229	8,594	8,990	9,223	9,224	9,372	8,834	9,258	9,193	8,917
Minority	184	174	159	141	125	112	83	74	76	84	77

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	69%	71%	72%	73%	73%	73%	74%	71%	74%	74%	71%
Female	72%	74%	76%	77%	77%	77%	78%	74%	77%	77%	74%
Male	64%	65%	66%	67%	67%	68%	70%	66%	69%	69%	67%
White	68%	70%	71%	72%	72%	73%	74%	70%	74%	74%	71%
Minority	21%	20%	18%	17%	15%	13%	10%	9%	9%	9%	8%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	814	855	897	969	1,020	1,028	1,094	1,031	1,093	1,120	1,126
Alcohol-related disorders	959	1,027	1,080	1,172	1,212	1,210	1,225	1,158	1,172	1,164	1,151
Anxiety disorders	5,964	6,321	6,589	6,984	7,212	7,225	7,388	7,033	7,247	7,200	7,074
Attention-deficit, conduct, and disruptive behavior disorders	273	295	327	371	426	468	497	491	547	629	659
Delirium, dementia, and amnestic and other cognitive disorders	1,418	1,456	1,435	1,387	1,360	1,330	1,266	1,140	1,101	1,025	944
Developmental disorders	657	674	688	718	762	790	825	791	838	859	898
Disorders usually diagnosed in infancy, childhood, adolescence	49	56	55	55	66	76	78	84	92	112	127
Impulse control disorders, NEC	142	156	155	170	185	194	192	174	185	196	205
Miscellaneous disorders	2,965	3,160	3,354	3,591	3,777	3,782	3,835	3,653	3,719	3,669	3,563
Mood disorders	5,174	5,465	5,702	6,048	6,280	6,337	6,468	6,226	6,429	6,382	6,267
Personality disorders	658	686	730	804	871	862	839	779	797	789	794
Schizophrenia and other psychotic disorders	1,608	1,666	1,677	1,680	1,666	1,641	1,606	1,505	1,481	1,412	1,355
Screening and history of mental health and substance abuse	5,165	5,538	5,866	6,208	6,446	6,528	6,696	6,342	6,582	6,533	6,332
Substance-related disorders	2,483	2,646	2,812	3,068	3,264	3,333	3,471	3,341	3,478	3,496	3,491
Suicide and intentional self-inflicted injury	303	323	345	371	393	420	448	433	453	454	446



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Caldwell, Crittenden, Livingston, Lyon, & Union

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,575	2,781	3,005	3,063	3,004	2,995	3,050	2,831	2,912	2,916	2,728
Female	1,282	1,380	1,462	1,466	1,425	1,418	1,442	1,337	1,366	1,363	1,252
Male	1,293	1,401	1,543	1,597	1,579	1,577	1,608	1,494	1,546	1,553	1,476
White	2,246	2,434	2,614	2,644	2,561	2,543	2,588	2,404	2,469	2,454	2,312
Minority	206	204	226	233	237	234	218	191	194	212	180

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	54%	59%	58%	56%	56%	57%	52%	52%	49%	44%
Female	54%	55%	57%	56%	54%	55%	55%	50%	50%	47%	42%
Male	54%	53%	61%	60%	58%	58%	58%	53%	54%	51%	47%
White	54%	55%	58%	57%	55%	55%	55%	50%	50%	47%	43%
Minority	34%	27%	38%	36%	35%	34%	33%	29%	28%	29%	24%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	392	412	446	437	427	438	449	403	398	387	365
Alcohol-related disorders	79	79	80	75	69	64	59	46	45	33	N.D.
Anxiety disorders	741	772	816	834	807	795	808	722	711	678	635
Attention-deficit, conduct, and disruptive behavior disorders	1,274	1,383	1,520	1,581	1,600	1,616	1,661	1,559	1,559	1,497	1,396
Delirium, dementia, and amnestic and other cognitive disorders	41	39	43	42	46	42	41	36	39	36	33
Developmental disorders	495	578	681	750	794	831	880	857	898	933	913
Disorders usually diagnosed in infancy, childhood, adolescence	126	136	159	162	171	191	213	206	225	237	227
Impulse control disorders, NEC	213	221	227	234	227	230	222	199	196	176	159
Miscellaneous disorders	336	354	367	362	347	345	335	298	305	303	267
Mood disorders	1,076	1,091	1,130	1,108	1,085	1,057	1,012	881	858	812	718
Personality disorders	62	66	69	70	64	60	55	54	51	44	40
Schizophrenia and other psychotic disorders	89	89	90	91	81	77	74	67	60	53	48
Screening and history of mental health and substance abuse	1,445	1,536	1,636	1,568	1,447	1,391	1,349	1,221	1,233	1,174	1,058
Substance-related disorders	205	206	213	207	201	192	173	137	137	120	101
Suicide and intentional self-inflicted injury	76	82	83	86	81	78	75	67	61	53	44



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Caldwell, Crittenden, Livingston, Lyon, & Union

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,790	3,019	3,170	3,238	3,330	3,367	3,468	3,158	3,478	3,496	3,393
Female	2,056	2,217	2,311	2,361	2,407	2,417	2,486	2,238	2,464	2,469	2,358
Male	734	802	859	877	923	950	982	920	1,014	1,027	1,035
White	2,484	2,690	2,833	2,886	2,973	2,994	3,087	2,791	3,099	3,108	2,994
Minority	202	214	215	219	212	215	205	193	205	202	202

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	61%	63%	67%	67%	68%	67%	69%	62%	68%	67%	63%
Female	64%	67%	69%	69%	70%	70%	72%	64%	70%	69%	65%
Male	55%	55%	62%	62%	62%	62%	63%	59%	64%	63%	60%
White	62%	65%	66%	66%	67%	67%	69%	62%	68%	67%	63%
Minority	40%	35%	45%	44%	42%	41%	39%	37%	39%	36%	33%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	191	210	232	231	236	234	255	239	277	286	264
Alcohol-related disorders	191	201	219	229	235	245	269	237	250	269	262
Anxiety disorders	1,408	1,541	1,682	1,801	1,890	1,972	2,070	1,894	2,071	2,090	2,013
Attention-deficit, conduct, and disruptive behavior disorders	193	217	254	275	299	318	354	354	419	455	457
Delirium, dementia, and amnestic and other cognitive disorders	783	803	788	749	730	686	679	633	632	576	521
Developmental disorders	279	287	308	301	313	321	330	317	332	332	338
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	36	38	37	39	41	44	53
Impulse control disorders, NEC	46	54	59	59	65	71	75	76	86	96	96
Miscellaneous disorders	567	631	660	697	740	791	803	717	799	820	787
Mood disorders	1,593	1,767	1,920	2,021	2,085	2,138	2,257	2,082	2,304	2,331	2,226
Personality disorders	113	141	145	150	168	173	186	162	172	175	167
Schizophrenia and other psychotic disorders	607	641	687	674	671	677	709	669	698	664	615
Screening and history of mental health and substance abuse	1,333	1,493	1,620	1,727	1,789	1,856	1,936	1,705	1,872	1,909	1,843
Substance-related disorders	369	393	449	482	513	540	580	538	574	589	570
Suicide and intentional self-inflicted injury	99	103	127	137	157	178	202	187	213	223	216



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Daviess

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,935	5,464	5,742	6,048	6,115	6,211	6,289	5,982	6,177	6,173	5,802
Female	2,436	2,664	2,735	2,862	2,856	2,920	2,883	2,712	2,788	2,752	2,561
Male	2,499	2,800	3,007	3,186	3,259	3,291	3,406	3,270	3,389	3,421	3,241
White	3,937	4,350	4,540	4,760	4,806	4,871	4,912	4,674	4,806	4,795	4,488
Minority	643	692	735	778	776	771	762	711	760	747	723

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	62%	63%	63%	62%	61%	61%	57%	56%	53%	48%
Female	61%	62%	62%	62%	60%	59%	58%	54%	52%	49%	44%
Male	60%	62%	64%	64%	64%	62%	64%	60%	60%	58%	52%
White	59%	61%	62%	61%	60%	59%	59%	55%	54%	52%	46%
Minority	43%	43%	43%	42%	41%	39%	38%	35%	35%	33%	31%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,085	1,182	1,225	1,279	1,263	1,277	1,275	1,160	1,127	1,078	1,004
Alcohol-related disorders	210	223	206	198	184	179	163	145	137	121	92
Anxiety disorders	1,493	1,600	1,628	1,745	1,732	1,757	1,768	1,664	1,685	1,608	1,469
Attention-deficit, conduct, and disruptive behavior disorders	2,423	2,679	2,832	3,010	3,068	3,092	3,109	2,914	2,922	2,827	2,614
Delirium, dementia, and amnestic and other cognitive disorders	70	72	75	77	78	77	73	67	70	67	59
Developmental disorders	874	1,045	1,219	1,371	1,513	1,668	1,837	1,900	2,022	2,094	2,053
Disorders usually diagnosed in infancy, childhood, adolescence	351	406	439	461	484	514	548	526	542	539	522
Impulse control disorders, NEC	499	543	564	580	583	579	564	510	487	456	390
Miscellaneous disorders	698	737	736	715	680	671	665	593	586	553	499
Mood disorders	2,114	2,245	2,284	2,297	2,237	2,181	2,113	1,913	1,886	1,787	1,574
Personality disorders	238	239	220	211	202	187	175	159	150	132	102
Schizophrenia and other psychotic disorders	273	282	281	281	273	261	246	236	228	200	167
Screening and history of mental health and substance abuse	3,425	3,725	3,833	3,813	3,675	3,571	3,442	3,140	3,132	3,074	2,755
Substance-related disorders	505	529	503	487	446	422	401	369	356	299	228
Suicide and intentional self-inflicted injury	146	157	141	133	131	124	127	107	95	85	70



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Daviess

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,346	4,790	5,146	5,491	5,744	5,983	6,152	5,739	6,320	6,451	6,259
Female	3,240	3,584	3,841	4,102	4,278	4,415	4,490	4,167	4,529	4,561	4,409
Male	1,106	1,206	1,305	1,389	1,466	1,568	1,662	1,572	1,791	1,890	1,850
White	3,745	4,138	4,432	4,707	4,940	5,145	5,291	4,901	5,459	5,545	5,309
Minority	421	455	491	538	539	555	548	530	548	576	587

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	68%	70%	72%	73%	74%	73%	68%	72%	71%	67%
Female	67%	70%	72%	75%	75%	76%	75%	69%	74%	73%	68%
Male	60%	63%	64%	66%	68%	68%	69%	65%	69%	68%	64%
White	65%	69%	70%	72%	74%	74%	74%	68%	72%	71%	67%
Minority	44%	45%	48%	49%	47%	48%	46%	45%	46%	45%	41%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	610	727	808	860	885	883	932	862	927	946	935
Alcohol-related disorders	442	514	542	607	659	691	744	716	774	791	776
Anxiety disorders	2,242	2,512	2,740	2,983	3,146	3,364	3,532	3,421	3,732	3,776	3,700
Attention-deficit, conduct, and disruptive behavior disorders	421	472	546	634	722	796	890	919	1,057	1,138	1,177
Delirium, dementia, and amnesic and other cognitive disorders	944	953	986	978	962	961	923	871	876	847	791
Developmental disorders	623	658	691	721	747	754	777	766	792	827	832
Disorders usually diagnosed in infancy, childhood, adolescence	49	55	61	70	76	79	90	95	106	117	139
Impulse control disorders, NEC	117	141	154	164	178	191	215	204	248	283	292
Miscellaneous disorders	1,028	1,168	1,325	1,450	1,571	1,672	1,704	1,517	1,631	1,620	1,558
Mood disorders	2,594	2,913	3,168	3,489	3,708	3,892	4,091	3,920	4,234	4,286	4,171
Personality disorders	517	564	627	657	682	714	724	695	744	756	727
Schizophrenia and other psychotic disorders	1,213	1,298	1,348	1,403	1,416	1,457	1,515	1,439	1,469	1,446	1,395
Screening and history of mental health and substance abuse	2,522	2,944	3,233	3,590	3,877	4,097	4,258	3,956	4,445	4,537	4,401
Substance-related disorders	730	847	917	1,010	1,101	1,190	1,286	1,239	1,370	1,423	1,374
Suicide and intentional self-inflicted injury	215	248	285	306	332	365	399	403	467	480	474



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Estill, Floyd, Johnson, Lee, Magoffin, & Powell

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	11,096	11,724	12,016	12,478	12,576	12,538	12,620	11,872	12,091	12,059	11,226
Female	5,630	5,822	5,870	6,036	6,056	5,962	5,964	5,535	5,605	5,558	5,193
Male	5,466	5,902	6,146	6,442	6,520	6,576	6,656	6,337	6,486	6,501	6,033
White	10,307	10,845	11,053	11,431	11,461	11,384	11,399	10,702	10,892	10,837	10,096
Minority	93	90	96	96	96	89	83	84	76	73	66

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	59%	60%	62%	63%	63%	63%	64%	60%	61%	60%	55%
Female	61%	62%	63%	63%	63%	62%	63%	59%	59%	58%	53%
Male	56%	59%	61%	63%	63%	64%	65%	62%	63%	62%	57%
White	57%	59%	61%	62%	61%	61%	62%	59%	59%	58%	53%
Minority	10%	9%	8%	8%	8%	7%	6%	6%	5%	5%	5%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,027	2,196	2,238	2,344	2,350	2,319	2,298	2,069	2,032	1,961	1,767
Alcohol-related disorders	475	463	449	430	370	336	319	256	232	197	157
Anxiety disorders	4,155	4,340	4,418	4,606	4,708	4,710	4,771	4,466	4,509	4,449	4,156
Attention-deficit, conduct, and disruptive behavior disorders	3,725	4,112	4,425	4,721	4,848	4,903	4,965	4,584	4,606	4,537	4,185
Delirium, dementia, and amnestic and other cognitive disorders	135	133	139	140	129	118	107	97	98	98	90
Developmental disorders	2,335	2,651	2,956	3,248	3,452	3,622	3,794	3,701	3,920	3,965	3,767
Disorders usually diagnosed in infancy, childhood, adolescence	687	780	849	924	989	1,033	1,082	1,028	1,052	1,051	992
Impulse control disorders, NEC	262	270	272	258	257	237	232	206	180	156	123
Miscellaneous disorders	1,870	1,897	1,939	1,990	2,009	2,000	2,018	1,874	1,862	1,779	1,662
Mood disorders	3,253	3,231	3,077	2,982	2,844	2,629	2,503	2,111	1,961	1,823	1,549
Personality disorders	250	251	240	237	245	227	212	184	167	143	116
Schizophrenia and other psychotic disorders	317	307	306	308	290	278	265	220	210	196	165
Screening and history of mental health and substance abuse	7,522	7,818	7,867	8,001	7,818	7,507	7,259	6,592	6,544	6,335	5,722
Substance-related disorders	1,313	1,268	1,165	1,100	1,002	867	804	654	637	590	506
Suicide and intentional self-inflicted injury	318	313	299	296	274	251	234	194	180	175	152



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Estill, Floyd, Johnson, Lee, Magoffin, & Powell

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	13,722	14,548	15,304	15,919	16,235	16,516	16,987	16,114	17,230	17,286	16,970
Female	9,079	9,628	10,069	10,426	10,601	10,816	11,148	10,506	11,231	11,161	10,902
Male	4,643	4,920	5,235	5,493	5,634	5,700	5,839	5,608	5,999	6,125	6,068
White	12,675	13,487	14,194	14,795	15,098	15,397	15,807	14,952	16,038	16,078	15,706
Minority	239	218	206	177	143	107	74	58	70	75	80

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	69%	71%	73%	74%	73%	74%	75%	71%	74%	74%	71%
Female	73%	75%	77%	78%	77%	78%	80%	74%	78%	78%	75%
Male	62%	65%	67%	67%	67%	67%	68%	64%	68%	67%	65%
White	68%	71%	73%	73%	73%	73%	75%	70%	74%	73%	70%
Minority	17%	16%	15%	13%	10%	8%	5%	4%	5%	5%	5%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,509	1,600	1,731	1,869	1,947	2,050	2,126	2,130	2,256	2,288	2,264
Alcohol-related disorders	1,305	1,404	1,479	1,567	1,620	1,615	1,722	1,647	1,708	1,705	1,622
Anxiety disorders	9,215	9,806	10,404	10,867	11,214	11,461	11,879	11,433	12,006	11,989	11,728
Attention-deficit, conduct, and disruptive behavior disorders	409	449	485	540	587	669	756	779	895	1,029	1,100
Delirium, dementia, and amnesic and other cognitive disorders	2,329	2,366	2,354	2,305	2,239	2,165	2,130	1,959	1,924	1,801	1,632
Developmental disorders	955	995	1,010	1,063	1,066	1,089	1,153	1,135	1,151	1,193	1,212
Disorders usually diagnosed in infancy, childhood, adolescence	133	146	148	165	175	193	200	207	217	234	239
Impulse control disorders, NEC	142	154	158	185	193	208	222	224	242	260	262
Miscellaneous disorders	4,283	4,646	4,915	5,218	5,399	5,562	5,718	5,442	5,597	5,497	5,231
Mood disorders	8,053	8,647	9,135	9,631	9,907	10,163	10,553	10,086	10,599	10,613	10,406
Personality disorders	765	827	875	928	963	988	1,019	999	1,015	1,003	974
Schizophrenia and other psychotic disorders	2,278	2,380	2,429	2,438	2,440	2,406	2,428	2,273	2,273	2,204	2,074
Screening and history of mental health and substance abuse	8,864	9,585	10,278	10,894	11,240	11,590	12,065	11,519	12,269	12,367	12,169
Substance-related disorders	3,017	3,243	3,469	3,715	3,891	4,080	4,342	4,243	4,550	4,594	4,529
Suicide and intentional self-inflicted injury	445	488	519	585	613	648	716	707	739	758	748



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Fayette

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	7,333	8,272	8,945	9,444	9,683	9,926	10,213	9,838	10,551	10,766	10,199
Female	3,699	4,094	4,356	4,473	4,537	4,602	4,667	4,370	4,705	4,810	4,484
Male	3,634	4,178	4,589	4,971	5,146	5,324	5,546	5,468	5,846	5,956	5,715
White	3,810	4,362	4,707	4,981	5,226	5,431	5,577	5,428	5,897	6,036	5,783
Minority	3,108	3,454	3,710	3,884	3,851	3,827	3,889	3,689	3,860	3,910	3,645

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	50%	51%	52%	52%	51%	50%	50%	46%	46%	43%	39%
Female	50%	50%	51%	50%	49%	48%	47%	42%	43%	40%	35%
Male	50%	52%	54%	54%	53%	52%	52%	49%	50%	47%	42%
White	51%	52%	53%	53%	51%	49%	47%	43%	43%	40%	36%
Minority	43%	44%	45%	45%	44%	44%	45%	42%	42%	40%	35%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,962	2,258	2,450	2,592	2,666	2,732	2,725	2,549	2,620	2,557	2,369
Alcohol-related disorders	342	384	374	355	352	360	344	282	276	250	194
Anxiety disorders	2,186	2,414	2,598	2,751	2,869	2,896	2,919	2,760	2,818	2,820	2,625
Attention-deficit, conduct, and disruptive behavior disorders	3,940	4,418	4,772	5,062	5,153	5,265	5,368	5,099	5,217	5,156	4,825
Delirium, dementia, and amnestic and other cognitive disorders	177	203	216	226	244	246	246	228	226	217	190
Developmental disorders	1,531	1,811	2,127	2,371	2,552	2,773	3,048	3,121	3,511	3,703	3,566
Disorders usually diagnosed in infancy, childhood, adolescence	775	887	1,006	1,088	1,165	1,243	1,290	1,249	1,310	1,305	1,251
Impulse control disorders, NEC	377	389	405	420	408	380	368	322	293	266	225
Miscellaneous disorders	1,126	1,215	1,264	1,268	1,286	1,284	1,264	1,175	1,246	1,196	1,080
Mood disorders	2,653	2,845	2,925	2,970	2,949	2,892	2,786	2,508	2,458	2,362	2,113
Personality disorders	258	267	263	256	254	231	223	198	173	165	137
Schizophrenia and other psychotic disorders	412	427	434	452	445	423	426	396	391	364	314
Screening and history of mental health and substance abuse	4,611	5,034	5,291	5,346	5,277	5,138	5,151	4,719	4,751	4,627	4,148
Substance-related disorders	995	1,062	1,070	1,010	976	945	895	763	744	666	553
Suicide and intentional self-inflicted injury	334	351	376	386	397	379	388	349	347	339	286



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Fayette

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	7,887	8,546	8,896	9,248	9,456	9,669	9,978	9,299	10,247	10,625	10,265
Female	5,923	6,409	6,645	6,882	7,044	7,176	7,359	6,813	7,573	7,787	7,412
Male	1,964	2,137	2,251	2,366	2,412	2,493	2,619	2,486	2,674	2,838	2,853
White	5,086	5,516	5,731	5,987	6,117	6,287	6,545	6,061	6,680	6,844	6,571
Minority	2,471	2,673	2,779	2,849	2,904	2,897	2,908	2,710	3,027	3,220	3,062

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	61%	64%	64%	65%	65%	65%	66%	61%	66%	64%	60%
Female	62%	65%	65%	64%	64%	64%	66%	61%	65%	64%	59%
Male	58%	61%	64%	66%	65%	66%	68%	63%	67%	65%	60%
White	64%	67%	68%	69%	69%	69%	70%	64%	68%	66%	61%
Minority	50%	51%	51%	51%	50%	50%	51%	48%	52%	52%	47%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	818	913	961	1,020	1,082	1,128	1,226	1,223	1,377	1,440	1,416
Alcohol-related disorders	974	1,078	1,163	1,240	1,274	1,278	1,372	1,279	1,409	1,456	1,421
Anxiety disorders	3,968	4,359	4,672	4,956	5,153	5,356	5,634	5,384	5,833	5,930	5,743
Attention-deficit, conduct, and disruptive behavior disorders	533	611	686	793	866	960	1,089	1,128	1,311	1,449	1,474
Delirium, dementia, and amnestic and other cognitive disorders	2,345	2,392	2,380	2,305	2,259	2,205	2,135	1,930	1,899	1,815	1,685
Developmental disorders	909	960	1,037	1,110	1,137	1,160	1,205	1,150	1,217	1,256	1,269
Disorders usually diagnosed in infancy, childhood, adolescence	112	129	139	159	177	177	211	216	251	280	300
Impulse control disorders, NEC	153	168	179	190	211	233	261	270	297	318	305
Miscellaneous disorders	2,003	2,262	2,378	2,518	2,657	2,802	2,936	2,721	3,036	3,138	2,978
Mood disorders	4,469	4,905	5,222	5,500	5,696	5,879	6,158	5,871	6,344	6,501	6,311
Personality disorders	806	864	937	966	960	994	1,056	998	1,062	1,072	1,041
Schizophrenia and other psychotic disorders	2,162	2,261	2,302	2,300	2,338	2,328	2,397	2,237	2,253	2,207	2,095
Screening and history of mental health and substance abuse	4,576	5,140	5,526	5,886	6,132	6,288	6,553	6,144	6,796	7,082	6,848
Substance-related disorders	1,643	1,825	1,944	2,155	2,295	2,371	2,580	2,415	2,741	2,915	2,795
Suicide and intentional self-inflicted injury	388	419	441	495	520	577	666	666	748	764	764



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Fleming & Lewis

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,898	2,026	2,149	2,235	2,262	2,217	2,270	2,093	2,167	2,203	2,003
Female	966	1,027	1,063	1,104	1,104	1,061	1,081	979	1,011	1,035	936
Male	932	999	1,086	1,131	1,158	1,156	1,189	1,114	1,156	1,168	1,067
White	1,784	1,892	2,007	2,081	2,103	2,047	2,088	1,919	1,978	2,016	1,832
Minority	37	42	42	43	40	37	36	33	37	36	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	53%	55%	55%	56%	54%	55%	50%	51%	50%	45%
Female	55%	54%	56%	55%	55%	53%	53%	48%	49%	48%	43%
Male	51%	51%	55%	55%	57%	56%	56%	53%	54%	52%	47%
White	52%	52%	54%	54%	55%	53%	53%	49%	50%	49%	44%
Minority	24%	24%	23%	21%	19%	17%	16%	15%	15%	13%	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	503	519	567	584	597	594	611	580	609	595	531
Alcohol-related disorders	84	79	80	73	68	60	57	44	43	33	31
Anxiety disorders	610	650	692	713	696	659	659	589	610	594	530
Attention-deficit, conduct, and disruptive behavior disorders	740	796	862	904	926	926	945	842	881	860	778
Delirium, dementia, and amnestic and other cognitive disorders	34	33	N.D.	N.D.	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	474	526	606	661	721	743	791	748	807	817	765
Disorders usually diagnosed in infancy, childhood, adolescence	161	173	196	214	230	226	229	205	222	219	210
Impulse control disorders, NEC	63	66	70	71	67	65	65	51	45	35	N.D.
Miscellaneous disorders	293	294	311	325	315	315	310	279	285	273	246
Mood disorders	647	669	688	674	639	599	577	477	480	456	399
Personality disorders	201	205	211	212	207	203	180	148	144	131	117
Schizophrenia and other psychotic disorders	124	114	119	120	120	119	114	99	95	92	85
Screening and history of mental health and substance abuse	1,312	1,373	1,432	1,465	1,441	1,379	1,369	1,215	1,219	1,191	1,083
Substance-related disorders	195	205	199	195	181	156	139	108	95	82	68
Suicide and intentional self-inflicted injury	44	44	46	44	39	39	36	N.D.	35	33	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Fleming & Lewis

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,172	2,297	2,417	2,510	2,620	2,623	2,756	2,593	2,802	2,784	2,693
Female	1,443	1,516	1,608	1,655	1,738	1,744	1,847	1,690	1,854	1,864	1,777
Male	729	781	809	855	882	879	909	903	948	920	916
White	1,986	2,103	2,206	2,294	2,406	2,410	2,534	2,380	2,572	2,557	2,455
Minority	48	50	50	50	42	31	N.D.	N.D.	34	33	32
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	65%	66%	67%	69%	66%	67%	63%	68%	66%	63%
Female	66%	68%	70%	70%	72%	70%	70%	65%	71%	70%	65%
Male	58%	60%	61%	63%	64%	61%	62%	61%	61%	60%	58%
White	62%	65%	66%	67%	68%	66%	66%	62%	67%	66%	62%
Minority	18%	18%	18%	18%	15%	11%	N.D.	N.D.	12%	11%	9%
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	187	220	231	261	281	291	323	321	341	356	354
Alcohol-related disorders	204	239	247	259	272	287	300	306	307	307	295
Anxiety disorders	1,283	1,383	1,470	1,564	1,604	1,644	1,733	1,653	1,728	1,695	1,634
Attention-deficit, conduct, and disruptive behavior disorders	77	92	107	124	138	162	176	171	197	199	221
Delirium, dementia, and amnestic and other cognitive disorders	524	519	529	502	498	467	454	427	425	387	365
Developmental disorders	226	237	250	262	266	280	290	277	274	280	292
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	33	41	45	49	49	49	52	55	57
Impulse control disorders, NEC	N.D.	32	33	43	43	46	46	48	64	65	62
Miscellaneous disorders	443	471	516	562	612	626	666	630	657	651	633
Mood disorders	1,149	1,239	1,315	1,402	1,465	1,483	1,576	1,483	1,576	1,569	1,525
Personality disorders	226	245	273	301	320	337	346	327	326	319	318
Schizophrenia and other psychotic disorders	488	501	521	518	534	532	549	513	501	476	456
Screening and history of mental health and substance abuse	1,236	1,369	1,496	1,583	1,690	1,693	1,796	1,699	1,861	1,850	1,817
Substance-related disorders	324	360	387	423	468	497	534	511	557	536	527
Suicide and intentional self-inflicted injury	55	60	61	67	76	78	91	90	90	100	101



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Grayson, Muhlenberg, Ohio, \$ Todd

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,257	5,811	6,093	6,439	6,573	6,707	6,786	6,404	6,613	6,615	6,161
Female	2,624	2,913	2,958	3,094	3,098	3,147	3,165	2,976	3,055	3,012	2,797
Male	2,633	2,898	3,135	3,345	3,475	3,560	3,621	3,428	3,558	3,603	3,364
White	4,753	5,237	5,479	5,773	5,851	5,935	5,971	5,624	5,812	5,809	5,413
Minority	217	243	240	240	239	249	258	236	247	239	224

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	52%	54%	55%	55%	55%	55%	52%	53%	51%	46%
Female	53%	54%	54%	54%	53%	53%	52%	49%	50%	48%	43%
Male	50%	50%	54%	56%	56%	57%	57%	54%	55%	53%	49%
White	52%	52%	53%	54%	54%	54%	53%	50%	51%	49%	44%
Minority	21%	21%	23%	24%	21%	21%	22%	20%	22%	19%	19%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,344	1,457	1,554	1,618	1,633	1,624	1,636	1,500	1,544	1,505	1,353
Alcohol-related disorders	164	160	164	161	145	145	129	108	104	95	73
Anxiety disorders	1,488	1,631	1,739	1,831	1,850	1,858	1,879	1,764	1,762	1,720	1,552
Attention-deficit, conduct, and disruptive behavior disorders	2,441	2,730	2,918	3,145	3,243	3,329	3,306	3,082	3,153	3,095	2,870
Delirium, dementia, and amnestic and other cognitive disorders	100	104	105	103	107	104	101	89	89	85	72
Developmental disorders	989	1,158	1,319	1,514	1,669	1,771	1,894	1,887	2,004	2,077	1,987
Disorders usually diagnosed in infancy, childhood, adolescence	289	339	364	417	468	497	535	533	563	575	543
Impulse control disorders, NEC	366	392	410	411	402	399	397	344	348	320	266
Miscellaneous disorders	752	802	818	817	787	773	739	644	639	591	533
Mood disorders	2,034	2,165	2,173	2,194	2,130	2,040	1,939	1,717	1,661	1,543	1,334
Personality disorders	174	186	187	181	183	172	164	148	132	111	90
Schizophrenia and other psychotic disorders	175	176	180	179	175	173	165	148	144	131	114
Screening and history of mental health and substance abuse	3,054	3,268	3,336	3,331	3,247	3,210	3,174	2,897	2,904	2,784	2,490
Substance-related disorders	478	489	487	477	441	415	382	330	316	280	231
Suicide and intentional self-inflicted injury	113	118	125	122	115	106	101	87	93	86	74



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Grayson, Muhlenberg, Ohio, \$ Todd

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,402	5,783	6,215	6,544	6,760	6,976	7,226	6,626	7,269	7,343	7,052
Female	3,850	4,094	4,392	4,604	4,765	4,890	5,087	4,584	5,050	5,100	4,820
Male	1,552	1,689	1,823	1,940	1,995	2,086	2,139	2,042	2,219	2,243	2,232
White	4,855	5,178	5,578	5,891	6,091	6,314	6,552	5,974	6,620	6,646	6,367
Minority	272	305	314	316	295	274	242	212	211	233	220

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	63%	65%	66%	67%	68%	68%	63%	68%	67%	62%
Female	63%	66%	69%	69%	70%	70%	72%	65%	70%	70%	65%
Male	53%	55%	57%	60%	60%	62%	62%	58%	62%	60%	57%
White	60%	63%	65%	65%	67%	67%	68%	62%	67%	66%	62%
Minority	30%	31%	32%	33%	31%	28%	25%	22%	23%	24%	22%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	590	644	734	783	793	843	890	835	909	929	921
Alcohol-related disorders	444	487	528	600	607	614	634	567	616	626	591
Anxiety disorders	2,736	2,945	3,214	3,506	3,707	3,827	4,048	3,838	4,181	4,228	4,113
Attention-deficit, conduct, and disruptive behavior disorders	270	294	330	398	450	519	605	611	709	764	821
Delirium, dementia, and amnesic and other cognitive disorders	1,336	1,332	1,358	1,315	1,263	1,235	1,238	1,142	1,165	1,082	1,015
Developmental disorders	418	434	459	475	480	504	524	509	528	528	545
Disorders usually diagnosed in infancy, childhood, adolescence	41	50	54	61	66	76	80	81	91	91	111
Impulse control disorders, NEC	82	91	98	110	108	123	145	147	156	159	184
Miscellaneous disorders	1,020	1,133	1,241	1,377	1,502	1,571	1,664	1,519	1,637	1,641	1,594
Mood disorders	3,002	3,297	3,620	3,898	4,118	4,299	4,528	4,248	4,657	4,787	4,639
Personality disorders	302	340	356	379	398	420	453	442	466	470	436
Schizophrenia and other psychotic disorders	1,119	1,161	1,203	1,232	1,236	1,254	1,294	1,198	1,226	1,209	1,160
Screening and history of mental health and substance abuse	2,727	3,014	3,371	3,670	3,901	4,114	4,306	3,932	4,319	4,391	4,199
Substance-related disorders	668	750	868	944	1,021	1,055	1,130	1,027	1,126	1,138	1,065
Suicide and intentional self-inflicted injury	134	153	180	192	223	235	268	259	291	303	296



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Greenup

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,695	1,922	2,011	2,073	2,146	2,159	2,179	2,050	2,133	2,105	1,896
Female	835	947	999	991	1,030	998	1,005	931	969	947	840
Male	860	975	1,012	1,082	1,116	1,161	1,174	1,119	1,164	1,158	1,056
White	1,611	1,819	1,883	1,932	1,981	1,981	1,981	1,865	1,926	1,895	1,710
Minority	N.D.	N.D.	32	N.D.	N.D.	32	35	N.D.	38	34	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	45%	48%	50%	50%	51%	51%	52%	48%	50%	48%	43%
Female	46%	49%	51%	49%	50%	49%	50%	46%	47%	45%	39%
Male	44%	48%	49%	51%	51%	53%	54%	51%	52%	52%	46%
White	44%	48%	49%	49%	49%	49%	50%	47%	48%	46%	41%
Minority	N.D.	N.D.	18%	N.D.	N.D.	13%	14%	N.D.	14%	12%	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	309	344	373	391	394	397	390	368	372	356	291
Alcohol-related disorders	89	81	79	73	69	62	53	50	45	39	N.D.
Anxiety disorders	514	568	566	561	565	558	547	519	530	523	467
Attention-deficit, conduct, and disruptive behavior disorders	766	866	927	984	1,058	1,082	1,076	1,020	1,043	1,027	925
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	282	324	388	426	464	494	522	522	584	589	551
Disorders usually diagnosed in infancy, childhood, adolescence	124	145	162	176	197	202	208	196	203	194	175
Impulse control disorders, NEC	59	62	59	57	57	54	52	47	43	36	34
Miscellaneous disorders	180	195	208	209	211	207	201	181	189	178	157
Mood disorders	687	729	721	707	695	645	613	540	518	483	391
Personality disorders	80	76	76	69	64	56	51	44	41	35	N.D.
Schizophrenia and other psychotic disorders	74	71	66	60	57	52	50	48	51	43	32
Screening and history of mental health and substance abuse	1,276	1,429	1,484	1,494	1,544	1,538	1,501	1,369	1,381	1,327	1,168
Substance-related disorders	230	241	216	197	194	164	145	128	133	116	99
Suicide and intentional self-inflicted injury	54	56	56	56	51	42	42	43	46	38	34



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Greenup

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,175	2,388	2,514	2,674	2,728	2,715	2,801	2,531	2,671	2,638	2,485
Female	1,531	1,651	1,763	1,861	1,883	1,891	1,947	1,760	1,842	1,801	1,692
Male	644	737	751	813	845	824	854	771	829	837	793
White	1,985	2,185	2,307	2,469	2,530	2,511	2,587	2,329	2,455	2,416	2,268
Minority	58	61	59	52	42	39	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	64%	64%	66%	65%	65%	67%	61%	66%	66%	61%
Female	64%	67%	68%	69%	68%	68%	70%	63%	68%	68%	64%
Male	56%	59%	57%	60%	60%	59%	62%	57%	61%	61%	56%
White	61%	63%	64%	65%	65%	64%	66%	60%	65%	65%	60%
Minority	21%	22%	21%	20%	16%	15%	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	148	152	174	191	201	218	248	213	233	245	229
Alcohol-related disorders	242	267	283	300	314	312	341	292	309	285	280
Anxiety disorders	1,155	1,309	1,390	1,494	1,551	1,572	1,648	1,522	1,611	1,566	1,468
Attention-deficit, conduct, and disruptive behavior disorders	184	204	216	249	258	270	300	270	310	314	302
Delirium, dementia, and amnestic and other cognitive disorders	558	582	589	589	572	562	570	528	511	464	408
Developmental disorders	264	281	296	312	313	321	322	297	304	281	265
Disorders usually diagnosed in infancy, childhood, adolescence	32	32	36	40	41	45	50	51	55	54	65
Impulse control disorders, NEC	43	52	56	60	66	66	70	65	65	65	62
Miscellaneous disorders	399	432	460	498	515	531	573	514	525	504	461
Mood disorders	1,242	1,422	1,531	1,644	1,701	1,743	1,819	1,655	1,748	1,714	1,613
Personality disorders	334	367	384	408	426	424	423	365	364	333	315
Schizophrenia and other psychotic disorders	596	638	649	671	668	655	686	623	623	602	544
Screening and history of mental health and substance abuse	1,250	1,446	1,566	1,729	1,781	1,749	1,818	1,627	1,731	1,725	1,647
Substance-related disorders	358	422	454	510	539	581	641	582	632	604	559
Suicide and intentional self-inflicted injury	86	98	101	114	114	130	150	137	150	134	130



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Hardin

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,454	3,959	4,305	4,510	4,596	4,713	4,905	4,615	4,869	5,028	4,703
Female	1,621	1,843	1,969	2,042	2,069	2,103	2,209	2,077	2,216	2,232	2,084
Male	1,833	2,116	2,336	2,468	2,527	2,610	2,696	2,538	2,653	2,796	2,619
White	2,592	2,971	3,230	3,366	3,425	3,483	3,592	3,368	3,550	3,677	3,423
Minority	633	711	760	779	756	783	813	748	800	826	789

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	47%	49%	51%	52%	52%	51%	53%	49%	49%	47%	43%
Female	45%	47%	49%	50%	49%	48%	49%	45%	46%	43%	39%
Male	48%	51%	53%	54%	55%	55%	56%	52%	53%	51%	46%
White	47%	50%	52%	53%	52%	51%	52%	48%	49%	47%	42%
Minority	33%	34%	34%	34%	32%	32%	33%	30%	31%	29%	27%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	826	963	1,073	1,138	1,166	1,189	1,238	1,148	1,206	1,260	1,190
Alcohol-related disorders	158	173	169	169	157	155	139	117	108	96	67
Anxiety disorders	979	1,101	1,204	1,231	1,244	1,267	1,337	1,269	1,340	1,339	1,239
Attention-deficit, conduct, and disruptive behavior disorders	1,957	2,255	2,451	2,571	2,695	2,756	2,872	2,679	2,762	2,755	2,551
Delirium, dementia, and amnestic and other cognitive disorders	48	53	59	58	59	60	64	56	61	61	55
Developmental disorders	724	882	1,026	1,156	1,248	1,350	1,452	1,467	1,614	1,708	1,651
Disorders usually diagnosed in infancy, childhood, adolescence	340	395	442	470	497	527	555	545	574	585	553
Impulse control disorders, NEC	453	489	497	515	514	499	493	440	416	381	320
Miscellaneous disorders	382	403	412	421	422	423	414	395	395	384	336
Mood disorders	1,291	1,436	1,469	1,458	1,438	1,372	1,360	1,249	1,211	1,135	978
Personality disorders	295	304	314	317	322	305	301	270	255	229	201
Schizophrenia and other psychotic disorders	140	135	155	155	155	147	160	143	141	136	120
Screening and history of mental health and substance abuse	2,302	2,555	2,686	2,684	2,557	2,443	2,413	2,125	2,101	2,083	1,921
Substance-related disorders	337	357	370	368	367	351	326	271	253	227	176
Suicide and intentional self-inflicted injury	74	83	84	83	86	84	85	84	84	91	81



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Hardin

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,895	3,198	3,439	3,656	3,871	3,966	4,087	3,770	4,164	4,278	4,036
Female	2,148	2,393	2,558	2,714	2,892	2,949	3,004	2,739	3,005	3,099	2,899
Male	747	805	881	942	979	1,017	1,083	1,031	1,159	1,179	1,137
White	2,330	2,566	2,748	2,927	3,127	3,210	3,335	3,094	3,416	3,498	3,282
Minority	409	462	510	536	533	530	520	447	502	532	498

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	58%	59%	61%	62%	62%	63%	58%	62%	61%	57%
Female	56%	60%	60%	63%	64%	64%	66%	59%	64%	63%	58%
Male	51%	53%	54%	56%	56%	56%	58%	55%	59%	57%	54%
White	56%	60%	60%	62%	63%	63%	65%	59%	64%	63%	58%
Minority	35%	39%	40%	41%	41%	39%	40%	35%	37%	38%	34%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	254	296	317	344	369	392	429	411	440	464	435
Alcohol-related disorders	242	291	334	351	365	382	407	366	389	395	374
Anxiety disorders	1,368	1,530	1,686	1,832	1,964	2,021	2,167	2,058	2,243	2,278	2,181
Attention-deficit, conduct, and disruptive behavior disorders	200	234	261	313	368	381	432	427	521	591	587
Delirium, dementia, and amnesic and other cognitive disorders	679	698	705	694	700	689	702	658	674	657	596
Developmental disorders	307	327	337	373	385	386	401	402	437	466	486
Disorders usually diagnosed in infancy, childhood, adolescence	55	61	66	76	80	79	95	93	106	125	137
Impulse control disorders, NEC	66	80	97	104	121	132	142	149	177	170	175
Miscellaneous disorders	616	709	778	875	914	963	1,002	921	953	963	906
Mood disorders	1,741	1,975	2,129	2,276	2,428	2,510	2,651	2,478	2,753	2,807	2,648
Personality disorders	357	400	431	463	470	480	496	462	514	531	517
Schizophrenia and other psychotic disorders	698	758	787	798	809	817	833	802	840	823	759
Screening and history of mental health and substance abuse	1,414	1,618	1,823	2,017	2,201	2,216	2,257	2,050	2,230	2,279	2,119
Substance-related disorders	362	428	460	475	511	546	592	532	579	600	567
Suicide and intentional self-inflicted injury	74	87	102	107	120	126	150	144	157	141	140



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Henderson & Webster

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,059	3,276	3,501	3,687	3,808	3,817	3,881	3,651	3,751	3,793	3,672
Female	1,497	1,589	1,699	1,754	1,779	1,777	1,812	1,686	1,696	1,689	1,623
Male	1,562	1,687	1,802	1,933	2,029	2,040	2,069	1,965	2,055	2,104	2,049
White	2,462	2,616	2,776	2,932	3,016	3,026	3,085	2,896	2,952	2,994	2,888
Minority	401	423	458	468	467	452	430	393	418	415	419

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	59%	59%	62%	62%	61%	60%	60%	55%	56%	53%	49%
Female	58%	58%	60%	60%	59%	57%	57%	52%	52%	48%	44%
Male	59%	61%	63%	64%	64%	63%	64%	59%	60%	57%	53%
White	58%	59%	62%	62%	61%	59%	59%	54%	54%	52%	47%
Minority	40%	39%	40%	39%	38%	36%	35%	32%	34%	31%	31%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	479	520	566	588	585	588	599	550	529	501	461
Alcohol-related disorders	110	112	104	102	83	74	72	72	65	57	52
Anxiety disorders	902	959	1,020	1,039	1,050	1,045	1,037	930	899	849	782
Attention-deficit, conduct, and disruptive behavior disorders	1,815	1,983	2,155	2,282	2,408	2,442	2,471	2,284	2,304	2,317	2,231
Delirium, dementia, and amnestic and other cognitive disorders	35	43	49	45	53	47	54	49	47	42	39
Developmental disorders	725	852	974	1,062	1,163	1,207	1,284	1,226	1,312	1,369	1,364
Disorders usually diagnosed in infancy, childhood, adolescence	208	228	272	299	334	357	361	340	365	377	364
Impulse control disorders, NEC	274	289	306	317	334	311	307	270	243	232	218
Miscellaneous disorders	456	479	502	520	550	573	578	551	586	582	574
Mood disorders	1,336	1,387	1,447	1,451	1,445	1,389	1,355	1,202	1,130	1,047	942
Personality disorders	105	107	110	103	114	102	102	81	75	64	61
Schizophrenia and other psychotic disorders	133	127	137	138	128	130	132	107	89	81	73
Screening and history of mental health and substance abuse	1,731	1,812	1,889	1,920	1,847	1,764	1,723	1,573	1,554	1,488	1,386
Substance-related disorders	323	328	314	309	279	249	234	201	181	158	142
Suicide and intentional self-inflicted injury	110	106	110	107	109	100	108	91	87	74	73



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Henderson & Webster

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,926	3,216	3,394	3,570	3,762	3,709	3,786	3,558	3,925	4,018	3,789
Female	2,133	2,346	2,486	2,626	2,747	2,703	2,779	2,598	2,847	2,924	2,747
Male	793	870	908	944	1,015	1,006	1,007	960	1,078	1,094	1,042
White	2,462	2,704	2,824	2,985	3,141	3,117	3,187	2,992	3,316	3,396	3,160
Minority	362	394	437	438	456	422	408	376	412	416	414

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	68%	69%	69%	68%	68%	63%	68%	67%	63%
Female	66%	69%	70%	71%	71%	70%	70%	65%	71%	69%	65%
Male	59%	62%	62%	64%	65%	63%	63%	59%	63%	63%	58%
White	65%	68%	68%	69%	70%	68%	68%	63%	69%	68%	63%
Minority	45%	47%	51%	48%	49%	46%	46%	42%	44%	42%	41%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	188	205	216	231	255	274	280	263	296	321	302
Alcohol-related disorders	263	273	294	306	329	327	348	334	351	356	343
Anxiety disorders	1,589	1,791	1,908	2,037	2,164	2,150	2,232	2,142	2,349	2,398	2,280
Attention-deficit, conduct, and disruptive behavior disorders	318	373	421	492	565	592	688	704	835	919	887
Delirium, dementia, and amnesic and other cognitive disorders	666	687	681	671	661	648	632	597	605	592	536
Developmental disorders	263	277	283	294	316	330	341	356	363	379	387
Disorders usually diagnosed in infancy, childhood, adolescence	37	38	45	47	56	57	66	74	81	91	94
Impulse control disorders, NEC	57	59	67	77	81	90	100	104	121	133	125
Miscellaneous disorders	614	700	743	796	836	835	873	837	898	950	876
Mood disorders	1,864	2,096	2,242	2,376	2,514	2,483	2,561	2,435	2,660	2,723	2,610
Personality disorders	200	215	234	236	253	254	257	261	287	284	259
Schizophrenia and other psychotic disorders	522	570	594	586	609	605	620	596	631	652	637
Screening and history of mental health and substance abuse	1,472	1,697	1,831	1,951	2,089	2,071	2,137	1,991	2,213	2,224	2,076
Substance-related disorders	449	521	565	626	680	710	742	742	848	867	796
Suicide and intentional self-inflicted injury	108	123	130	139	166	168	193	202	230	252	242



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Jackson, Laurel, & Rockcastle

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	6,545	7,268	7,630	7,909	8,187	8,340	8,486	8,045	8,404	8,356	7,777
Female	3,321	3,631	3,792	3,875	3,982	4,011	4,055	3,767	3,931	3,893	3,621
Male	3,224	3,637	3,838	4,034	4,205	4,329	4,431	4,278	4,473	4,463	4,156
White	6,165	6,837	7,165	7,409	7,667	7,785	7,886	7,466	7,800	7,734	7,198
Minority	88	98	97	106	101	97	100	99	105	109	98

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	60%	62%	62%	63%	64%	64%	60%	61%	59%	54%
Female	58%	60%	62%	63%	63%	64%	63%	59%	59%	57%	52%
Male	55%	59%	61%	62%	63%	64%	64%	62%	62%	61%	56%
White	56%	58%	60%	61%	62%	63%	62%	59%	60%	58%	53%
Minority	19%	19%	18%	18%	16%	15%	15%	14%	15%	13%	12%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,305	1,415	1,470	1,505	1,530	1,510	1,475	1,313	1,284	1,240	1,094
Alcohol-related disorders	457	487	498	484	483	445	411	352	324	280	229
Anxiety disorders	3,286	3,658	3,878	4,050	4,260	4,378	4,427	4,208	4,300	4,210	3,979
Attention-deficit, conduct, and disruptive behavior disorders	2,551	2,873	3,026	3,152	3,308	3,386	3,377	3,185	3,245	3,163	2,913
Delirium, dementia, and amnestic and other cognitive disorders	108	107	101	98	91	91	89	74	81	73	62
Developmental disorders	1,590	1,858	2,083	2,280	2,541	2,682	2,774	2,634	2,812	2,858	2,727
Disorders usually diagnosed in infancy, childhood, adolescence	379	453	493	529	574	600	626	585	584	584	558
Impulse control disorders, NEC	255	260	263	259	255	238	227	203	196	184	149
Miscellaneous disorders	981	1,047	1,057	1,063	1,075	1,060	1,038	939	944	941	847
Mood disorders	2,154	2,188	2,159	2,082	2,003	1,902	1,780	1,565	1,487	1,341	1,125
Personality disorders	356	365	363	364	360	341	319	279	244	226	189
Schizophrenia and other psychotic disorders	222	225	222	232	220	208	190	179	178	161	133
Screening and history of mental health and substance abuse	3,475	3,744	3,775	3,784	3,724	3,621	3,561	3,250	3,304	3,227	2,835
Substance-related disorders	967	981	956	915	901	822	743	643	598	552	441
Suicide and intentional self-inflicted injury	206	210	205	209	193	181	167	156	153	145	123



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Jackson, Laurel, & Rockcastle

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	7,439	8,047	8,401	8,688	8,921	9,041	9,609	8,999	9,710	9,909	9,579
Female	5,096	5,451	5,683	5,856	6,011	6,041	6,441	6,029	6,512	6,622	6,334
Male	2,343	2,596	2,718	2,832	2,910	3,000	3,168	2,970	3,198	3,287	3,245
White	6,808	7,391	7,757	8,036	8,270	8,392	8,924	8,339	9,051	9,229	8,909
Minority	190	191	159	154	129	101	72	56	58	74	73

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	70%	71%	72%	72%	71%	73%	67%	72%	71%	67%
Female	71%	73%	74%	75%	75%	74%	77%	70%	75%	74%	70%
Male	61%	64%	65%	66%	66%	65%	67%	62%	66%	65%	62%
White	67%	69%	71%	71%	71%	70%	72%	66%	71%	70%	66%
Minority	23%	23%	19%	19%	16%	13%	9%	7%	7%	9%	8%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	878	1,007	1,071	1,151	1,196	1,241	1,381	1,321	1,426	1,469	1,427
Alcohol-related disorders	709	785	828	881	953	970	1,050	957	1,020	1,037	991
Anxiety disorders	4,880	5,313	5,631	5,885	6,083	6,234	6,654	6,328	6,700	6,824	6,610
Attention-deficit, conduct, and disruptive behavior disorders	340	375	431	492	559	599	661	618	720	782	827
Delirium, dementia, and amnesic and other cognitive disorders	1,517	1,553	1,517	1,478	1,428	1,382	1,412	1,266	1,247	1,174	1,061
Developmental disorders	674	715	763	798	848	867	920	875	929	941	953
Disorders usually diagnosed in infancy, childhood, adolescence	108	118	120	131	135	144	154	145	159	171	184
Impulse control disorders, NEC	177	194	211	223	244	248	266	243	266	271	280
Miscellaneous disorders	1,962	2,140	2,337	2,451	2,591	2,670	2,818	2,718	2,891	2,957	2,852
Mood disorders	4,470	4,860	5,135	5,359	5,533	5,626	5,991	5,673	6,042	6,193	6,043
Personality disorders	558	634	666	716	765	787	827	773	814	812	806
Schizophrenia and other psychotic disorders	1,448	1,537	1,564	1,574	1,582	1,577	1,648	1,512	1,549	1,513	1,416
Screening and history of mental health and substance abuse	4,282	4,729	5,072	5,332	5,576	5,717	6,164	5,823	6,370	6,547	6,369
Substance-related disorders	1,409	1,580	1,706	1,848	1,988	2,087	2,287	2,156	2,366	2,472	2,373
Suicide and intentional self-inflicted injury	225	271	289	308	330	357	429	412	455	482	459



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Jefferson

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	27,005	29,923	32,167	33,983	35,145	35,410	35,746	33,926	34,849	35,138	33,399
Female	12,855	14,031	14,990	15,689	16,109	16,033	16,048	15,164	15,477	15,462	14,503
Male	14,150	15,892	17,177	18,294	19,036	19,377	19,698	18,762	19,372	19,676	18,896
White	11,662	13,218	14,222	15,210	15,817	16,057	16,241	15,424	15,760	16,025	15,360
Minority	13,733	14,833	15,768	16,352	16,633	16,494	16,432	15,368	15,779	15,645	14,690

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	48%	50%	52%	52%	52%	50%	49%	46%	45%	43%	39%
Female	47%	48%	50%	49%	48%	47%	45%	42%	41%	39%	35%
Male	50%	52%	54%	55%	55%	54%	53%	49%	49%	47%	43%
White	48%	50%	53%	53%	52%	50%	48%	44%	43%	41%	37%
Minority	43%	44%	45%	45%	45%	43%	43%	39%	39%	36%	33%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	8,198	9,197	9,909	10,344	10,588	10,544	10,425	9,548	9,393	8,995	8,275
Alcohol-related disorders	1,056	1,077	1,056	1,012	980	914	849	730	655	574	471
Anxiety disorders	7,205	7,820	8,178	8,443	8,580	8,386	8,222	7,634	7,463	7,170	6,436
Attention-deficit, conduct, and disruptive behavior disorders	13,367	15,095	16,483	17,571	18,326	18,626	18,732	17,608	17,642	17,401	16,345
Delirium, dementia, and amnesic and other cognitive disorders	488	524	549	571	565	554	548	514	519	493	464
Developmental disorders	5,663	6,764	7,718	8,679	9,434	9,918	10,479	10,575	11,173	11,658	11,531
Disorders usually diagnosed in infancy, childhood, adolescence	2,347	2,729	3,012	3,316	3,553	3,730	3,928	3,843	3,978	3,953	3,837
Impulse control disorders, NEC	2,520	2,747	2,876	2,932	2,918	2,810	2,715	2,445	2,332	2,157	1,852
Miscellaneous disorders	2,919	3,049	3,172	3,258	3,278	3,269	3,195	2,962	2,947	2,875	2,660
Mood disorders	9,844	10,536	10,788	10,875	10,820	10,413	10,055	9,066	8,715	8,173	7,203
Personality disorders	791	836	868	878	876	842	810	744	733	667	563
Schizophrenia and other psychotic disorders	1,551	1,636	1,662	1,700	1,716	1,676	1,649	1,547	1,547	1,437	1,250
Screening and history of mental health and substance abuse	16,460	17,732	18,518	18,794	18,684	17,990	17,513	16,008	15,772	15,100	13,668
Substance-related disorders	2,936	3,042	3,010	2,933	2,851	2,700	2,515	2,229	2,166	1,985	1,679
Suicide and intentional self-inflicted injury	923	954	1,002	1,043	1,050	1,023	1,010	951	959	903	792



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Jefferson

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	22,608	24,718	26,599	28,305	29,831	30,885	31,491	30,153	32,964	34,141	33,348
Female	17,078	18,540	19,897	21,176	22,260	22,811	23,167	21,904	23,895	24,432	23,632
Male	5,530	6,178	6,702	7,129	7,571	8,074	8,324	8,249	9,069	9,709	9,716
White	12,720	13,911	14,902	15,855	16,725	17,440	17,892	16,900	18,435	18,886	18,153
Minority	8,884	9,698	10,491	11,150	11,749	12,030	12,093	11,704	12,866	13,498	13,243

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	58%	60%	62%	62%	63%	63%	59%	62%	62%	58%
Female	56%	59%	61%	63%	63%	64%	64%	59%	63%	62%	59%
Male	51%	55%	57%	59%	59%	61%	61%	59%	61%	60%	57%
White	57%	60%	62%	65%	65%	66%	65%	61%	65%	64%	60%
Minority	46%	49%	51%	52%	53%	53%	53%	50%	53%	52%	49%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,739	3,040	3,278	3,473	3,748	3,913	4,102	4,023	4,454	4,709	4,714
Alcohol-related disorders	2,667	2,999	3,254	3,484	3,690	3,856	3,974	3,873	4,116	4,177	4,006
Anxiety disorders	9,081	10,109	10,995	11,860	12,560	13,135	13,752	13,570	14,612	15,161	14,772
Attention-deficit, conduct, and disruptive behavior disorders	1,521	1,749	1,997	2,222	2,472	2,777	3,145	3,358	3,914	4,248	4,349
Delirium, dementia, and amnesic and other cognitive disorders	5,914	6,225	6,383	6,358	6,262	6,350	6,389	6,126	6,171	5,936	5,423
Developmental disorders	2,021	2,171	2,320	2,427	2,495	2,576	2,674	2,713	2,897	3,056	3,129
Disorders usually diagnosed in infancy, childhood, adolescence	290	328	360	405	449	495	564	613	696	801	878
Impulse control disorders, NEC	523	580	660	729	808	871	978	1,024	1,173	1,295	1,385
Miscellaneous disorders	3,362	3,843	4,323	4,942	5,513	5,899	6,121	5,641	6,233	6,302	5,940
Mood disorders	12,322	13,531	14,690	15,769	16,710	17,361	17,967	17,508	18,937	19,587	19,281
Personality disorders	1,333	1,476	1,586	1,618	1,686	1,760	1,854	1,817	1,904	1,944	1,878
Schizophrenia and other psychotic disorders	5,753	6,175	6,440	6,599	6,764	6,937	7,033	6,797	6,949	6,932	6,608
Screening and history of mental health and substance abuse	12,370	13,989	15,544	16,957	18,433	19,202	19,608	18,776	20,742	21,489	20,910
Substance-related disorders	3,885	4,436	4,877	5,380	5,783	6,066	6,399	6,229	6,818	7,130	6,889
Suicide and intentional self-inflicted injury	859	975	1,105	1,222	1,326	1,446	1,595	1,599	1,858	1,949	1,950



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Kenton

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,505	6,191	6,634	7,040	7,162	7,429	7,800	7,436	8,019	8,225	7,802
Female	2,652	2,966	3,156	3,299	3,298	3,403	3,552	3,352	3,641	3,739	3,493
Male	2,853	3,225	3,478	3,741	3,864	4,026	4,248	4,084	4,378	4,486	4,309
White	4,320	4,872	5,198	5,513	5,588	5,845	6,093	5,801	6,266	6,436	6,111
Minority	941	1,036	1,106	1,141	1,139	1,131	1,174	1,133	1,205	1,241	1,175

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	58%	60%	59%	58%	58%	59%	54%	55%	52%	46%
Female	56%	58%	59%	57%	56%	55%	55%	50%	52%	49%	44%
Male	57%	59%	61%	62%	60%	60%	62%	58%	59%	55%	49%
White	57%	59%	60%	60%	59%	59%	59%	54%	56%	52%	46%
Minority	45%	45%	44%	43%	40%	38%	39%	36%	37%	36%	32%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,283	1,410	1,506	1,604	1,608	1,626	1,657	1,529	1,554	1,518	1,373
Alcohol-related disorders	287	306	307	301	305	288	283	257	248	235	200
Anxiety disorders	1,666	1,813	1,870	1,895	1,860	1,880	1,892	1,730	1,760	1,713	1,563
Attention-deficit, conduct, and disruptive behavior disorders	3,117	3,507	3,873	4,217	4,390	4,601	4,875	4,656	4,869	4,890	4,646
Delirium, dementia, and amnestic and other cognitive disorders	126	121	139	147	151	151	161	160	169	153	137
Developmental disorders	1,449	1,676	1,887	2,116	2,310	2,487	2,725	2,661	2,979	3,058	2,947
Disorders usually diagnosed in infancy, childhood, adolescence	704	807	928	1,010	1,074	1,130	1,193	1,128	1,183	1,167	1,103
Impulse control disorders, NEC	319	340	357	355	361	361	347	316	297	286	252
Miscellaneous disorders	1,137	1,258	1,338	1,410	1,444	1,462	1,508	1,379	1,454	1,421	1,328
Mood disorders	2,076	2,256	2,309	2,373	2,303	2,268	2,267	2,030	1,995	1,936	1,719
Personality disorders	354	354	348	331	314	285	257	217	192	166	144
Schizophrenia and other psychotic disorders	268	272	274	293	280	286	292	263	268	275	250
Screening and history of mental health and substance abuse	3,789	4,201	4,384	4,465	4,359	4,338	4,381	4,056	4,173	4,136	3,732
Substance-related disorders	653	680	683	663	659	623	604	533	514	485	413
Suicide and intentional self-inflicted injury	240	240	240	259	264	270	276	278	296	285	245



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Kenton

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,722	5,275	5,698	6,005	6,343	6,545	6,952	6,347	7,147	7,368	6,983
Female	3,617	3,999	4,330	4,533	4,738	4,925	5,203	4,699	5,296	5,417	5,072
Male	1,105	1,276	1,368	1,472	1,605	1,620	1,749	1,648	1,851	1,951	1,911
White	4,032	4,484	4,815	5,103	5,423	5,595	5,946	5,432	6,115	6,296	5,940
Minority	515	609	686	699	704	716	747	670	773	796	739

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	69%	69%	70%	70%	73%	65%	71%	70%	65%
Female	66%	69%	71%	71%	72%	72%	74%	66%	71%	71%	66%
Male	58%	61%	63%	64%	67%	66%	69%	64%	69%	67%	63%
White	65%	68%	69%	70%	72%	72%	74%	66%	72%	72%	66%
Minority	44%	48%	51%	50%	48%	48%	49%	44%	48%	46%	41%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	654	732	831	862	908	950	1,008	957	1,016	1,081	1,042
Alcohol-related disorders	706	784	830	868	931	975	1,021	943	1,022	1,026	964
Anxiety disorders	2,592	2,882	3,143	3,356	3,542	3,681	3,963	3,714	4,043	4,185	3,993
Attention-deficit, conduct, and disruptive behavior disorders	360	424	482	596	698	782	895	898	1,053	1,151	1,118
Delirium, dementia, and amnesic and other cognitive disorders	1,004	1,136	1,180	1,213	1,226	1,253	1,258	1,197	1,227	1,153	1,049
Developmental disorders	367	406	445	455	468	484	545	569	620	640	659
Disorders usually diagnosed in infancy, childhood, adolescence	60	75	82	94	100	113	137	134	154	185	197
Impulse control disorders, NEC	72	88	100	112	122	138	149	145	170	180	180
Miscellaneous disorders	1,130	1,274	1,457	1,642	1,770	1,835	1,962	1,749	1,914	1,946	1,775
Mood disorders	2,910	3,252	3,528	3,833	4,076	4,245	4,571	4,265	4,720	4,883	4,611
Personality disorders	561	638	688	762	834	880	1,018	909	956	921	823
Schizophrenia and other psychotic disorders	1,028	1,151	1,194	1,245	1,290	1,356	1,415	1,360	1,435	1,416	1,332
Screening and history of mental health and substance abuse	3,031	3,433	3,822	4,100	4,346	4,511	4,880	4,434	5,054	5,256	4,963
Substance-related disorders	936	1,074	1,156	1,278	1,370	1,494	1,603	1,474	1,652	1,751	1,674
Suicide and intentional self-inflicted injury	280	310	350	378	398	440	526	501	554	613	576



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Knox & Whitley

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	7,216	7,815	8,167	8,666	8,901	8,969	9,005	8,504	8,764	8,572	8,043
Female	3,630	3,916	4,021	4,253	4,332	4,315	4,321	4,073	4,200	4,089	3,822
Male	3,586	3,899	4,146	4,413	4,569	4,654	4,684	4,431	4,564	4,483	4,221
White	6,671	7,204	7,511	7,955	8,152	8,187	8,181	7,721	7,958	7,776	7,289
Minority	79	90	90	88	94	97	105	104	114	108	98

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	62%	64%	66%	66%	68%	68%	64%	65%	62%	58%
Female	62%	64%	65%	67%	66%	67%	67%	63%	63%	60%	56%
Male	57%	60%	63%	65%	66%	68%	68%	65%	66%	64%	59%
White	58%	61%	63%	65%	65%	66%	66%	62%	63%	61%	57%
Minority	12%	12%	11%	10%	11%	11%	11%	11%	12%	10%	10%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,376	1,455	1,530	1,623	1,639	1,611	1,552	1,412	1,403	1,300	1,174
Alcohol-related disorders	676	709	710	714	705	657	608	519	489	427	369
Anxiety disorders	3,756	4,096	4,358	4,713	4,925	5,015	5,105	4,879	5,005	4,912	4,647
Attention-deficit, conduct, and disruptive behavior disorders	3,020	3,256	3,495	3,702	3,764	3,741	3,656	3,392	3,390	3,227	2,989
Delirium, dementia, and amnestic and other cognitive disorders	114	126	123	118	113	105	98	90	87	86	73
Developmental disorders	1,943	2,155	2,327	2,524	2,655	2,703	2,699	2,535	2,575	2,478	2,320
Disorders usually diagnosed in infancy, childhood, adolescence	615	692	746	810	876	898	895	836	851	813	773
Impulse control disorders, NEC	352	368	370	376	381	367	368	324	310	282	244
Miscellaneous disorders	1,387	1,416	1,398	1,405	1,385	1,328	1,302	1,213	1,196	1,110	1,003
Mood disorders	2,467	2,477	2,441	2,419	2,312	2,180	2,012	1,765	1,642	1,453	1,250
Personality disorders	518	524	529	549	547	520	490	425	383	344	314
Schizophrenia and other psychotic disorders	293	296	288	285	274	265	246	214	201	172	151
Screening and history of mental health and substance abuse	3,689	3,854	3,904	3,972	3,924	3,797	3,705	3,401	3,409	3,238	2,955
Substance-related disorders	1,312	1,320	1,310	1,298	1,246	1,117	1,005	868	800	718	601
Suicide and intentional self-inflicted injury	209	215	212	195	181	171	165	153	145	124	112



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Knox & Whitley

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	8,855	9,422	10,006	10,400	10,580	10,670	11,002	10,678	11,349	11,413	11,051
Female	5,969	6,303	6,630	6,849	6,950	6,974	7,154	6,947	7,393	7,386	7,088
Male	2,886	3,119	3,376	3,551	3,630	3,696	3,848	3,731	3,956	4,027	3,963
White	8,102	8,628	9,174	9,546	9,741	9,829	10,127	9,830	10,480	10,539	10,172
Minority	184	182	175	171	144	125	101	88	84	93	91

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	69%	71%	72%	73%	73%	73%	75%	72%	74%	74%	71%
Female	72%	75%	76%	76%	76%	76%	78%	75%	78%	77%	73%
Male	63%	64%	66%	68%	67%	68%	70%	67%	69%	68%	66%
White	68%	70%	72%	72%	72%	73%	74%	71%	74%	74%	70%
Minority	19%	18%	17%	16%	14%	12%	10%	8%	8%	8%	8%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	866	924	1,002	1,051	1,123	1,181	1,251	1,254	1,370	1,402	1,342
Alcohol-related disorders	1,016	1,070	1,149	1,234	1,280	1,300	1,365	1,293	1,355	1,355	1,326
Anxiety disorders	6,119	6,551	7,040	7,353	7,545	7,670	7,934	7,785	8,202	8,198	7,991
Attention-deficit, conduct, and disruptive behavior disorders	358	415	463	532	631	698	777	821	927	995	995
Delirium, dementia, and amnesic and other cognitive disorders	1,677	1,719	1,714	1,700	1,652	1,613	1,591	1,491	1,432	1,345	1,225
Developmental disorders	723	759	802	847	870	908	983	980	1,023	1,090	1,100
Disorders usually diagnosed in infancy, childhood, adolescence	93	98	105	113	115	129	142	153	170	182	192
Impulse control disorders, NEC	174	192	202	217	233	244	258	259	282	291	296
Miscellaneous disorders	2,842	3,093	3,298	3,530	3,667	3,763	3,887	3,814	3,997	3,989	3,814
Mood disorders	5,108	5,474	5,845	6,161	6,330	6,425	6,719	6,546	6,915	6,904	6,701
Personality disorders	786	841	884	959	1,007	1,036	1,086	1,064	1,113	1,115	1,066
Schizophrenia and other psychotic disorders	1,795	1,884	1,913	1,966	1,950	1,973	2,034	1,953	1,922	1,838	1,709
Screening and history of mental health and substance abuse	4,832	5,243	5,693	6,055	6,257	6,410	6,769	6,598	7,106	7,222	7,057
Substance-related disorders	2,104	2,294	2,506	2,690	2,820	2,911	3,085	2,996	3,226	3,291	3,214
Suicide and intentional self-inflicted injury	319	353	380	427	452	475	539	534	568	562	540



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: McCracken

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,300	3,593	3,840	3,902	3,952	3,901	3,783	3,438	3,578	3,528	3,218
Female	1,594	1,728	1,841	1,846	1,829	1,779	1,721	1,562	1,630	1,590	1,437
Male	1,706	1,865	1,999	2,056	2,123	2,122	2,062	1,876	1,948	1,938	1,781
White	2,136	2,358	2,501	2,542	2,588	2,521	2,448	2,221	2,323	2,324	2,107
Minority	974	1,014	1,083	1,078	1,054	1,045	982	903	919	883	794

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	58%	59%	59%	58%	56%	54%	50%	51%	49%	43%
Female	53%	56%	57%	57%	54%	52%	50%	46%	47%	44%	40%
Male	55%	60%	61%	61%	61%	59%	58%	54%	55%	54%	47%
White	56%	59%	60%	59%	58%	55%	54%	50%	50%	49%	42%
Minority	44%	47%	47%	46%	44%	43%	40%	38%	38%	37%	33%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	348	388	395	375	372	370	378	342	348	327	273
Alcohol-related disorders	103	99	100	89	89	80	87	71	69	55	45
Anxiety disorders	714	734	763	743	758	729	708	622	602	579	514
Attention-deficit, conduct, and disruptive behavior disorders	1,774	1,969	2,113	2,178	2,307	2,292	2,245	2,011	2,049	1,995	1,821
Delirium, dementia, and amnestic and other cognitive disorders	45	40	42	50	45	45	43	44	48	46	41
Developmental disorders	689	804	927	1,022	1,076	1,129	1,175	1,090	1,183	1,208	1,154
Disorders usually diagnosed in infancy, childhood, adolescence	202	226	241	281	305	317	332	297	312	313	301
Impulse control disorders, NEC	235	244	248	239	257	245	238	216	199	175	156
Miscellaneous disorders	407	419	417	394	400	390	368	334	335	320	262
Mood disorders	1,156	1,191	1,224	1,186	1,220	1,159	1,124	986	969	937	832
Personality disorders	89	83	78	71	68	60	60	52	53	42	33
Schizophrenia and other psychotic disorders	117	106	111	107	110	110	108	95	102	91	79
Screening and history of mental health and substance abuse	2,111	2,299	2,401	2,342	2,213	2,060	1,951	1,731	1,731	1,642	1,444
Substance-related disorders	255	255	250	238	216	196	196	163	156	133	121
Suicide and intentional self-inflicted injury	107	108	107	98	100	91	88	79	74	75	68




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: McCracken

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,353	3,660	3,875	3,997	4,025	4,107	4,171	3,804	4,089	4,103	3,895
Female	2,579	2,787	2,928	3,021	3,046	3,102	3,147	2,826	3,009	3,005	2,808
Male	774	873	947	976	979	1,005	1,024	978	1,080	1,098	1,087
White	2,380	2,615	2,763	2,877	2,897	2,953	3,012	2,740	2,944	2,971	2,750
Minority	825	878	934	932	932	947	928	831	901	888	879
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	66%	68%	69%	69%	69%	70%	66%	71%	69%	64%
Female	63%	67%	69%	70%	70%	71%	73%	67%	72%	71%	65%
Male	57%	60%	63%	65%	64%	63%	63%	61%	66%	65%	63%
White	62%	66%	68%	70%	69%	69%	71%	65%	71%	70%	64%
Minority	51%	55%	56%	56%	55%	56%	55%	51%	55%	52%	50%
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	197	216	243	248	257	272	283	268	276	298	278
Alcohol-related disorders	358	400	441	485	491	492	492	455	487	481	465
Anxiety disorders	1,711	1,888	2,004	2,149	2,204	2,270	2,349	2,214	2,312	2,339	2,217
Attention-deficit, conduct, and disruptive behavior disorders	150	170	195	236	249	294	334	321	381	462	480
Delirium, dementia, and amnestic and other cognitive disorders	767	794	793	751	721	718	723	682	706	674	618
Developmental disorders	257	271	275	286	277	288	304	299	320	338	324
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	32	35	40	40	45	51	57
Impulse control disorders, NEC	48	49	59	66	68	83	98	90	85	117	115
Miscellaneous disorders	662	757	809	874	922	961	1,013	920	984	1,011	948
Mood disorders	1,887	2,098	2,244	2,393	2,439	2,532	2,622	2,421	2,602	2,640	2,514
Personality disorders	254	278	308	320	333	338	360	349	379	401	397
Schizophrenia and other psychotic disorders	592	628	641	661	674	682	689	648	672	688	663
Screening and history of mental health and substance abuse	1,910	2,182	2,402	2,510	2,559	2,616	2,636	2,370	2,523	2,530	2,407
Substance-related disorders	467	536	562	632	654	683	713	650	703	713	698
Suicide and intentional self-inflicted injury	144	169	183	206	230	247	292	273	332	341	327



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Mason

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	943	1,026	1,123	1,130	1,162	1,177	1,212	1,150	1,245	1,246	1,149
Female	465	490	540	535	552	558	571	544	578	571	506
Male	478	536	583	595	610	619	641	606	667	675	643
White	787	859	943	943	968	980	999	941	1,019	1,020	940
Minority	108	111	112	117	117	119	124	122	141	138	133

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	57%	60%	59%	59%	56%	56%	53%	54%	49%	44%
Female	58%	58%	60%	57%	56%	55%	53%	52%	53%	47%	40%
Male	56%	56%	61%	60%	61%	58%	59%	53%	55%	51%	47%
White	57%	58%	61%	58%	58%	56%	56%	51%	52%	48%	43%
Minority	38%	36%	35%	38%	37%	35%	34%	34%	38%	33%	31%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	204	219	241	236	242	255	267	274	302	291	282
Alcohol-related disorders	78	77	75	64	65	53	51	37	39	38	N.D.
Anxiety disorders	310	320	351	352	354	355	353	330	332	319	285
Attention-deficit, conduct, and disruptive behavior disorders	457	512	554	566	564	565	582	526	551	546	494
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	299	337	384	403	440	465	487	471	522	530	502
Disorders usually diagnosed in infancy, childhood, adolescence	84	89	109	115	120	123	126	122	126	130	117
Impulse control disorders, NEC	N.D.	N.D.	32	33	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	128	135	140	129	129	121	124	112	129	132	115
Mood disorders	341	346	388	369	355	341	330	293	288	260	215
Personality disorders	71	82	86	90	96	96	88	67	54	46	34
Schizophrenia and other psychotic disorders	50	48	47	51	51	54	55	47	45	45	37
Screening and history of mental health and substance abuse	699	748	820	813	816	789	790	727	750	724	652
Substance-related disorders	130	126	130	121	114	106	97	76	73	65	47
Suicide and intentional self-inflicted injury	N.D.	N.D.	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Mason

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,017	1,093	1,142	1,230	1,269	1,323	1,391	1,276	1,414	1,483	1,383
Female	770	813	841	892	914	960	1,012	913	1,007	1,049	955
Male	247	280	301	338	355	363	379	363	407	434	428
White	891	952	1,010	1,075	1,113	1,175	1,214	1,121	1,245	1,310	1,205
Minority	92	105	96	116	114	106	123	106	118	116	116
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	64%	63%	66%	68%	67%	68%	62%	68%	68%	63%
Female	67%	68%	68%	70%	71%	71%	72%	65%	71%	71%	64%
Male	52%	55%	54%	57%	61%	60%	60%	56%	62%	62%	60%
White	64%	65%	65%	67%	68%	68%	68%	63%	68%	69%	64%
Minority	38%	42%	40%	44%	48%	45%	46%	42%	49%	40%	38%
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	94	98	112	132	139	150	166	148	173	187	176
Alcohol-related disorders	95	99	111	142	138	140	153	137	144	162	160
Anxiety disorders	548	584	647	685	704	724	779	722	779	823	789
Attention-deficit, conduct, and disruptive behavior disorders	86	88	101	127	142	154	160	163	180	191	185
Delirium, dementia, and amnestic and other cognitive disorders	206	206	198	201	204	201	195	189	190	174	162
Developmental disorders	102	113	117	117	134	137	143	139	149	151	154
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	33	N.D.	34	34	35	38	35
Miscellaneous disorders	186	207	227	242	259	275	276	269	293	310	291
Mood disorders	547	581	632	685	707	730	769	712	781	819	793
Personality disorders	127	146	161	175	177	187	199	180	195	210	197
Schizophrenia and other psychotic disorders	195	212	220	235	238	228	234	226	226	235	220
Screening and history of mental health and substance abuse	656	725	770	860	931	974	1,006	914	1,037	1,100	1,014
Substance-related disorders	140	147	172	202	217	240	253	229	282	283	279
Suicide and intentional self-inflicted injury	N.D.	33	33	36	41	45	51	46	48	53	53



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Metcalfe, Monroe, Taylor & Wayne

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,412	4,692	4,977	5,206	5,253	5,292	5,400	5,060	5,315	5,267	4,949
Female	2,221	2,310	2,430	2,486	2,502	2,504	2,544	2,358	2,492	2,456	2,292
Male	2,191	2,382	2,547	2,720	2,751	2,788	2,856	2,702	2,823	2,811	2,657
White	3,983	4,229	4,463	4,650	4,668	4,706	4,779	4,471	4,696	4,661	4,370
Minority	219	228	233	238	233	222	221	215	224	204	189

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	57%	60%	61%	60%	60%	62%	56%	57%	54%	50%
Female	57%	58%	60%	60%	60%	59%	60%	54%	55%	52%	48%
Male	54%	56%	60%	62%	61%	61%	63%	59%	59%	57%	52%
White	55%	56%	59%	60%	59%	59%	60%	55%	56%	53%	48%
Minority	34%	33%	32%	31%	28%	27%	27%	25%	26%	22%	20%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	907	955	1,034	1,078	1,079	1,065	1,070	952	958	929	837
Alcohol-related disorders	173	181	178	170	144	120	112	90	77	67	53
Anxiety disorders	2,022	2,158	2,265	2,372	2,401	2,435	2,495	2,355	2,420	2,362	2,191
Attention-deficit, conduct, and disruptive behavior disorders	1,860	2,002	2,140	2,263	2,299	2,289	2,321	2,100	2,163	2,125	1,971
Delirium, dementia, and amnestic and other cognitive disorders	63	70	73	74	72	72	67	60	59	55	50
Developmental disorders	974	1,106	1,269	1,409	1,502	1,583	1,674	1,616	1,759	1,790	1,725
Disorders usually diagnosed in infancy, childhood, adolescence	306	359	401	444	460	454	480	452	459	444	412
Impulse control disorders, NEC	289	312	331	330	315	294	283	237	229	218	191
Miscellaneous disorders	648	667	675	650	619	584	559	488	489	468	419
Mood disorders	1,742	1,753	1,748	1,713	1,621	1,550	1,483	1,278	1,238	1,170	1,014
Personality disorders	127	121	116	110	99	89	78	65	57	58	48
Schizophrenia and other psychotic disorders	154	150	146	151	145	137	131	118	119	107	93
Screening and history of mental health and substance abuse	2,508	2,602	2,653	2,647	2,555	2,441	2,387	2,115	2,153	2,034	1,839
Substance-related disorders	463	440	432	404	358	314	288	228	216	197	171
Suicide and intentional self-inflicted injury	135	130	123	122	106	100	94	79	81	83	74



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Metcalfe, Monroe, Taylor & Wayne

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,979	6,281	6,496	6,689	6,775	6,857	7,031	6,586	6,956	7,090	6,790
Female	3,995	4,181	4,297	4,420	4,478	4,513	4,654	4,391	4,623	4,690	4,433
Male	1,984	2,100	2,199	2,269	2,297	2,344	2,377	2,195	2,333	2,400	2,357
White	5,253	5,530	5,726	5,927	6,024	6,104	6,259	5,839	6,202	6,311	6,023
Minority	389	388	397	363	337	310	296	279	299	308	304

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	70%	71%	72%	71%	71%	72%	67%	70%	69%	66%
Female	70%	73%	74%	75%	74%	73%	75%	70%	73%	72%	67%
Male	63%	65%	67%	66%	67%	68%	68%	63%	65%	65%	62%
White	67%	70%	71%	71%	71%	71%	72%	66%	69%	69%	65%
Minority	38%	37%	38%	36%	33%	31%	29%	28%	30%	28%	27%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	570	640	681	740	736	754	790	767	828	849	806
Alcohol-related disorders	596	662	685	701	740	734	747	704	734	729	690
Anxiety disorders	3,733	3,987	4,212	4,377	4,523	4,624	4,813	4,580	4,780	4,878	4,743
Attention-deficit, conduct, and disruptive behavior disorders	206	227	256	284	330	374	435	459	539	604	588
Delirium, dementia, and amnesic and other cognitive disorders	1,420	1,453	1,421	1,365	1,289	1,255	1,206	1,111	1,064	981	897
Developmental disorders	529	545	577	585	601	614	628	598	631	642	654
Disorders usually diagnosed in infancy, childhood, adolescence	44	48	55	51	51	58	63	62	71	84	80
Impulse control disorders, NEC	107	121	141	161	169	190	202	189	203	211	214
Miscellaneous disorders	1,458	1,522	1,617	1,724	1,774	1,825	1,888	1,772	1,852	1,865	1,753
Mood disorders	3,670	3,916	4,123	4,353	4,449	4,522	4,691	4,446	4,641	4,735	4,552
Personality disorders	475	500	524	550	572	584	599	551	561	572	537
Schizophrenia and other psychotic disorders	1,389	1,441	1,481	1,457	1,421	1,408	1,424	1,347	1,326	1,254	1,166
Screening and history of mental health and substance abuse	3,100	3,351	3,564	3,726	3,830	3,905	4,043	3,829	4,078	4,191	4,035
Substance-related disorders	913	1,002	1,061	1,132	1,198	1,232	1,269	1,208	1,294	1,344	1,282
Suicide and intentional self-inflicted injury	182	215	235	242	267	284	321	306	329	336	312



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Montgomery

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,544	1,666	1,788	1,821	1,861	1,903	1,985	1,869	1,988	1,990	1,827
Female	800	833	901	898	888	902	929	867	910	903	818
Male	744	833	887	923	973	1,001	1,056	1,002	1,078	1,087	1,009
White	1,420	1,525	1,636	1,660	1,685	1,729	1,786	1,673	1,778	1,781	1,634
Minority	48	58	58	50	53	51	52	49	52	57	50

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	60%	62%	63%	63%	62%	62%	57%	58%	55%	48%
Female	59%	60%	62%	63%	62%	60%	60%	55%	55%	53%	46%
Male	54%	61%	62%	63%	64%	62%	64%	59%	60%	56%	50%
White	56%	59%	61%	62%	62%	60%	61%	55%	57%	53%	47%
Minority	24%	29%	27%	22%	22%	22%	20%	18%	18%	19%	17%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	327	343	385	410	421	442	443	387	397	394	349
Alcohol-related disorders	88	78	83	79	73	66	60	44	43	35	N.D.
Anxiety disorders	636	692	766	765	797	799	835	796	844	840	788
Attention-deficit, conduct, and disruptive behavior disorders	566	620	672	715	733	741	778	708	734	714	639
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	302	334	374	406	451	478	530	528	570	576	544
Disorders usually diagnosed in infancy, childhood, adolescence	160	175	201	224	238	241	258	247	258	254	234
Impulse control disorders, NEC	46	50	57	55	55	52	55	45	43	38	N.D.
Miscellaneous disorders	225	236	253	243	250	256	276	261	266	266	240
Mood disorders	626	634	662	650	646	616	611	526	522	487	420
Personality disorders	56	57	51	50	47	42	44	34	33	32	N.D.
Schizophrenia and other psychotic disorders	67	63	67	70	72	66	70	63	63	59	49
Screening and history of mental health and substance abuse	1,193	1,267	1,354	1,356	1,342	1,326	1,349	1,199	1,232	1,205	1,059
Substance-related disorders	221	208	215	201	193	173	180	137	131	111	87
Suicide and intentional self-inflicted injury	67	66	68	66	67	64	64	56	60	59	46



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Montgomery

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,586	1,713	1,818	1,869	1,920	1,943	1,980	1,895	2,100	2,087	2,048
Female	1,159	1,250	1,313	1,356	1,404	1,420	1,468	1,367	1,534	1,505	1,464
Male	427	463	505	513	516	523	512	528	566	582	584
White	1,425	1,543	1,645	1,690	1,737	1,761	1,791	1,720	1,920	1,912	1,863
Minority	78	83	79	77	76	72	68	58	61	56	58

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	69%	71%	71%	69%	70%	66%	70%	68%	65%
Female	68%	71%	72%	74%	74%	72%	74%	68%	73%	71%	67%
Male	57%	58%	61%	63%	64%	63%	62%	62%	64%	61%	60%
White	65%	67%	69%	71%	71%	70%	70%	66%	71%	68%	65%
Minority	31%	33%	31%	29%	29%	26%	26%	23%	23%	21%	21%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	149	164	177	189	208	212	228	221	246	234	250
Alcohol-related disorders	184	202	213	220	232	219	233	223	238	227	219
Anxiety disorders	917	1,010	1,056	1,120	1,169	1,192	1,249	1,180	1,288	1,296	1,284
Attention-deficit, conduct, and disruptive behavior disorders	59	71	75	88	101	108	130	137	148	173	182
Delirium, dementia, and amnesic and other cognitive disorders	350	344	332	313	305	305	296	275	287	264	225
Developmental disorders	141	157	171	169	179	187	204	203	205	218	224
Disorders usually diagnosed in infancy, childhood, adolescence	33	39	38	35	34	35	46	48	52	58	58
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	36	38	40	38	35	38	41
Miscellaneous disorders	409	443	469	496	519	554	575	539	580	594	580
Mood disorders	924	1,021	1,095	1,148	1,190	1,214	1,276	1,201	1,316	1,307	1,302
Personality disorders	145	174	189	196	205	198	214	193	195	192	194
Schizophrenia and other psychotic disorders	331	326	344	335	343	348	360	339	326	307	296
Screening and history of mental health and substance abuse	1,003	1,133	1,259	1,301	1,352	1,368	1,408	1,355	1,527	1,531	1,515
Substance-related disorders	307	352	370	404	430	450	480	449	505	511	528
Suicide and intentional self-inflicted injury	66	77	80	86	91	91	106	106	113	114	113



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Pike

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,987	5,241	5,329	5,378	5,407	5,354	5,294	4,922	4,980	4,896	4,532
Female	2,454	2,547	2,574	2,590	2,592	2,530	2,468	2,314	2,316	2,261	2,071
Male	2,533	2,694	2,755	2,788	2,815	2,824	2,826	2,608	2,664	2,635	2,461
White	4,667	4,895	4,954	4,981	4,993	4,922	4,835	4,475	4,527	4,428	4,098
Minority	73	72	70	61	59	53	49	47	45	46	47

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	50%	52%	54%	54%	54%	54%	55%	52%	53%	52%	48%
Female	51%	53%	54%	53%	53%	53%	53%	50%	50%	49%	45%
Male	49%	52%	54%	54%	55%	55%	57%	53%	55%	54%	50%
White	49%	51%	53%	53%	53%	53%	53%	50%	51%	50%	46%
Minority	18%	17%	15%	12%	11%	10%	9%	8%	8%	7%	7%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	611	650	670	704	691	684	681	622	614	588	525
Alcohol-related disorders	196	184	178	161	147	132	117	85	66	53	42
Anxiety disorders	1,611	1,648	1,676	1,698	1,712	1,687	1,685	1,549	1,517	1,456	1,329
Attention-deficit, conduct, and disruptive behavior disorders	1,751	1,879	1,975	2,033	2,063	2,044	2,071	1,901	1,882	1,828	1,682
Delirium, dementia, and amnestic and other cognitive disorders	51	50	51	47	45	50	49	45	41	42	39
Developmental disorders	1,010	1,137	1,248	1,329	1,433	1,501	1,547	1,507	1,562	1,550	1,491
Disorders usually diagnosed in infancy, childhood, adolescence	279	304	322	342	366	379	401	387	379	370	350
Impulse control disorders, NEC	83	86	88	86	81	80	78	66	64	60	47
Miscellaneous disorders	1,011	1,032	1,028	1,040	1,032	1,014	990	911	894	859	783
Mood disorders	1,661	1,635	1,594	1,533	1,449	1,363	1,272	1,110	1,025	949	807
Personality disorders	61	64	70	65	57	60	53	42	41	39	31
Schizophrenia and other psychotic disorders	149	154	159	154	144	139	133	106	101	97	80
Screening and history of mental health and substance abuse	3,162	3,274	3,326	3,289	3,208	3,108	3,030	2,739	2,697	2,564	2,305
Substance-related disorders	607	557	522	479	435	397	356	335	313	295	263
Suicide and intentional self-inflicted injury	124	113	117	114	105	105	96	87	75	68	54



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Pike

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	6,535	6,898	7,253	7,467	7,681	7,767	7,872	7,390	7,894	7,972	7,732
Female	4,348	4,579	4,783	4,935	5,071	5,161	5,208	4,897	5,230	5,263	5,083
Male	2,187	2,319	2,470	2,532	2,610	2,606	2,664	2,493	2,664	2,709	2,649
White	6,010	6,358	6,692	6,904	7,109	7,186	7,288	6,805	7,295	7,390	7,149
Minority	142	135	132	116	95	76	46	36	41	40	41

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	67%	70%	71%	72%	72%	73%	68%	72%	72%	69%
Female	68%	70%	73%	74%	75%	75%	76%	71%	76%	75%	72%
Male	60%	62%	65%	66%	67%	65%	67%	63%	66%	66%	64%
White	64%	66%	69%	71%	71%	71%	72%	67%	71%	72%	69%
Minority	21%	19%	19%	17%	14%	11%	7%	5%	6%	5%	5%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	343	374	399	406	433	447	465	426	465	474	460
Alcohol-related disorders	718	783	826	856	872	878	921	903	946	928	857
Anxiety disorders	4,217	4,522	4,765	4,985	5,127	5,195	5,313	5,046	5,289	5,298	5,112
Attention-deficit, conduct, and disruptive behavior disorders	154	188	204	223	256	283	322	319	402	439	468
Delirium, dementia, and amnesic and other cognitive disorders	956	959	961	939	916	904	881	802	805	782	708
Developmental disorders	390	410	423	437	445	454	463	470	499	511	510
Disorders usually diagnosed in infancy, childhood, adolescence	35	42	42	43	46	45	46	51	64	71	73
Impulse control disorders, NEC	46	48	53	60	64	68	69	68	71	73	79
Miscellaneous disorders	2,113	2,284	2,465	2,667	2,787	2,819	2,850	2,652	2,720	2,697	2,565
Mood disorders	4,246	4,558	4,887	5,083	5,216	5,297	5,401	5,159	5,421	5,402	5,234
Personality disorders	254	286	293	307	314	312	321	307	306	302	303
Schizophrenia and other psychotic disorders	994	1,056	1,076	1,088	1,083	1,088	1,076	997	993	974	918
Screening and history of mental health and substance abuse	4,256	4,582	4,960	5,243	5,458	5,527	5,736	5,532	5,952	5,996	5,829
Substance-related disorders	1,490	1,641	1,752	1,869	1,998	2,082	2,228	2,180	2,367	2,453	2,405
Suicide and intentional self-inflicted injury	249	274	294	295	323	346	380	379	389	388	383



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Pulaski

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,869	4,281	4,472	4,714	4,790	4,824	4,957	4,696	4,829	4,737	4,363
Female	1,936	2,122	2,210	2,287	2,304	2,311	2,359	2,214	2,294	2,265	2,067
Male	1,933	2,159	2,262	2,427	2,486	2,513	2,598	2,482	2,535	2,472	2,296
White	3,586	3,965	4,124	4,336	4,366	4,379	4,471	4,220	4,320	4,238	3,894
Minority	98	103	106	103	112	112	115	121	134	133	127

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	58%	59%	59%	59%	59%	59%	56%	56%	53%	47%
Female	58%	59%	59%	58%	58%	58%	57%	53%	54%	50%	45%
Male	56%	58%	59%	60%	60%	61%	61%	58%	59%	55%	49%
White	56%	57%	58%	58%	58%	58%	58%	55%	54%	51%	46%
Minority	26%	24%	23%	20%	21%	19%	18%	18%	20%	18%	17%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	764	844	875	889	889	884	874	790	773	715	616
Alcohol-related disorders	210	202	193	187	169	151	143	111	98	76	56
Anxiety disorders	1,816	1,996	2,119	2,216	2,248	2,262	2,321	2,185	2,215	2,119	1,953
Attention-deficit, conduct, and disruptive behavior disorders	2,104	2,306	2,410	2,512	2,530	2,500	2,487	2,259	2,231	2,141	1,911
Delirium, dementia, and amnestic and other cognitive disorders	67	70	71	71	66	65	66	64	61	54	45
Developmental disorders	773	861	909	1,000	1,053	1,116	1,209	1,220	1,319	1,334	1,257
Disorders usually diagnosed in infancy, childhood, adolescence	337	389	417	433	450	450	457	408	419	416	382
Impulse control disorders, NEC	263	273	272	274	249	229	229	178	166	149	116
Miscellaneous disorders	497	522	521	510	503	476	482	413	397	351	305
Mood disorders	1,651	1,745	1,742	1,706	1,623	1,511	1,451	1,263	1,190	1,078	898
Personality disorders	121	123	111	105	95	94	86	69	62	62	40
Schizophrenia and other psychotic disorders	186	187	186	178	168	159	161	135	130	122	99
Screening and history of mental health and substance abuse	2,067	2,232	2,233	2,216	2,171	2,049	2,038	1,845	1,813	1,707	1,510
Substance-related disorders	539	533	520	495	454	415	386	314	281	244	187
Suicide and intentional self-inflicted injury	178	177	172	160	149	146	153	139	141	134	104



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Pulaski

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,132	5,434	5,664	5,783	5,983	6,109	6,349	6,026	6,427	6,485	6,291
Female	3,468	3,652	3,771	3,856	3,969	4,061	4,228	4,022	4,291	4,268	4,060
Male	1,664	1,782	1,893	1,927	2,014	2,048	2,121	2,004	2,136	2,217	2,231
White	4,719	4,997	5,229	5,338	5,541	5,660	5,879	5,592	5,991	6,048	5,825
Minority	163	170	156	157	139	134	129	109	116	114	127

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	69%	72%	70%	70%	70%	72%	68%	70%	69%	66%
Female	71%	72%	73%	72%	72%	72%	74%	69%	72%	71%	67%
Male	63%	65%	68%	67%	68%	67%	69%	65%	66%	65%	63%
White	68%	69%	71%	70%	70%	70%	72%	67%	70%	69%	66%
Minority	28%	29%	27%	26%	23%	21%	21%	18%	20%	18%	17%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	448	502	528	578	588	621	660	636	710	740	729
Alcohol-related disorders	447	506	530	547	594	583	599	569	593	614	590
Anxiety disorders	3,290	3,516	3,712	3,860	4,007	4,121	4,342	4,184	4,454	4,514	4,368
Attention-deficit, conduct, and disruptive behavior disorders	342	382	420	473	511	556	623	642	727	783	800
Delirium, dementia, and amnesic and other cognitive disorders	1,122	1,133	1,132	1,087	1,056	1,023	1,002	953	931	888	834
Developmental disorders	779	789	805	816	840	851	894	848	877	885	901
Disorders usually diagnosed in infancy, childhood, adolescence	186	193	206	210	214	219	230	219	238	260	263
Impulse control disorders, NEC	161	182	193	212	229	231	256	258	274	287	295
Miscellaneous disorders	1,041	1,131	1,225	1,285	1,334	1,387	1,470	1,419	1,512	1,501	1,465
Mood disorders	3,309	3,526	3,689	3,827	4,000	4,072	4,276	4,112	4,377	4,423	4,300
Personality disorders	423	439	455	461	480	474	491	455	487	466	472
Schizophrenia and other psychotic disorders	1,068	1,103	1,156	1,153	1,148	1,104	1,120	1,064	1,085	1,052	1,024
Screening and history of mental health and substance abuse	2,822	3,086	3,281	3,427	3,620	3,734	3,960	3,760	4,025	4,060	3,922
Substance-related disorders	799	878	935	981	1,064	1,092	1,199	1,141	1,255	1,301	1,299
Suicide and intentional self-inflicted injury	287	296	322	335	353	374	403	400	447	462	457



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Rowan

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,287	1,384	1,451	1,512	1,542	1,547	1,614	1,524	1,574	1,566	1,480
Female	670	712	718	738	749	732	765	716	722	701	651
Male	617	672	733	774	793	815	849	808	852	865	829
White	1,199	1,287	1,337	1,388	1,404	1,404	1,454	1,382	1,423	1,412	1,333
Minority	N.D.	N.D.	38	37	38	38	38	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	58%	60%	62%	61%	60%	61%	57%	57%	55%	51%
Female	60%	60%	60%	63%	61%	58%	59%	54%	55%	52%	48%
Male	55%	57%	60%	62%	62%	62%	62%	59%	59%	59%	54%
White	57%	58%	59%	61%	60%	59%	59%	55%	55%	54%	50%
Minority	N.D.	N.D.	26%	24%	22%	20%	20%	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	287	317	341	347	357	357	369	328	325	317	294
Alcohol-related disorders	54	62	54	54	50	44	39	33	N.D.	N.D.	N.D.
Anxiety disorders	495	520	529	568	599	591	624	598	596	588	548
Attention-deficit, conduct, and disruptive behavior disorders	534	580	613	642	661	675	721	671	674	665	620
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	330	379	426	462	477	504	539	535	543	563	557
Disorders usually diagnosed in infancy, childhood, adolescence	117	134	144	166	182	188	207	184	183	173	167
Impulse control disorders, NEC	59	62	67	70	70	72	76	64	63	63	57
Miscellaneous disorders	170	177	178	184	191	190	186	171	157	153	137
Mood disorders	551	580	569	581	580	553	555	477	458	422	366
Personality disorders	59	55	57	57	54	45	47	39	36	32	N.D.
Schizophrenia and other psychotic disorders	51	49	54	54	56	55	56	44	43	43	42
Screening and history of mental health and substance abuse	1,036	1,080	1,123	1,140	1,132	1,084	1,113	1,018	1,035	1,000	929
Substance-related disorders	152	148	142	142	132	116	113	106	89	77	54
Suicide and intentional self-inflicted injury	40	43	40	44	44	39	42	35	33	33	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Rowan

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,591	1,701	1,759	1,815	1,889	1,940	2,024	1,903	2,016	1,991	1,917
Female	1,068	1,115	1,176	1,223	1,257	1,293	1,355	1,272	1,342	1,311	1,257
Male	523	586	583	592	632	647	669	631	674	680	660
White	1,447	1,553	1,598	1,638	1,711	1,764	1,855	1,734	1,838	1,819	1,753
Minority	41	36	40	46	44	35	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	71%	73%	73%	74%	74%	73%	76%	71%	73%	70%	67%
Female	72%	75%	76%	77%	76%	76%	79%	73%	76%	73%	69%
Male	68%	70%	68%	70%	71%	69%	71%	67%	68%	66%	64%
White	70%	73%	73%	74%	74%	73%	75%	70%	72%	70%	67%
Minority	21%	19%	20%	22%	20%	16%	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	149	169	188	196	214	226	249	245	258	241	236
Alcohol-related disorders	208	228	243	242	264	263	279	264	272	255	244
Anxiety disorders	962	1,043	1,105	1,167	1,221	1,250	1,320	1,252	1,310	1,282	1,240
Attention-deficit, conduct, and disruptive behavior disorders	74	74	88	95	113	121	138	136	156	170	178
Delirium, dementia, and amnesic and other cognitive disorders	414	417	416	414	392	371	358	330	307	288	246
Developmental disorders	277	292	302	326	348	354	358	331	326	337	335
Disorders usually diagnosed in infancy, childhood, adolescence	40	43	47	47	50	56	63	58	62	67	66
Impulse control disorders, NEC	63	71	72	73	78	88	88	84	90	99	92
Miscellaneous disorders	355	385	410	423	444	465	498	462	484	473	445
Mood disorders	984	1,063	1,110	1,166	1,253	1,296	1,369	1,297	1,356	1,333	1,280
Personality disorders	237	265	282	290	307	310	323	299	309	291	296
Schizophrenia and other psychotic disorders	345	355	361	364	363	361	392	363	362	355	335
Screening and history of mental health and substance abuse	934	1,053	1,124	1,180	1,249	1,307	1,387	1,333	1,457	1,456	1,417
Substance-related disorders	279	316	336	354	388	412	438	435	459	459	452
Suicide and intentional self-inflicted injury	52	56	58	59	69	78	90	90	95	93	104



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Warren

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,465	5,005	5,297	5,467	5,562	5,621	5,692	5,414	5,775	5,947	5,645
Female	2,144	2,386	2,530	2,600	2,640	2,632	2,650	2,428	2,613	2,720	2,564
Male	2,321	2,619	2,767	2,867	2,922	2,989	3,042	2,986	3,162	3,227	3,081
White	3,287	3,694	3,870	3,976	4,000	4,041	4,080	3,887	4,129	4,282	4,087
Minority	973	1,071	1,145	1,184	1,223	1,234	1,219	1,144	1,235	1,245	1,170

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	54%	55%	55%	54%	53%	53%	49%	50%	48%	42%
Female	51%	53%	54%	54%	52%	51%	50%	46%	47%	45%	40%
Male	52%	54%	56%	56%	56%	55%	55%	52%	53%	50%	45%
White	52%	54%	55%	55%	55%	53%	52%	49%	50%	49%	44%
Minority	42%	43%	43%	43%	42%	41%	41%	38%	38%	33%	29%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	774	849	929	955	952	936	894	814	809	797	728
Alcohol-related disorders	158	160	159	156	146	143	141	107	98	83	61
Anxiety disorders	1,173	1,343	1,458	1,506	1,615	1,680	1,731	1,689	1,778	1,803	1,699
Attention-deficit, conduct, and disruptive behavior disorders	2,193	2,440	2,636	2,758	2,767	2,791	2,826	2,632	2,659	2,621	2,421
Delirium, dementia, and amnestic and other cognitive disorders	64	63	65	66	64	62	63	57	59	60	57
Developmental disorders	1,097	1,308	1,451	1,578	1,711	1,769	1,867	1,885	1,998	2,063	2,022
Disorders usually diagnosed in infancy, childhood, adolescence	393	457	518	566	591	627	655	643	677	683	656
Impulse control disorders, NEC	398	437	451	460	452	432	392	352	332	314	267
Miscellaneous disorders	642	694	728	741	729	731	698	635	633	605	557
Mood disorders	1,613	1,700	1,739	1,703	1,642	1,537	1,467	1,307	1,249	1,163	1,034
Personality disorders	93	101	95	95	91	85	68	53	56	51	39
Schizophrenia and other psychotic disorders	185	199	202	193	201	189	176	164	155	147	123
Screening and history of mental health and substance abuse	3,315	3,553	3,637	3,547	3,412	3,252	3,142	2,813	2,949	2,918	2,708
Substance-related disorders	394	415	409	406	356	335	322	264	246	222	168
Suicide and intentional self-inflicted injury	99	100	100	103	94	84	81	64	62	57	46




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

BRFSS Group: Warren

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,621	5,191	5,555	5,626	5,685	5,754	5,951	5,419	5,851	6,079	5,941
Female	3,281	3,660	3,924	3,997	4,050	4,112	4,213	3,803	4,148	4,252	4,106
Male	1,340	1,531	1,631	1,629	1,635	1,642	1,738	1,616	1,703	1,827	1,835
White	3,656	4,107	4,391	4,443	4,475	4,537	4,686	4,261	4,592	4,773	4,648
Minority	757	845	905	918	926	918	938	834	904	937	917
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	69%	70%	70%	69%	70%	64%	68%	65%	60%
Female	65%	68%	70%	71%	71%	69%	70%	63%	68%	66%	61%
Male	62%	66%	68%	69%	68%	68%	69%	65%	67%	64%	59%
White	65%	69%	71%	72%	72%	71%	72%	66%	70%	69%	63%
Minority	47%	48%	49%	50%	49%	47%	47%	42%	43%	40%	37%
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	438	476	512	524	537	545	565	522	569	602	592
Alcohol-related disorders	532	604	650	651	678	675	692	629	663	680	657
Anxiety disorders	2,431	2,693	2,944	3,067	3,154	3,231	3,365	3,195	3,412	3,519	3,426
Attention-deficit, conduct, and disruptive behavior disorders	361	424	460	522	599	636	710	705	831	899	910
Delirium, dementia, and amnesic and other cognitive disorders	1,082	1,125	1,145	1,103	1,054	1,042	1,044	964	943	895	831
Developmental disorders	480	515	532	542	547	545	611	592	611	651	636
Disorders usually diagnosed in infancy, childhood, adolescence	60	69	77	82	89	95	119	114	134	152	155
Impulse control disorders, NEC	119	124	139	142	161	160	186	184	210	220	229
Miscellaneous disorders	979	1,129	1,209	1,299	1,327	1,404	1,504	1,381	1,436	1,511	1,417
Mood disorders	2,973	3,346	3,636	3,748	3,865	3,912	4,060	3,762	4,032	4,158	4,044
Personality disorders	328	347	364	366	366	354	373	356	373	361	358
Schizophrenia and other psychotic disorders	1,089	1,166	1,224	1,237	1,244	1,246	1,283	1,233	1,265	1,215	1,142
Screening and history of mental health and substance abuse	2,940	3,373	3,717	3,804	3,880	3,877	4,019	3,655	3,978	4,112	4,010
Substance-related disorders	692	780	832	858	921	953	1,026	962	1,038	1,053	1,026
Suicide and intentional self-inflicted injury	184	201	228	248	269	273	316	310	344	346	347



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Adair County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,120	1,203	1,231	1,264	1,294	1,336	1,320	1,242	1,293	1,288	1,169
Female	582	626	615	617	624	632	625	580	606	599	526
Male	538	577	616	647	670	704	695	662	687	689	643
White	1,021	1,098	1,129	1,148	1,177	1,210	1,193	1,126	1,176	1,165	1,053
Minority	56	56	53	60	52	54	43	40	38	38	36

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	56%	59%	59%	59%	59%	58%	54%	54%	50%	44%
Female	55%	56%	59%	59%	58%	58%	56%	51%	51%	46%	39%
Male	52%	55%	58%	59%	59%	60%	60%	58%	58%	53%	49%
White	53%	55%	58%	58%	58%	58%	57%	53%	53%	49%	43%
Minority	36%	36%	34%	36%	30%	31%	26%	25%	24%	19%	18%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	265	272	275	289	292	296	289	270	262	256	232
Alcohol-related disorders	49	49	49	39	33	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	587	618	611	627	661	681	679	633	621	602	542
Attention-deficit, conduct, and disruptive behavior disorders	533	551	590	615	629	634	628	587	579	575	530
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	194	246	272	294	309	343	358	348	377	379	363
Disorders usually diagnosed in infancy, childhood, adolescence	87	90	97	105	107	113	116	112	112	106	99
Impulse control disorders, NEC	86	81	85	88	86	86	80	63	55	46	41
Miscellaneous disorders	205	220	204	199	192	189	179	167	166	160	137
Mood disorders	451	444	420	408	399	389	367	313	302	280	237
Personality disorders	45	40	41	38	34	31	32	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	56	54	51	47	52	49	47	41	36	31	N.D.
Screening and history of mental health and substance abuse	682	695	707	702	670	655	627	558	559	528	477
Substance-related disorders	146	135	121	111	105	103	95	73	55	53	43
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Adair County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,492	1,582	1,645	1,657	1,680	1,672	1,732	1,637	1,768	1,802	1,645
Female	1,023	1,087	1,117	1,120	1,138	1,148	1,195	1,111	1,220	1,223	1,117
Male	469	495	528	537	542	524	537	526	548	579	528
White	1,342	1,424	1,479	1,504	1,524	1,521	1,582	1,497	1,629	1,664	1,512
Minority	79	83	80	68	71	60	51	47	56	51	44

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	69%	72%	73%	74%	74%	73%	75%	70%	72%	72%	65%
Female	70%	75%	75%	77%	76%	75%	77%	72%	75%	74%	69%
Male	65%	67%	68%	69%	70%	68%	69%	66%	66%	67%	58%
White	68%	72%	73%	74%	74%	73%	75%	70%	72%	72%	65%
Minority	37%	40%	37%	32%	34%	30%	25%	24%	28%	26%	22%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	172	189	213	228	233	251	264	260	295	293	284
Alcohol-related disorders	124	135	145	160	166	164	165	153	158	167	150
Anxiety disorders	929	1,002	1,069	1,099	1,124	1,172	1,236	1,172	1,240	1,283	1,207
Attention-deficit, conduct, and disruptive behavior disorders	56	58	79	84	90	94	122	133	155	166	171
Delirium, dementia, and amnesic and other cognitive disorders	352	353	355	334	326	305	311	287	281	263	229
Developmental disorders	172	177	185	186	191	190	198	204	200	204	188
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	36	36	35	37	38	43	53	45	52	62	58
Miscellaneous disorders	330	367	396	413	439	448	464	426	460	464	435
Mood disorders	956	1,027	1,076	1,081	1,109	1,134	1,184	1,130	1,199	1,207	1,142
Personality disorders	179	186	196	205	210	214	213	204	208	212	204
Schizophrenia and other psychotic disorders	414	429	434	415	409	397	398	366	368	359	328
Screening and history of mental health and substance abuse	747	815	858	897	908	912	969	933	1,027	1,042	951
Substance-related disorders	256	281	309	323	332	329	353	339	386	387	348
Suicide and intentional self-inflicted injury	62	64	68	77	75	76	73	76	78	90	83



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Allen County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	815	902	961	1,007	1,013	1,048	1,118	1,117	1,141	1,181	1,124
Female	408	440	479	508	516	538	556	548	561	572	532
Male	407	462	482	499	497	510	562	569	580	609	592
White	761	836	897	938	937	962	1,028	1,022	1,039	1,084	1,022
Minority	N.D.	34	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	53%	53%	52%	51%	51%	51%	50%	49%	46%	42%
Female	55%	54%	55%	54%	53%	51%	50%	49%	48%	44%	40%
Male	50%	52%	51%	50%	50%	50%	51%	51%	50%	47%	44%
White	52%	52%	52%	51%	50%	50%	50%	49%	48%	45%	41%
Minority	N.D.	32%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	165	174	189	201	199	209	219	212	208	205	190
Alcohol-related disorders	31	33	31	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	280	308	363	390	397	423	448	446	448	431	414
Attention-deficit, conduct, and disruptive behavior disorders	369	409	448	479	491	515	546	526	518	532	498
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	97	119	131	156	183	187	227	246	283	305	292
Disorders usually diagnosed in infancy, childhood, adolescence	61	62	74	77	83	85	96	94	93	94	89
Impulse control disorders, NEC	43	47	53	52	52	50	55	47	44	37	33
Miscellaneous disorders	110	121	131	129	124	129	131	123	117	116	108
Mood disorders	311	322	349	361	347	344	342	319	287	264	237
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	33	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	549	601	606	594	552	537	543	512	503	501	462
Substance-related disorders	76	77	73	74	62	58	55	47	38	33	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Allen County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	997	1,108	1,166	1,192	1,215	1,226	1,292	1,255	1,343	1,375	1,326
Female	684	762	801	803	817	820	874	844	893	912	896
Male	313	346	365	389	398	406	418	411	450	463	430
White	885	988	1,051	1,071	1,094	1,104	1,167	1,147	1,233	1,260	1,210
Minority	53	58	49	52	48	40	37	N.D.	32	38	35

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	71%	69%	69%	68%	69%	71%	67%	70%	67%	63%
Female	69%	72%	71%	71%	70%	70%	72%	68%	72%	69%	65%
Male	66%	68%	65%	67%	65%	66%	68%	65%	66%	64%	60%
White	68%	70%	69%	69%	68%	68%	70%	66%	69%	67%	63%
Minority	33%	35%	30%	33%	30%	26%	24%	N.D.	22%	23%	21%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	80	95	105	106	112	115	125	123	133	122	129
Alcohol-related disorders	87	105	110	112	119	112	127	127	130	135	123
Anxiety disorders	582	654	693	717	746	762	823	833	894	895	870
Attention-deficit, conduct, and disruptive behavior disorders	47	53	64	70	88	81	107	122	127	134	131
Delirium, dementia, and amnesic and other cognitive disorders	320	331	328	306	288	272	257	241	244	229	202
Developmental disorders	101	102	102	109	111	114	124	124	135	139	128
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	34	36	41	34
Miscellaneous disorders	158	188	216	225	249	259	289	281	299	289	281
Mood disorders	603	675	736	765	806	793	857	860	915	932	891
Personality disorders	49	53	59	64	66	67	75	73	74	80	76
Schizophrenia and other psychotic disorders	244	253	258	266	262	264	273	271	273	262	235
Screening and history of mental health and substance abuse	491	572	611	638	674	680	733	713	751	762	739
Substance-related disorders	99	121	140	145	159	149	170	180	189	202	181
Suicide and intentional self-inflicted injury	N.D.	N.D.	40	39	46	47	61	70	67	71	70



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Anderson County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	633	715	787	838	893	901	935	891	974	1,028	1,011
Female	305	334	371	392	415	421	447	419	459	490	474
Male	328	381	416	446	478	480	488	472	515	538	537
White	568	645	708	755	800	803	828	791	862	919	901
Minority	37	38	40	38	39	41	45	47	56	51	51

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	56%	60%	60%	61%	61%	60%	55%	53%	50%	48%
Female	54%	54%	58%	58%	59%	57%	58%	53%	51%	48%	46%
Male	56%	59%	63%	63%	62%	64%	63%	57%	56%	52%	50%
White	54%	55%	60%	60%	60%	60%	59%	55%	52%	50%	46%
Minority	42%	37%	34%	30%	30%	29%	29%	27%	31%	28%	29%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	217	246	258	280	291	289	308	278	280	278	271
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	223	237	252	277	283	280	277	264	274	292	269
Attention-deficit, conduct, and disruptive behavior disorders	340	386	419	451	482	482	499	459	472	486	463
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	120	157	191	221	259	277	294	299	339	386	388
Disorders usually diagnosed in infancy, childhood, adolescence	78	94	101	106	133	138	140	120	133	135	129
Impulse control disorders, NEC	40	45	40	36	34	34	35	33	31	N.D.	N.D.
Miscellaneous disorders	143	155	166	178	177	179	183	165	177	182	171
Mood disorders	242	256	282	282	289	280	271	255	246	247	227
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	33	34	34	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	367	398	437	442	459	436	436	397	422	409	404
Substance-related disorders	69	69	72	69	66	61	58	57	55	57	45
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	34	35	34	34	31



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Anderson County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	509	549	552	623	658	661	684	692	765	816	803
Female	374	402	398	455	475	482	493	496	547	569	559
Male	135	147	154	168	183	179	191	196	218	247	244
White	447	479	484	555	584	587	609	619	692	739	723
Minority	39	43	40	39	43	43	39	36	36	40	43

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	65%	64%	67%	68%	68%	67%	66%	71%	69%	64%
Female	64%	68%	66%	70%	69%	69%	68%	66%	71%	69%	66%
Male	58%	58%	60%	60%	65%	64%	65%	65%	69%	67%	61%
White	62%	64%	64%	67%	68%	68%	68%	66%	71%	69%	65%
Minority	43%	42%	39%	36%	37%	38%	33%	31%	31%	33%	34%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	55	62	65	81	82	86	91	97	99	110	109
Alcohol-related disorders	39	45	59	63	65	61	70	71	79	71	62
Anxiety disorders	243	288	307	351	367	365	398	416	446	467	487
Attention-deficit, conduct, and disruptive behavior disorders	34	44	55	57	61	72	74	81	93	111	114
Delirium, dementia, and amnestic and other cognitive disorders	153	150	150	148	147	135	135	128	123	125	130
Developmental disorders	43	48	44	47	48	55	62	63	64	66	67
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	113	142	161	190	215	228	247	242	271	279	267
Mood disorders	289	319	342	382	405	420	438	436	487	510	500
Personality disorders	50	58	66	78	78	81	84	85	85	87	83
Schizophrenia and other psychotic disorders	121	125	133	145	155	149	159	153	150	150	138
Screening and history of mental health and substance abuse	274	305	329	387	405	410	439	441	492	509	496
Substance-related disorders	78	82	98	117	121	129	145	151	178	191	192
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	36	36	40	39	40



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Ballard County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	365	424	454	488	469	471	462	419	427	416	407
Female	171	187	196	207	197	194	193	176	182	172	165
Male	194	237	258	281	272	277	269	243	245	244	242
White	310	355	377	411	378	378	362	324	329	323	318
Minority	41	45	47	46	51	51	50	45	45	41	37

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	51%	55%	54%	56%	56%	58%	57%	51%	52%	48%	45%
Female	52%	54%	53%	53%	50%	52%	52%	48%	51%	44%	39%
Male	50%	56%	56%	59%	60%	62%	61%	53%	53%	52%	50%
White	49%	53%	52%	55%	52%	55%	53%	47%	48%	45%	42%
Minority	48%	48%	43%	40%	41%	40%	38%	35%	34%	29%	25%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	43	56	56	53	50	44	46	44	46	55	53
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	74	88	94	102	103	96	96	86	86	73	66
Attention-deficit, conduct, and disruptive behavior disorders	202	238	249	272	277	274	274	249	240	243	248
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	105	124	142	160	166	178	179	168	175	165	160
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	34	36	43	44	47	46	41	36	37	33
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	38	45	44	51	49	43	41	38	36	33	31
Mood disorders	127	142	147	154	145	142	131	122	121	100	90
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	176	206	214	208	201	195	181	159	160	152	150
Substance-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Ballard County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	407	468	506	514	516	525	519	482	514	528	511
Female	312	351	370	376	379	387	376	344	363	372	367
Male	95	117	136	138	137	138	143	138	151	156	144
White	356	421	453	458	453	464	451	426	458	470	454
Minority	38	33	37	37	41	36	36	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	61%	65%	69%	69%	67%	70%	68%	66%	69%	69%	63%
Female	66%	70%	74%	72%	71%	73%	72%	69%	72%	71%	64%
Male	50%	53%	60%	61%	58%	62%	61%	59%	62%	64%	60%
White	61%	64%	69%	69%	66%	69%	67%	65%	69%	69%	63%
Minority	49%	47%	49%	46%	48%	46%	43%	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	36	38	42	49	47	47	51	45	40	38	40
Alcohol-related disorders	N.D.	N.D.	32	31	32	33	34	N.D.	37	35	40
Anxiety disorders	214	239	256	279	282	289	301	280	294	307	306
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	40	44	42	47	47
Delirium, dementia, and amnesic and other cognitive disorders	125	132	132	123	113	103	103	96	98	88	80
Developmental disorders	34	38	39	36	33	38	41	39	42	45	45
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	75	89	102	111	116	116	120	106	108	115	110
Mood disorders	248	290	318	334	333	345	350	324	332	344	343
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	84	87	99	98	93	88	91	86	88	86	83
Screening and history of mental health and substance abuse	193	235	265	279	282	298	298	268	286	301	295
Substance-related disorders	36	44	47	58	59	68	77	68	72	75	72
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	32



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Barren County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,956	2,235	2,388	2,473	2,546	2,661	2,737	2,585	2,649	2,734	2,549
Female	979	1,117	1,174	1,205	1,246	1,282	1,310	1,243	1,272	1,265	1,172
Male	977	1,118	1,214	1,268	1,300	1,379	1,427	1,342	1,377	1,469	1,377
White	1,719	1,975	2,114	2,174	2,239	2,348	2,395	2,278	2,321	2,396	2,237
Minority	157	170	171	183	173	168	171	150	158	153	147

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	58%	60%	59%	58%	57%	58%	54%	54%	51%	46%
Female	56%	58%	59%	58%	57%	55%	55%	52%	52%	49%	43%
Male	56%	57%	60%	60%	59%	59%	60%	56%	56%	54%	49%
White	56%	57%	59%	58%	57%	57%	57%	53%	53%	50%	45%
Minority	39%	40%	40%	41%	36%	33%	33%	28%	29%	26%	24%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	396	464	468	484	491	508	511	480	501	498	438
Alcohol-related disorders	75	76	75	69	65	58	52	42	44	38	N.D.
Anxiety disorders	736	787	834	851	905	936	964	913	922	920	851
Attention-deficit, conduct, and disruptive behavior disorders	974	1,108	1,166	1,216	1,265	1,334	1,369	1,255	1,278	1,282	1,170
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	33	34	33	39	37	40	38	38	36	33
Developmental disorders	363	439	508	563	618	682	755	740	782	826	807
Disorders usually diagnosed in infancy, childhood, adolescence	153	174	187	211	236	251	271	252	274	285	276
Impulse control disorders, NEC	147	156	163	159	152	141	142	121	123	118	92
Miscellaneous disorders	375	398	399	410	409	406	402	360	347	324	292
Mood disorders	748	783	787	759	740	714	680	600	587	534	439
Personality disorders	49	54	54	52	54	46	46	41	36	N.D.	N.D.
Schizophrenia and other psychotic disorders	84	87	90	92	94	88	85	64	64	50	45
Screening and history of mental health and substance abuse	1,335	1,513	1,548	1,503	1,449	1,422	1,393	1,237	1,256	1,228	1,103
Substance-related disorders	198	204	203	187	178	165	150	127	131	109	79
Suicide and intentional self-inflicted injury	69	65	69	56	61	53	58	52	47	43	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Barren County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,588	2,779	2,973	3,033	3,115	3,193	3,291	3,028	3,186	3,278	3,180
Female	1,793	1,947	2,091	2,139	2,184	2,243	2,294	2,077	2,192	2,255	2,194
Male	795	832	882	894	931	950	997	951	994	1,023	986
White	2,262	2,444	2,620	2,682	2,765	2,832	2,908	2,680	2,827	2,889	2,808
Minority	215	217	223	214	213	217	225	191	205	230	209

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	68%	69%	70%	70%	69%	70%	66%	69%	66%	63%
Female	66%	70%	71%	71%	71%	71%	72%	67%	71%	69%	65%
Male	63%	63%	64%	66%	66%	65%	66%	64%	65%	61%	57%
White	65%	68%	69%	70%	70%	69%	70%	66%	69%	67%	63%
Minority	43%	43%	44%	42%	41%	41%	41%	36%	39%	37%	33%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	162	179	192	219	229	252	283	258	274	290	296
Alcohol-related disorders	208	229	242	240	257	258	273	258	268	276	268
Anxiety disorders	1,523	1,631	1,781	1,845	1,935	2,002	2,094	1,957	2,057	2,108	2,061
Attention-deficit, conduct, and disruptive behavior disorders	115	135	154	169	195	221	249	238	264	313	320
Delirium, dementia, and amnesic and other cognitive disorders	769	804	818	781	735	719	684	610	589	562	505
Developmental disorders	251	266	286	288	290	295	315	301	305	321	310
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	34	42	43
Impulse control disorders, NEC	N.D.	35	37	49	54	59	63	55	61	71	74
Miscellaneous disorders	574	630	702	745	768	801	833	785	809	847	823
Mood disorders	1,638	1,782	1,938	2,016	2,087	2,148	2,224	2,075	2,184	2,213	2,139
Personality disorders	114	135	158	163	169	178	177	167	173	180	176
Schizophrenia and other psychotic disorders	684	732	772	776	773	770	784	735	725	723	671
Screening and history of mental health and substance abuse	1,252	1,374	1,532	1,583	1,668	1,711	1,811	1,685	1,765	1,826	1,767
Substance-related disorders	284	323	362	381	410	420	452	429	448	470	460
Suicide and intentional self-inflicted injury	76	85	102	112	116	125	132	121	143	148	160



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Bath County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,013	1,094	1,129	1,126	1,104	1,117	1,162	1,110	1,166	1,161	1,087
Female	523	570	558	552	534	546	574	535	559	545	507
Male	490	524	571	574	570	571	588	575	607	616	580
White	931	1,004	1,036	1,022	998	1,011	1,047	1,007	1,057	1,051	988
Minority	N.D.	N.D.	N.D.	N.D.	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	65%	68%	67%	65%	65%	64%	59%	60%	57%	52%
Female	67%	67%	67%	67%	64%	64%	64%	59%	61%	55%	50%
Male	61%	62%	68%	67%	66%	65%	64%	59%	60%	58%	53%
White	63%	64%	67%	65%	64%	63%	62%	58%	58%	55%	50%
Minority	N.D.	N.D.	N.D.	N.D.	25%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	191	199	208	212	190	204	214	204	200	209	189
Alcohol-related disorders	50	44	41	40	35	34	36	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	406	438	448	473	461	474	504	474	485	479	450
Attention-deficit, conduct, and disruptive behavior disorders	366	393	407	434	422	421	443	410	421	413	376
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	224	249	270	298	309	317	342	329	342	356	347
Disorders usually diagnosed in infancy, childhood, adolescence	85	91	94	113	110	116	122	111	120	118	113
Impulse control disorders, NEC	N.D.	N.D.	N.D.	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	118	118	119	123	102	99	103	100	98	100	91
Mood disorders	389	398	390	394	354	335	339	306	301	279	234
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	50	47	54	47	42	39	42	39	36	35	N.D.
Screening and history of mental health and substance abuse	793	847	860	831	786	774	791	733	746	715	651
Substance-related disorders	117	109	108	100	87	81	84	69	58	49	42
Suicide and intentional self-inflicted injury	39	36	34	34	33	33	33	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Bath County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,160	1,218	1,276	1,319	1,339	1,415	1,470	1,375	1,471	1,490	1,445
Female	769	800	840	860	885	929	976	886	956	944	923
Male	391	418	436	459	454	486	494	489	515	546	522
White	1,022	1,081	1,143	1,187	1,212	1,279	1,329	1,248	1,348	1,373	1,317
Minority	53	55	51	44	38	41	34	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	69%	70%	72%	72%	73%	75%	70%	74%	73%	68%
Female	71%	72%	74%	75%	76%	77%	79%	73%	77%	76%	73%
Male	62%	63%	64%	67%	65%	66%	68%	67%	69%	67%	60%
White	66%	68%	69%	71%	72%	72%	74%	70%	74%	72%	67%
Minority	31%	33%	30%	27%	23%	24%	20%	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	85	86	102	103	112	120	129	127	137	131	130
Alcohol-related disorders	115	123	135	149	164	169	181	171	180	184	170
Anxiety disorders	610	643	705	732	787	821	872	835	901	901	885
Attention-deficit, conduct, and disruptive behavior disorders	41	42	48	55	62	72	83	82	103	111	117
Delirium, dementia, and amnesic and other cognitive disorders	273	281	272	263	255	247	235	208	197	183	154
Developmental disorders	87	91	100	102	111	119	120	110	124	125	118
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	224	246	274	285	306	315	342	316	337	329	326
Mood disorders	638	687	735	786	846	884	931	878	952	956	934
Personality disorders	112	116	135	139	146	151	148	141	137	142	142
Schizophrenia and other psychotic disorders	255	271	286	288	273	266	268	247	254	258	231
Screening and history of mental health and substance abuse	709	746	820	882	899	971	1,042	989	1,082	1,080	1,053
Substance-related disorders	175	192	215	234	255	280	306	297	319	326	326
Suicide and intentional self-inflicted injury	32	32	38	39	44	50	61	60	65	71	76



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Bell County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,128	3,293	3,407	3,498	3,513	3,511	3,511	3,351	3,383	3,354	3,122
Female	1,594	1,642	1,683	1,707	1,701	1,707	1,698	1,635	1,626	1,589	1,477
Male	1,534	1,651	1,724	1,791	1,812	1,804	1,813	1,716	1,757	1,765	1,645
White	2,877	3,025	3,111	3,174	3,180	3,169	3,156	2,994	3,014	2,982	2,766
Minority	107	103	114	116	106	102	97	99	95	100	93

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	59%	61%	63%	63%	63%	64%	62%	63%	61%	57%
Female	60%	61%	62%	64%	63%	63%	64%	61%	61%	59%	55%
Male	55%	57%	60%	61%	62%	63%	65%	62%	64%	62%	59%
White	56%	58%	60%	61%	61%	62%	63%	60%	61%	59%	56%
Minority	31%	28%	30%	28%	25%	24%	22%	21%	21%	20%	18%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	544	571	609	623	614	617	599	576	574	547	486
Alcohol-related disorders	219	215	219	216	209	193	170	141	129	106	93
Anxiety disorders	1,677	1,765	1,837	1,918	1,945	1,967	1,958	1,896	1,892	1,877	1,729
Attention-deficit, conduct, and disruptive behavior disorders	1,120	1,202	1,260	1,318	1,330	1,327	1,311	1,213	1,200	1,165	1,088
Delirium, dementia, and amnestic and other cognitive disorders	32	N.D.	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	929	995	1,084	1,180	1,268	1,278	1,291	1,252	1,264	1,230	1,164
Disorders usually diagnosed in infancy, childhood, adolescence	175	194	206	231	246	256	254	251	249	241	230
Impulse control disorders, NEC	83	83	84	84	78	77	68	58	55	50	46
Miscellaneous disorders	430	426	414	398	378	350	330	313	315	294	271
Mood disorders	879	864	844	810	751	696	633	544	517	455	389
Personality disorders	189	189	190	194	182	167	147	127	123	104	85
Schizophrenia and other psychotic disorders	133	130	130	122	113	108	99	87	82	79	64
Screening and history of mental health and substance abuse	1,472	1,522	1,537	1,503	1,402	1,339	1,279	1,180	1,160	1,115	1,026
Substance-related disorders	602	592	588	567	524	472	416	352	324	275	225
Suicide and intentional self-inflicted injury	96	91	89	82	79	74	70	54	55	46	35



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Bell County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,221	4,502	4,650	4,775	4,869	4,842	4,896	4,625	4,741	4,770	4,662
Female	2,781	2,979	3,064	3,129	3,195	3,172	3,204	3,035	3,105	3,129	3,044
Male	1,440	1,523	1,586	1,646	1,674	1,670	1,692	1,590	1,636	1,641	1,618
White	3,823	4,085	4,226	4,351	4,443	4,432	4,483	4,244	4,356	4,377	4,273
Minority	177	181	171	163	154	129	111	98	103	106	96

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	69%	71%	71%	72%	71%	70%	71%	68%	70%	70%	67%
Female	72%	74%	75%	75%	74%	74%	75%	71%	74%	74%	70%
Male	64%	65%	65%	65%	65%	64%	65%	62%	64%	63%	61%
White	69%	70%	71%	71%	70%	70%	70%	67%	70%	69%	66%
Minority	32%	32%	31%	29%	28%	24%	22%	19%	20%	19%	17%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	352	372	394	402	418	434	443	416	425	443	434
Alcohol-related disorders	410	450	454	487	517	501	503	455	481	469	438
Anxiety disorders	3,103	3,308	3,445	3,517	3,588	3,591	3,682	3,516	3,579	3,609	3,549
Attention-deficit, conduct, and disruptive behavior disorders	133	147	163	189	226	235	277	258	282	315	331
Delirium, dementia, and amnesic and other cognitive disorders	749	753	759	737	705	685	671	628	605	569	501
Developmental disorders	305	329	344	372	405	407	422	407	430	452	456
Disorders usually diagnosed in infancy, childhood, adolescence	49	56	56	50	51	54	60	48	50	59	52
Impulse control disorders, NEC	52	58	65	64	78	77	86	77	86	87	86
Miscellaneous disorders	1,190	1,290	1,345	1,416	1,503	1,523	1,546	1,466	1,474	1,482	1,434
Mood disorders	2,271	2,441	2,598	2,716	2,792	2,781	2,822	2,690	2,757	2,792	2,745
Personality disorders	396	428	464	482	517	513	512	476	481	488	476
Schizophrenia and other psychotic disorders	696	736	747	757	768	741	743	691	673	654	640
Screening and history of mental health and substance abuse	2,130	2,352	2,501	2,612	2,700	2,721	2,777	2,610	2,682	2,699	2,650
Substance-related disorders	838	933	1,009	1,089	1,181	1,186	1,209	1,124	1,175	1,202	1,159
Suicide and intentional self-inflicted injury	159	167	179	189	202	210	232	235	236	249	263



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Boone County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,875	2,151	2,414	2,656	2,853	3,105	3,387	3,268	3,723	3,924	3,805
Female	916	1,001	1,112	1,222	1,302	1,416	1,545	1,478	1,670	1,700	1,672
Male	959	1,150	1,302	1,434	1,551	1,689	1,842	1,790	2,053	2,224	2,133
White	1,642	1,882	2,105	2,313	2,486	2,692	2,918	2,821	3,216	3,389	3,300
Minority	135	149	166	184	178	193	207	213	228	246	234

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	50%	52%	54%	54%	53%	53%	53%	49%	51%	46%	41%
Female	51%	51%	52%	52%	51%	49%	51%	46%	47%	42%	38%
Male	50%	53%	56%	56%	55%	56%	56%	52%	54%	49%	43%
White	51%	52%	55%	55%	53%	53%	53%	49%	50%	46%	41%
Minority	28%	26%	27%	27%	24%	25%	26%	25%	25%	21%	20%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	379	435	462	505	512	544	581	539	600	608	548
Alcohol-related disorders	90	98	99	95	98	93	88	78	81	69	61
Anxiety disorders	566	619	701	742	736	777	791	717	775	760	683
Attention-deficit, conduct, and disruptive behavior disorders	1,046	1,211	1,393	1,576	1,746	1,911	2,121	2,010	2,196	2,236	2,137
Delirium, dementia, and amnestic and other cognitive disorders	43	50	58	57	60	65	67	59	76	66	54
Developmental disorders	542	648	741	836	938	1,028	1,179	1,176	1,373	1,479	1,471
Disorders usually diagnosed in infancy, childhood, adolescence	252	292	356	404	423	461	506	486	521	537	507
Impulse control disorders, NEC	80	92	98	103	98	96	107	85	88	84	71
Miscellaneous disorders	308	330	364	389	388	396	427	378	416	422	391
Mood disorders	727	795	860	891	885	903	924	800	846	802	710
Personality disorders	86	94	102	100	84	83	83	69	69	56	46
Schizophrenia and other psychotic disorders	87	95	111	111	108	107	116	101	110	102	90
Screening and history of mental health and substance abuse	1,069	1,184	1,298	1,326	1,324	1,343	1,376	1,250	1,363	1,350	1,215
Substance-related disorders	188	205	217	211	210	199	184	172	183	169	145
Suicide and intentional self-inflicted injury	85	97	97	96	100	101	109	95	113	108	96



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Boone County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,755	1,992	2,231	2,387	2,577	2,754	3,070	2,788	3,146	3,341	3,202
Female	1,344	1,527	1,711	1,827	1,982	2,093	2,315	2,088	2,364	2,500	2,362
Male	411	465	520	560	595	661	755	700	782	841	840
White	1,597	1,807	2,049	2,191	2,367	2,536	2,833	2,575	2,915	3,072	2,939
Minority	85	111	99	110	114	112	112	97	108	134	114

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	64%	66%	67%	68%	68%	72%	63%	67%	65%	60%
Female	63%	66%	68%	69%	69%	70%	73%	64%	68%	66%	61%
Male	59%	58%	61%	63%	63%	63%	69%	63%	65%	63%	59%
White	63%	65%	68%	69%	69%	69%	74%	65%	68%	67%	62%
Minority	30%	33%	27%	29%	28%	28%	27%	22%	25%	26%	22%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	262	300	326	339	374	397	454	430	465	470	447
Alcohol-related disorders	218	253	277	299	325	347	400	359	381	392	364
Anxiety disorders	1,004	1,128	1,273	1,370	1,484	1,634	1,830	1,702	1,893	1,979	1,886
Attention-deficit, conduct, and disruptive behavior disorders	160	197	250	296	321	353	422	411	492	517	507
Delirium, dementia, and amnesic and other cognitive disorders	466	503	532	538	560	562	600	547	558	555	514
Developmental disorders	216	238	261	274	299	324	356	348	369	403	390
Disorders usually diagnosed in infancy, childhood, adolescence	33	35	38	46	50	61	72	71	83	102	102
Impulse control disorders, NEC	40	48	55	58	62	71	83	86	97	107	105
Miscellaneous disorders	410	479	569	623	671	731	825	713	793	831	734
Mood disorders	1,178	1,365	1,540	1,658	1,803	1,928	2,149	1,983	2,198	2,288	2,215
Personality disorders	182	202	211	236	266	287	334	302	313	311	292
Schizophrenia and other psychotic disorders	439	493	540	559	581	608	655	622	642	631	580
Screening and history of mental health and substance abuse	974	1,118	1,294	1,426	1,569	1,715	1,944	1,771	2,034	2,163	2,065
Substance-related disorders	334	399	454	520	558	612	714	660	749	790	755
Suicide and intentional self-inflicted injury	100	110	126	128	145	166	199	195	211	222	219



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Bourbon County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	841	936	1,011	1,046	1,087	1,093	1,157	1,102	1,181	1,219	1,149
Female	435	469	510	515	523	521	548	510	528	561	517
Male	406	467	501	531	564	572	609	592	653	658	632
White	685	763	823	858	894	895	941	896	962	1,002	935
Minority	119	130	137	136	131	134	139	128	139	135	129

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	58%	62%	61%	60%	59%	60%	55%	55%	50%	45%
Female	60%	59%	63%	61%	60%	59%	59%	52%	51%	47%	42%
Male	56%	57%	60%	61%	60%	59%	60%	57%	58%	53%	48%
White	57%	58%	62%	60%	60%	59%	59%	53%	53%	49%	43%
Minority	45%	45%	45%	45%	40%	39%	42%	38%	39%	37%	34%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	222	254	277	281	308	291	307	262	282	262	228
Alcohol-related disorders	41	39	42	42	42	41	39	33	34	35	31
Anxiety disorders	321	363	415	440	460	466	508	496	530	531	511
Attention-deficit, conduct, and disruptive behavior disorders	348	398	438	454	467	449	476	428	441	439	410
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	132	158	198	212	241	251	289	286	297	312	319
Disorders usually diagnosed in infancy, childhood, adolescence	79	98	106	114	125	121	131	119	117	112	106
Impulse control disorders, NEC	35	40	47	49	47	46	47	41	38	31	N.D.
Miscellaneous disorders	124	129	141	148	151	138	134	129	130	130	122
Mood disorders	296	324	354	348	343	323	316	261	271	266	226
Personality disorders	33	32	33	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	35	33	36	36	38	32	35	32	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	514	556	573	571	556	553	550	480	485	480	430
Substance-related disorders	118	119	122	121	108	108	101	86	89	85	71
Suicide and intentional self-inflicted injury	N.D.	N.D.	34	37	33	N.D.	36	32	39	39	32



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Bourbon County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	794	898	938	990	1,020	1,059	1,099	1,016	1,158	1,220	1,171
Female	626	707	741	783	809	815	857	781	893	936	907
Male	168	191	197	207	211	244	242	235	265	284	264
White	660	758	789	837	864	888	931	855	981	1,031	993
Minority	99	103	109	113	114	123	112	107	120	125	117

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	66%	70%	67%	67%	68%	67%	68%	61%	66%	66%	62%
Female	69%	73%	70%	69%	71%	70%	71%	62%	68%	68%	65%
Male	57%	62%	58%	59%	59%	60%	61%	57%	58%	61%	54%
White	67%	71%	70%	69%	71%	70%	70%	63%	68%	68%	63%
Minority	46%	49%	42%	42%	42%	41%	39%	35%	38%	40%	39%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	63	70	78	87	91	99	108	108	131	140	135
Alcohol-related disorders	59	79	84	98	99	105	108	106	108	114	109
Anxiety disorders	391	459	484	503	525	551	591	557	641	678	646
Attention-deficit, conduct, and disruptive behavior disorders	31	37	41	50	54	66	64	73	89	99	97
Delirium, dementia, and amnesic and other cognitive disorders	201	210	211	214	216	202	188	151	158	160	146
Developmental disorders	63	71	71	68	67	72	75	65	75	86	79
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	36	34	31
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	168	205	212	214	231	246	277	258	295	319	308
Mood disorders	448	495	526	555	580	613	669	635	710	740	727
Personality disorders	91	107	111	119	131	137	133	126	141	140	134
Schizophrenia and other psychotic disorders	154	170	172	174	163	171	176	155	166	158	150
Screening and history of mental health and substance abuse	452	533	574	624	664	676	729	690	794	849	808
Substance-related disorders	152	178	187	207	233	257	274	268	309	330	331
Suicide and intentional self-inflicted injury	32	42	39	38	45	50	60	58	72	66	66



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Boyd County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,772	3,026	3,102	3,230	3,324	3,367	3,409	3,132	3,283	3,225	3,032
Female	1,334	1,413	1,455	1,517	1,536	1,520	1,533	1,427	1,507	1,463	1,376
Male	1,438	1,613	1,647	1,713	1,788	1,847	1,876	1,705	1,776	1,762	1,656
White	2,514	2,748	2,811	2,902	2,971	3,004	3,016	2,776	2,894	2,844	2,678
Minority	134	134	123	121	122	113	114	101	115	112	108

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	56%	58%	58%	58%	57%	59%	55%	56%	54%	49%
Female	54%	55%	56%	56%	55%	53%	55%	51%	52%	49%	45%
Male	54%	58%	59%	61%	61%	61%	63%	58%	60%	58%	53%
White	53%	55%	57%	57%	56%	56%	57%	53%	54%	52%	48%
Minority	35%	33%	30%	27%	25%	23%	24%	21%	23%	20%	19%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	559	604	642	675	697	714	732	679	695	677	613
Alcohol-related disorders	164	150	147	147	135	121	111	89	77	74	59
Anxiety disorders	957	992	1,006	1,029	1,026	1,030	1,015	909	915	883	804
Attention-deficit, conduct, and disruptive behavior disorders	1,486	1,630	1,696	1,805	1,892	1,948	1,964	1,809	1,846	1,738	1,627
Delirium, dementia, and amnestic and other cognitive disorders	37	33	36	34	33	31	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	622	727	831	943	1,058	1,142	1,210	1,173	1,274	1,289	1,237
Disorders usually diagnosed in infancy, childhood, adolescence	268	307	345	372	407	419	444	419	420	400	381
Impulse control disorders, NEC	125	134	137	141	131	125	126	109	97	82	63
Miscellaneous disorders	285	301	299	309	314	318	342	335	343	323	306
Mood disorders	1,217	1,277	1,269	1,282	1,247	1,175	1,136	987	932	825	746
Personality disorders	125	124	122	112	109	100	94	72	67	63	50
Schizophrenia and other psychotic disorders	130	129	126	123	129	133	124	98	89	82	72
Screening and history of mental health and substance abuse	2,062	2,224	2,265	2,303	2,335	2,311	2,286	2,034	2,077	1,982	1,835
Substance-related disorders	421	411	386	352	320	302	272	226	220	204	182
Suicide and intentional self-inflicted injury	108	111	107	102	100	98	98	85	79	72	65



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Boyd County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,120	3,405	3,654	3,819	3,979	4,083	4,326	3,985	4,245	4,258	4,151
Female	2,234	2,437	2,630	2,713	2,823	2,901	3,043	2,771	2,962	2,964	2,878
Male	886	968	1,024	1,106	1,156	1,182	1,283	1,214	1,283	1,294	1,273
White	2,822	3,085	3,326	3,472	3,628	3,724	3,973	3,649	3,884	3,893	3,770
Minority	133	141	134	139	121	100	81	77	97	91	91

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	67%	68%	69%	68%	68%	70%	65%	70%	68%	65%
Female	67%	69%	71%	72%	71%	71%	73%	68%	72%	71%	68%
Male	59%	60%	62%	63%	62%	63%	65%	60%	65%	62%	61%
White	64%	66%	68%	69%	68%	68%	70%	65%	69%	68%	65%
Minority	32%	31%	30%	31%	26%	21%	18%	16%	20%	17%	15%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	257	288	317	344	375	401	436	403	437	446	445
Alcohol-related disorders	459	509	510	571	594	623	653	599	617	604	584
Anxiety disorders	1,918	2,138	2,289	2,421	2,525	2,621	2,767	2,611	2,715	2,696	2,635
Attention-deficit, conduct, and disruptive behavior disorders	193	223	265	290	314	345	414	403	446	484	498
Delirium, dementia, and amnesic and other cognitive disorders	564	586	625	642	639	625	676	589	586	571	538
Developmental disorders	347	370	403	437	457	459	487	454	489	507	526
Disorders usually diagnosed in infancy, childhood, adolescence	45	46	50	55	60	68	72	83	90	104	118
Impulse control disorders, NEC	65	68	79	92	101	103	116	122	130	147	154
Miscellaneous disorders	590	648	694	741	762	784	838	771	810	788	754
Mood disorders	2,065	2,291	2,502	2,617	2,783	2,893	3,051	2,837	2,943	2,946	2,854
Personality disorders	520	563	607	646	675	669	684	641	638	653	641
Schizophrenia and other psychotic disorders	727	788	854	905	935	925	955	898	910	863	819
Screening and history of mental health and substance abuse	1,917	2,101	2,317	2,457	2,607	2,681	2,861	2,673	2,857	2,900	2,880
Substance-related disorders	707	804	861	951	1,013	1,073	1,169	1,117	1,207	1,227	1,181
Suicide and intentional self-inflicted injury	119	135	152	168	194	209	243	239	261	278	270



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Boyle County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,165	1,331	1,446	1,517	1,545	1,600	1,596	1,509	1,631	1,618	1,525
Female	564	659	718	737	720	758	765	706	766	768	718
Male	601	672	728	780	825	842	831	803	865	850	807
White	934	1,069	1,161	1,223	1,245	1,276	1,255	1,203	1,303	1,283	1,222
Minority	163	184	198	201	206	220	223	192	206	208	192

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	58%	62%	61%	60%	62%	60%	57%	59%	54%	48%
Female	56%	58%	62%	60%	57%	60%	58%	53%	55%	50%	44%
Male	58%	59%	62%	62%	62%	64%	62%	60%	63%	58%	52%
White	57%	58%	62%	61%	60%	62%	60%	56%	58%	53%	47%
Minority	40%	41%	44%	42%	40%	42%	40%	37%	38%	36%	34%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	437	491	528	529	533	541	538	491	512	480	442
Alcohol-related disorders	46	48	50	57	59	51	45	37	35	N.D.	N.D.
Anxiety disorders	471	531	592	646	654	702	697	652	707	668	627
Attention-deficit, conduct, and disruptive behavior disorders	612	687	729	772	788	805	805	724	744	727	684
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	308	386	424	459	485	514	521	506	545	559	547
Disorders usually diagnosed in infancy, childhood, adolescence	118	137	143	146	161	169	181	168	170	169	158
Impulse control disorders, NEC	50	52	54	45	52	47	46	46	45	31	N.D.
Miscellaneous disorders	199	226	244	243	248	250	261	247	257	260	230
Mood disorders	395	405	415	403	385	375	356	296	284	258	226
Personality disorders	41	46	45	40	43	38	32	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	51	53	52	50	53	53	50	44	47	40	38
Screening and history of mental health and substance abuse	586	632	681	653	629	613	594	543	542	521	475
Substance-related disorders	113	115	114	108	101	93	79	65	60	51	39
Suicide and intentional self-inflicted injury	46	53	47	48	48	45	49	46	46	43	34



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Boyle County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,394	1,498	1,587	1,677	1,705	1,734	1,770	1,648	1,784	1,822	1,806
Female	991	1,078	1,149	1,205	1,242	1,247	1,277	1,181	1,284	1,291	1,263
Male	403	420	438	472	463	487	493	467	500	531	543
White	1,187	1,276	1,354	1,430	1,459	1,471	1,520	1,409	1,534	1,546	1,520
Minority	154	170	176	189	186	196	177	163	176	196	191

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	66%	68%	69%	69%	67%	69%	64%	68%	67%	64%
Female	66%	68%	70%	71%	71%	69%	70%	65%	70%	69%	65%
Male	62%	62%	64%	65%	63%	64%	65%	61%	63%	62%	60%
White	66%	68%	70%	70%	70%	69%	71%	65%	69%	67%	64%
Minority	44%	46%	45%	48%	46%	44%	42%	40%	44%	46%	43%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	208	230	269	292	312	330	337	306	349	370	368
Alcohol-related disorders	125	138	140	152	158	168	168	151	158	161	159
Anxiety disorders	795	888	980	1,041	1,071	1,100	1,155	1,066	1,131	1,162	1,147
Attention-deficit, conduct, and disruptive behavior disorders	112	126	141	156	156	173	189	171	200	219	238
Delirium, dementia, and amnesic and other cognitive disorders	434	432	420	414	390	375	351	313	324	322	292
Developmental disorders	168	172	185	195	188	193	213	201	213	228	239
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33	36	36	38	43
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	N.D.	32	42	48
Miscellaneous disorders	331	361	414	451	469	470	482	437	448	479	469
Mood disorders	843	933	1,022	1,074	1,115	1,147	1,188	1,108	1,162	1,178	1,155
Personality disorders	189	208	236	251	251	252	260	236	239	242	242
Schizophrenia and other psychotic disorders	426	457	463	463	460	452	444	409	408	404	389
Screening and history of mental health and substance abuse	667	750	831	904	908	913	941	876	938	939	932
Substance-related disorders	192	220	246	272	284	300	317	278	328	328	339
Suicide and intentional self-inflicted injury	68	79	86	97	101	107	116	97	117	127	143



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Bracken County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	437	471	531	552	583	600	622	586	575	598	548
Female	230	239	267	278	281	287	301	283	274	277	259
Male	207	232	264	274	302	313	321	303	301	321	289
White	399	427	480	497	519	537	560	532	521	549	499
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	58%	62%	60%	59%	61%	59%	54%	53%	52%	47%
Female	63%	61%	64%	61%	58%	60%	60%	54%	52%	51%	45%
Male	53%	55%	61%	59%	60%	61%	58%	54%	54%	53%	48%
White	57%	56%	60%	58%	56%	58%	57%	52%	52%	50%	45%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	108	115	129	128	141	133	142	145	133	129	116
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	124	131	147	143	151	147	157	143	134	131	127
Attention-deficit, conduct, and disruptive behavior disorders	182	195	222	234	254	265	268	246	243	253	229
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	134	155	180	200	241	254	261	243	250	244	237
Disorders usually diagnosed in infancy, childhood, adolescence	35	44	51	54	60	57	59	49	51	50	44
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	120	134	145	151	162	162	158	145	141	132	119
Mood disorders	137	150	162	152	148	138	140	123	113	103	91
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	332	353	383	383	389	381	375	345	328	332	296
Substance-related disorders	37	45	44	43	44	45	43	36	32	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Bracken County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	416	469	497	525	548	554	576	511	602	630	602
Female	300	331	359	388	396	404	424	375	440	455	429
Male	116	138	138	137	152	150	152	136	162	175	173
White	384	436	464	488	514	522	542	N.D.	570	599	566
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	69%	67%	68%	69%	69%	72%	59%	67%	68%	66%
Female	67%	70%	72%	74%	74%	73%	75%	63%	71%	73%	69%
Male	60%	66%	58%	55%	59%	60%	64%	51%	56%	59%	59%
White	64%	69%	67%	67%	69%	69%	71%	59%	67%	68%	66%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	42	49	60	56	62	63	69	60	77	98	97
Alcohol-related disorders	48	48	51	52	50	54	58	54	65	63	61
Anxiety disorders	218	249	277	299	311	321	331	294	331	343	321
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	32	35	41	43	43	45	57	55	59
Delirium, dementia, and amnestic and other cognitive disorders	80	89	91	86	87	77	73	72	78	73	70
Developmental disorders	49	52	51	54	56	63	62	52	60	66	61
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	68	75	84	103	116	127	129	102	132	136	129
Mood disorders	209	238	260	274	291	312	319	277	335	357	337
Personality disorders	33	34	40	43	49	50	48	43	47	51	56
Schizophrenia and other psychotic disorders	64	74	76	81	82	86	86	78	78	77	77
Screening and history of mental health and substance abuse	271	317	344	376	391	405	413	361	432	460	435
Substance-related disorders	44	54	62	65	70	74	74	63	80	81	77
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	34	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Breathitt County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,712	1,776	1,801	1,828	1,804	1,792	1,813	1,683	1,739	1,701	1,576
Female	885	906	890	877	857	840	834	775	801	788	718
Male	827	870	911	951	947	952	979	908	938	913	858
White	1,593	1,653	1,670	1,693	1,658	1,635	1,629	1,510	1,559	1,518	1,407
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	60%	62%	63%	62%	62%	63%	60%	61%	60%	56%
Female	61%	62%	63%	63%	62%	62%	61%	58%	59%	58%	56%
Male	54%	57%	62%	63%	62%	62%	65%	61%	63%	61%	56%
White	56%	58%	61%	61%	61%	60%	61%	58%	59%	58%	54%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	297	311	332	339	328	335	352	318	327	319	289
Alcohol-related disorders	136	129	117	119	108	101	94	78	78	72	56
Anxiety disorders	732	751	745	769	746	747	742	696	685	658	585
Attention-deficit, conduct, and disruptive behavior disorders	586	602	627	651	646	651	670	614	608	582	530
Delirium, dementia, and amnesic and other cognitive disorders	33	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	463	506	566	623	647	656	690	673	718	717	684
Disorders usually diagnosed in infancy, childhood, adolescence	129	138	157	164	167	164	180	172	176	168	156
Impulse control disorders, NEC	53	53	58	59	47	46	46	45	42	40	34
Miscellaneous disorders	205	198	193	179	168	157	160	145	138	140	128
Mood disorders	556	543	515	500	459	434	400	349	330	304	252
Personality disorders	105	102	109	114	108	106	99	90	80	73	65
Schizophrenia and other psychotic disorders	70	66	63	58	51	48	46	39	37	35	N.D.
Screening and history of mental health and substance abuse	1,146	1,176	1,153	1,141	1,090	1,041	1,017	902	900	856	777
Substance-related disorders	275	271	247	239	208	202	197	174	167	159	135
Suicide and intentional self-inflicted injury	38	37	35	34	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Breathitt County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,558	2,705	2,852	2,897	2,919	2,901	2,906	2,652	2,808	2,786	2,686
Female	1,595	1,685	1,764	1,785	1,783	1,789	1,783	1,612	1,698	1,695	1,627
Male	963	1,020	1,088	1,112	1,136	1,112	1,123	1,040	1,110	1,091	1,059
White	2,385	2,524	2,668	2,713	2,743	2,725	2,742	2,489	2,651	2,631	2,529
Minority	46	44	36	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	70%	71%	73%	73%	73%	73%	75%	69%	74%	75%	72%
Female	74%	76%	78%	78%	78%	78%	79%	73%	78%	80%	76%
Male	63%	64%	66%	68%	66%	67%	69%	64%	68%	69%	66%
White	69%	71%	73%	73%	73%	73%	74%	69%	74%	75%	72%
Minority	20%	19%	15%	14%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	212	223	245	265	279	281	300	285	301	302	299
Alcohol-related disorders	246	269	284	306	315	314	323	292	300	302	292
Anxiety disorders	1,824	1,928	2,072	2,132	2,177	2,169	2,173	1,996	2,107	2,064	2,004
Attention-deficit, conduct, and disruptive behavior disorders	75	84	94	115	125	129	149	141	170	177	197
Delirium, dementia, and amnesic and other cognitive disorders	541	550	554	519	513	468	435	399	387	358	326
Developmental disorders	252	260	265	280	277	281	297	275	286	298	300
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	31	33	36	34	35	34	34	41	45	48
Impulse control disorders, NEC	52	56	59	70	76	75	78	71	75	75	74
Miscellaneous disorders	686	738	806	858	878	879	898	830	835	813	763
Mood disorders	1,478	1,573	1,679	1,745	1,777	1,770	1,810	1,654	1,737	1,720	1,655
Personality disorders	103	108	118	120	124	123	128	116	120	118	128
Schizophrenia and other psychotic disorders	588	607	612	605	593	586	583	540	539	513	488
Screening and history of mental health and substance abuse	1,527	1,663	1,780	1,852	1,901	1,925	1,955	1,754	1,884	1,878	1,810
Substance-related disorders	577	631	689	753	781	793	829	762	805	802	799
Suicide and intentional self-inflicted injury	101	104	113	116	121	128	139	124	137	137	138



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Breckinridge County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	944	1,044	1,120	1,178	1,220	1,218	1,234	1,172	1,213	1,238	1,173
Female	463	527	572	581	595	596	595	571	598	581	560
Male	481	517	548	597	625	622	639	601	615	657	613
White	843	926	983	1,040	1,082	1,079	1,083	1,021	1,064	1,082	1,027
Minority	52	56	63	60	57	55	57	53	51	53	49

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	51%	52%	52%	53%	54%	54%	53%	50%	50%	47%	43%
Female	51%	52%	53%	53%	54%	55%	52%	49%	48%	45%	41%
Male	51%	52%	51%	53%	53%	53%	54%	52%	51%	49%	44%
White	50%	50%	50%	51%	52%	52%	51%	48%	48%	45%	41%
Minority	34%	31%	33%	31%	27%	27%	29%	26%	25%	23%	21%


CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	238	269	298	306	306	311	312	295	303	309	291
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	287	321	353	365	378	371	384	371	379	379	354
Attention-deficit, conduct, and disruptive behavior disorders	403	448	507	539	565	565	593	556	579	578	553
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	269	293	336	364	378	390	419	416	414	429	405
Disorders usually diagnosed in infancy, childhood, adolescence	74	81	99	112	122	118	124	126	137	131	117
Impulse control disorders, NEC	85	95	99	95	92	92	88	80	71	64	57
Miscellaneous disorders	98	101	101	102	105	100	101	95	91	89	90
Mood disorders	313	335	357	356	343	322	305	275	265	254	234
Personality disorders	40	40	45	41	42	42	41	36	37	33	31
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	617	668	663	682	668	634	602	559	544	530	488
Substance-related disorders	66	65	72	70	70	63	52	46	38	39	33
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Breckinridge County											
ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	923	990	1,093	1,243	1,286	1,312	1,342	1,280	1,368	1,355	1,270
Female	678	721	801	917	927	933	923	898	958	959	889
Male	245	269	292	326	359	379	419	382	410	396	381
White	821	881	982	1,128	1,166	1,201	1,233	1,174	1,252	1,237	1,145
Minority	55	54	52	53	55	43	38	32	39	44	46
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	56%	59%	63%	65%	64%	65%	60%	62%	63%	59%
Female	59%	61%	64%	68%	69%	68%	68%	64%	67%	68%	64%
Male	46%	47%	50%	52%	55%	56%	59%	52%	54%	54%	50%
White	55%	56%	59%	63%	65%	64%	65%	59%	62%	63%	58%
Minority	30%	29%	29%	30%	30%	26%	23%	19%	23%	26%	24%
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	97	107	118	130	131	139	143	140	147	152	143
Alcohol-related disorders	63	83	83	101	107	111	113	101	113	100	91
Anxiety disorders	509	569	647	728	748	787	794	770	817	815	767
Attention-deficit, conduct, and disruptive behavior disorders	43	50	58	67	78	87	100	108	116	128	123
Delirium, dementia, and amnestic and other cognitive disorders	221	228	232	243	243	237	232	209	194	190	169
Developmental disorders	110	106	112	127	138	129	133	130	145	138	146
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	31	38	42	47	48	48	53	49	49
Miscellaneous disorders	159	184	211	245	245	246	249	233	253	251	235
Mood disorders	522	575	648	733	760	800	835	806	879	869	824
Personality disorders	72	81	90	102	107	110	115	106	111	115	115
Schizophrenia and other psychotic disorders	191	202	214	230	239	240	239	229	228	219	217
Screening and history of mental health and substance abuse	438	488	551	669	718	731	737	691	738	739	673
Substance-related disorders	115	133	158	188	190	202	208	186	202	203	185
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	33	38	50	45



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Bullitt County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,937	2,217	2,403	2,625	2,741	2,763	2,842	2,673	2,822	2,882	2,735
Female	942	1,070	1,158	1,257	1,289	1,293	1,315	1,217	1,277	1,295	1,204
Male	995	1,147	1,245	1,368	1,452	1,470	1,527	1,456	1,545	1,587	1,531
White	1,814	2,065	2,241	2,447	2,533	2,554	2,612	2,436	2,575	2,622	2,490
Minority	38	54	56	55	62	52	48	43	49	52	49

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	47%	47%	50%	50%	50%	49%	49%	46%	47%	44%	39%
Female	46%	47%	48%	48%	48%	46%	45%	42%	42%	40%	35%
Male	48%	48%	51%	53%	52%	52%	53%	50%	51%	48%	43%
White	46%	47%	49%	50%	49%	48%	48%	44%	45%	43%	37%
Minority	19%	22%	21%	19%	20%	16%	14%	12%	14%	13%	12%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	552	625	701	752	754	750	744	657	696	702	642
Alcohol-related disorders	78	89	92	105	101	98	95	75	69	58	57
Anxiety disorders	574	650	683	737	756	758	781	691	704	684	615
Attention-deficit, conduct, and disruptive behavior disorders	1,008	1,153	1,260	1,399	1,487	1,494	1,532	1,426	1,477	1,478	1,389
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	33	38	41	42	40	46	47	48	51	43
Developmental disorders	275	351	406	470	516	551	597	623	673	721	702
Disorders usually diagnosed in infancy, childhood, adolescence	191	234	273	301	354	381	421	407	439	423	405
Impulse control disorders, NEC	153	161	175	188	186	188	185	164	167	149	122
Miscellaneous disorders	239	277	282	310	320	319	344	324	329	326	300
Mood disorders	781	873	924	974	980	974	953	842	833	772	695
Personality disorders	53	59	62	63	57	63	57	50	55	53	47
Schizophrenia and other psychotic disorders	82	89	99	104	101	107	107	89	95	91	77
Screening and history of mental health and substance abuse	961	1,075	1,137	1,162	1,179	1,124	1,101	978	994	941	854
Substance-related disorders	208	226	234	257	228	234	218	170	166	148	134
Suicide and intentional self-inflicted injury	77	95	111	115	115	118	120	109	111	111	94



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Bullitt County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,239	1,383	1,539	1,645	1,816	1,912	2,062	1,943	2,154	2,289	2,289
Female	945	1,063	1,181	1,262	1,386	1,449	1,547	1,438	1,600	1,724	1,707
Male	294	320	358	383	430	463	515	505	554	565	582
White	1,143	1,274	1,415	1,523	1,691	1,790	1,928	1,813	2,017	2,148	2,142
Minority	41	44	50	44	42	40	36	34	41	42	34

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	56%	58%	60%	63%	62%	64%	60%	65%	64%	61%
Female	55%	58%	60%	62%	65%	65%	66%	61%	67%	67%	63%
Male	49%	50%	52%	53%	58%	55%	60%	57%	61%	58%	56%
White	53%	55%	57%	59%	62%	62%	64%	59%	65%	64%	61%
Minority	25%	25%	28%	25%	24%	22%	20%	18%	21%	20%	15%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	129	150	174	185	212	229	259	247	257	280	287
Alcohol-related disorders	115	120	146	160	182	189	215	192	209	230	229
Anxiety disorders	616	684	781	860	947	1,011	1,142	1,103	1,209	1,300	1,292
Attention-deficit, conduct, and disruptive behavior disorders	102	111	137	155	162	182	220	241	293	329	340
Delirium, dementia, and amnestic and other cognitive disorders	272	278	283	268	264	258	264	257	241	242	252
Developmental disorders	102	102	125	126	130	136	151	149	169	171	183
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	35	41	45	54	61	69
Impulse control disorders, NEC	32	31	41	45	51	47	64	61	68	85	85
Miscellaneous disorders	204	238	275	333	391	434	466	406	469	494	479
Mood disorders	749	837	938	983	1,086	1,162	1,286	1,241	1,390	1,504	1,485
Personality disorders	82	87	92	101	107	109	129	119	130	133	125
Schizophrenia and other psychotic disorders	274	279	312	308	316	330	365	345	350	364	352
Screening and history of mental health and substance abuse	691	779	883	992	1,162	1,254	1,350	1,263	1,392	1,465	1,429
Substance-related disorders	195	227	258	287	325	352	411	390	442	467	453
Suicide and intentional self-inflicted injury	63	71	84	94	103	111	129	133	153	163	162



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Butler County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	737	794	845	879	900	934	920	855	903	895	833
Female	361	387	416	427	420	432	423	393	414	413	393
Male	376	407	429	452	480	502	497	462	489	482	440
White	701	755	798	831	848	878	862	803	848	843	785
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	55%	56%	55%	53%	54%	54%	51%	53%	50%	45%
Female	57%	58%	59%	56%	52%	54%	53%	48%	50%	46%	43%
Male	51%	53%	53%	54%	54%	55%	55%	53%	56%	53%	47%
White	53%	55%	55%	54%	52%	54%	53%	50%	52%	49%	44%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	189	198	213	205	211	208	193	163	169	158	136
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	278	296	314	330	343	357	360	328	345	322	298
Attention-deficit, conduct, and disruptive behavior disorders	346	374	394	417	442	452	450	414	421	409	371
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	99	115	129	148	172	189	204	190	212	207	204
Disorders usually diagnosed in infancy, childhood, adolescence	45	53	55	58	63	69	67	60	66	59	60
Impulse control disorders, NEC	31	35	36	38	37	36	35	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	84	93	100	94	97	92	91	83	84	77	62
Mood disorders	272	282	310	295	286	273	253	220	208	193	168
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	480	515	533	536	516	500	474	408	427	415	355
Substance-related disorders	75	75	70	65	59	57	53	44	46	47	37
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Butler County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	906	955	1,031	1,046	1,075	1,072	1,108	1,037	1,107	1,111	1,082
Female	635	664	719	734	755	753	761	716	766	784	743
Male	271	291	312	312	320	319	347	321	341	327	339
White	835	882	956	966	1,003	996	1,025	954	1,021	1,024	998
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	66%	68%	71%	70%	71%	69%	70%	64%	70%	70%	67%
Female	70%	71%	74%	74%	73%	72%	72%	66%	74%	74%	70%
Male	60%	61%	64%	62%	65%	62%	64%	61%	62%	63%	62%
White	66%	68%	71%	70%	71%	69%	69%	64%	69%	70%	67%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	102	130	137	144	147	153	174	164	178	190	185
Alcohol-related disorders	92	99	113	106	96	103	96	84	95	95	98
Anxiety disorders	513	553	610	626	644	656	695	665	703	711	709
Attention-deficit, conduct, and disruptive behavior disorders	48	48	55	59	71	79	94	96	119	131	142
Delirium, dementia, and amnesic and other cognitive disorders	266	267	256	232	225	222	227	226	213	186	174
Developmental disorders	83	89	91	97	87	87	83	81	86	86	86
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	168	194	210	231	234	253	271	237	256	264	262
Mood disorders	546	589	647	674	687	698	752	713	755	768	747
Personality disorders	51	47	50	51	49	50	55	50	49	54	53
Schizophrenia and other psychotic disorders	198	206	210	208	202	208	213	210	205	189	185
Screening and history of mental health and substance abuse	524	568	639	659	684	690	697	642	695	709	700
Substance-related disorders	109	129	152	171	165	165	167	153	166	166	172
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	38	34	37	42	51



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Caldwell County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	715	760	806	833	810	803	809	752	775	769	711
Female	357	383	391	396	390	393	397	361	372	364	325
Male	358	377	415	437	420	410	412	391	403	405	386
White	613	652	687	708	683	673	681	637	651	637	594
Minority	66	66	73	75	73	72	65	54	54	60	51

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	58%	58%	60%	57%	57%	56%	51%	51%	47%	41%
Female	56%	57%	55%	55%	54%	55%	55%	48%	48%	46%	39%
Male	58%	60%	62%	64%	59%	59%	58%	53%	54%	49%	43%
White	56%	57%	57%	59%	55%	55%	54%	49%	49%	46%	40%
Minority	41%	40%	39%	38%	39%	39%	35%	29%	28%	25%	21%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	102	110	118	123	115	123	128	122	117	104	102
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	212	225	223	237	229	233	239	217	212	193	174
Attention-deficit, conduct, and disruptive behavior disorders	324	344	374	392	387	382	393	377	389	366	339
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	154	185	207	223	230	238	237	225	226	228	225
Disorders usually diagnosed in infancy, childhood, adolescence	32	35	38	40	37	42	47	46	50	62	60
Impulse control disorders, NEC	77	81	84	89	88	90	85	80	74	65	57
Miscellaneous disorders	112	112	119	120	106	103	104	93	90	88	74
Mood disorders	277	270	285	282	269	255	243	218	205	190	164
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	406	418	440	422	400	393	378	354	349	321	276
Substance-related disorders	45	41	47	49	50	49	44	39	41	31	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Caldwell County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	798	876	934	943	999	1,013	1,009	924	1,043	1,051	1,011
Female	576	627	663	664	704	697	687	618	706	701	667
Male	222	249	271	279	295	316	322	306	337	350	344
White	691	762	819	825	877	888	884	808	911	918	868
Minority	80	86	88	89	89	90	84	77	90	92	95

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	67%	69%	69%	70%	70%	71%	63%	70%	70%	64%
Female	66%	69%	71%	70%	72%	71%	72%	63%	70%	70%	64%
Male	56%	61%	66%	67%	68%	69%	68%	65%	71%	70%	63%
White	63%	67%	69%	69%	70%	70%	70%	62%	69%	70%	64%
Minority	46%	50%	52%	53%	52%	51%	49%	47%	52%	47%	43%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	52	61	65	59	63	60	63	62	82	92	80
Alcohol-related disorders	48	55	63	66	71	75	78	69	80	84	77
Anxiety disorders	392	437	473	503	543	559	581	533	597	596	571
Attention-deficit, conduct, and disruptive behavior disorders	41	54	62	72	78	85	94	88	110	129	126
Delirium, dementia, and amnestic and other cognitive disorders	213	226	223	215	216	206	198	181	181	162	141
Developmental disorders	117	120	127	125	132	138	137	134	145	144	147
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33	43	49	49
Miscellaneous disorders	190	205	219	224	254	259	254	219	249	248	240
Mood disorders	468	529	563	572	609	615	644	593	669	674	641
Personality disorders	33	40	34	36	41	48	46	43	45	47	47
Schizophrenia and other psychotic disorders	196	208	221	226	228	232	239	222	238	228	217
Screening and history of mental health and substance abuse	378	419	483	496	535	571	570	516	593	601	594
Substance-related disorders	80	91	108	112	117	127	128	110	128	134	124
Suicide and intentional self-inflicted injury	N.D.	32	40	41	45	50	51	47	57	61	58



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Calloway County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,066	1,170	1,309	1,363	1,384	1,417	1,387	1,281	1,357	1,381	1,281
Female	550	608	674	674	673	687	681	608	630	645	613
Male	516	562	635	689	711	730	706	673	727	736	668
White	927	1,013	1,132	1,177	1,189	1,199	1,162	1,087	1,151	1,171	1,085
Minority	96	104	109	108	110	123	125	102	112	112	103

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	46%	47%	50%	50%	50%	49%	47%	44%	45%	43%	39%
Female	47%	50%	52%	50%	49%	48%	46%	42%	43%	41%	37%
Male	44%	45%	49%	50%	50%	49%	47%	45%	47%	45%	40%
White	45%	47%	49%	49%	49%	48%	45%	43%	44%	41%	37%
Minority	34%	35%	35%	33%	32%	30%	30%	26%	28%	28%	25%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	196	216	244	252	250	244	229	207	199	184	171
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	252	270	291	289	287	287	272	259	267	250	220
Attention-deficit, conduct, and disruptive behavior disorders	513	552	604	657	677	685	661	623	640	649	600
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	137	160	195	220	240	272	296	306	349	368	339
Disorders usually diagnosed in infancy, childhood, adolescence	77	85	101	112	112	122	122	115	121	122	111
Impulse control disorders, NEC	74	72	80	84	85	77	67	58	55	55	40
Miscellaneous disorders	204	216	233	238	243	248	233	202	197	192	172
Mood disorders	358	379	407	394	387	379	360	314	330	323	279
Personality disorders	52	57	52	46	41	33	35	34	34	33	N.D.
Schizophrenia and other psychotic disorders	47	37	42	40	38	37	35	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	528	573	646	653	660	657	619	558	580	581	521
Substance-related disorders	73	68	72	64	68	56	51	50	41	47	37
Suicide and intentional self-inflicted injury	31	N.D.	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Calloway County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,101	1,212	1,285	1,291	1,352	1,413	1,478	1,327	1,453	1,470	1,419
Female	844	916	975	965	1,003	1,050	1,098	970	1,065	1,067	1,015
Male	257	296	310	326	349	363	380	357	388	403	404
White	982	1,087	1,151	1,164	1,220	1,273	1,327	1,195	1,310	1,316	1,265
Minority	83	90	90	81	83	85	87	70	82	87	87

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	60%	61%	60%	61%	60%	61%	57%	60%	60%	56%
Female	57%	62%	63%	61%	62%	62%	64%	58%	62%	61%	56%
Male	51%	56%	55%	55%	56%	55%	55%	53%	56%	56%	54%
White	56%	61%	62%	61%	61%	61%	62%	57%	60%	60%	56%
Minority	37%	41%	38%	34%	35%	31%	34%	30%	33%	31%	29%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	65	73	83	89	101	107	132	120	128	135	119
Alcohol-related disorders	56	61	62	58	81	86	102	91	92	97	97
Anxiety disorders	490	525	568	583	634	682	747	696	735	773	768
Attention-deficit, conduct, and disruptive behavior disorders	61	67	78	95	122	130	150	152	180	188	194
Delirium, dementia, and amnestic and other cognitive disorders	290	305	310	314	297	289	290	256	270	243	230
Developmental disorders	91	104	113	115	116	120	130	124	126	134	136
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33	35
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	39	38	44	43	57	58	61
Miscellaneous disorders	198	226	253	262	285	299	327	281	316	319	305
Mood disorders	567	628	686	722	792	799	867	810	865	909	888
Personality disorders	75	111	118	101	109	110	118	109	114	114	111
Schizophrenia and other psychotic disorders	194	218	228	236	238	252	261	259	264	245	226
Screening and history of mental health and substance abuse	385	465	495	505	559	582	625	551	599	622	601
Substance-related disorders	110	122	127	127	144	154	175	151	183	195	189
Suicide and intentional self-inflicted injury	39	45	49	52	54	58	77	82	88	93	92



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Campbell County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,673	2,987	3,127	3,223	3,316	3,461	3,570	3,326	3,502	3,546	3,346
Female	1,291	1,433	1,486	1,534	1,552	1,602	1,655	1,515	1,600	1,597	1,483
Male	1,382	1,554	1,641	1,689	1,764	1,859	1,915	1,811	1,902	1,949	1,863
White	2,295	2,545	2,645	2,722	2,803	2,929	2,993	2,781	2,911	2,941	2,764
Minority	234	267	284	278	278	267	275	273	300	304	305

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	58%	59%	59%	58%	58%	58%	53%	54%	51%	45%
Female	54%	57%	59%	58%	56%	56%	55%	50%	51%	47%	42%
Male	57%	59%	60%	59%	59%	60%	61%	56%	57%	55%	49%
White	55%	58%	58%	58%	58%	58%	58%	53%	54%	50%	45%
Minority	35%	36%	38%	33%	32%	30%	29%	27%	30%	27%	25%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	525	570	592	606	606	642	639	580	578	564	506
Alcohol-related disorders	144	153	153	146	150	136	120	98	94	86	65
Anxiety disorders	719	769	787	777	786	782	794	691	672	640	590
Attention-deficit, conduct, and disruptive behavior disorders	1,378	1,564	1,697	1,790	1,886	1,989	2,058	1,924	2,011	2,009	1,881
Delirium, dementia, and amnestic and other cognitive disorders	65	64	71	69	69	74	73	60	60	67	58
Developmental disorders	727	852	952	1,033	1,103	1,191	1,280	1,219	1,312	1,348	1,270
Disorders usually diagnosed in infancy, childhood, adolescence	235	289	316	334	357	373	398	392	409	400	378
Impulse control disorders, NEC	145	146	165	163	147	150	159	141	127	126	114
Miscellaneous disorders	560	616	620	617	642	621	625	570	565	557	511
Mood disorders	973	1,041	1,053	1,023	1,003	974	942	804	789	749	669
Personality disorders	124	130	119	100	102	86	79	57	50	46	37
Schizophrenia and other psychotic disorders	113	121	117	115	116	109	102	96	92	90	78
Screening and history of mental health and substance abuse	1,715	1,837	1,851	1,843	1,787	1,769	1,764	1,566	1,600	1,556	1,423
Substance-related disorders	275	287	283	255	255	249	229	201	204	186	154
Suicide and intentional self-inflicted injury	92	96	97	93	94	101	90	91	85	74	72



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Campbell County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,748	2,948	3,170	3,312	3,369	3,488	3,637	3,201	3,526	3,641	3,495
Female	2,073	2,218	2,386	2,487	2,543	2,616	2,717	2,420	2,645	2,725	2,572
Male	675	730	784	825	826	872	920	781	881	916	923
White	2,486	2,668	2,883	3,004	3,065	3,173	3,320	2,921	3,202	3,312	3,183
Minority	163	183	181	198	191	196	185	151	184	187	168

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	67%	70%	70%	71%	72%	73%	65%	72%	72%	66%
Female	67%	70%	71%	72%	74%	74%	75%	67%	74%	73%	67%
Male	61%	62%	65%	65%	65%	66%	68%	60%	66%	67%	63%
White	65%	68%	70%	71%	72%	72%	74%	66%	72%	73%	68%
Minority	39%	42%	40%	41%	41%	41%	39%	31%	38%	35%	29%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	293	319	351	386	392	419	461	397	446	484	450
Alcohol-related disorders	383	427	476	497	508	540	552	469	534	559	524
Anxiety disorders	1,539	1,686	1,820	1,885	1,932	2,019	2,133	1,911	2,095	2,167	2,082
Attention-deficit, conduct, and disruptive behavior disorders	207	232	276	317	360	407	465	439	511	568	563
Delirium, dementia, and amnesic and other cognitive disorders	696	729	730	728	690	677	656	590	618	584	540
Developmental disorders	194	214	241	250	246	266	296	288	306	323	331
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	38	41	47	49	54	67	68	77	79	86
Impulse control disorders, NEC	41	49	53	59	65	71	86	85	97	103	114
Miscellaneous disorders	548	615	711	800	869	939	988	830	902	889	832
Mood disorders	1,736	1,870	2,048	2,143	2,213	2,310	2,450	2,213	2,401	2,467	2,372
Personality disorders	270	301	327	349	347	377	408	380	401	400	371
Schizophrenia and other psychotic disorders	616	662	701	718	704	733	744	663	718	705	652
Screening and history of mental health and substance abuse	1,637	1,784	2,010	2,153	2,210	2,342	2,477	2,185	2,402	2,500	2,407
Substance-related disorders	500	552	633	662	689	729	791	706	779	827	771
Suicide and intentional self-inflicted injury	138	158	182	195	201	223	268	246	288	309	298



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Carlisle County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	228	261	275	290	300	285	283	277	287	286	265
Female	115	133	137	139	138	142	141	137	139	142	127
Male	113	128	138	151	162	143	142	140	148	144	138
White	N.D.	229	243	256	256	246	240	237	245	244	229
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	50%	54%	54%	55%	55%	54%	53%	50%	50%	47%	41%
Female	49%	54%	52%	52%	51%	52%	49%	45%	44%	42%	36%
Male	51%	54%	55%	57%	59%	56%	57%	56%	57%	53%	47%
White	48%	53%	52%	53%	53%	52%	50%	47%	47%	44%	39%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	40	43	45	38	36	31	N.D.	N.D.	N.D.	N.D.	N.D.
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	43	45	47	47	46	43	43	44	42	39	36
Attention-deficit, conduct, and disruptive behavior disorders	125	144	157	157	167	160	162	152	161	160	152
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	50	63	79	91	100	99	105	97	102	100	96
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	32	33	31	33	N.D.	31	31	N.D.	N.D.	N.D.	N.D.
Mood disorders	75	81	82	78	76	72	71	62	59	55	51
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	110	116	116	115	111	103	97	104	108	106	101
Substance-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Carlisle County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	274	285	279	301	313	318	321	301	335	344	350
Female	213	222	214	231	240	238	229	221	250	241	245
Male	61	63	65	70	73	80	92	80	85	103	105
White	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	66%	69%	65%	66%	69%	68%	69%	62%	67%	65%	62%
Female	70%	73%	70%	73%	72%	71%	72%	65%	71%	66%	62%
Male	55%	58%	53%	52%	61%	60%	63%	55%	58%	62%	60%
White	66%	70%	67%	68%	70%	69%	69%	64%	68%	65%	61%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	139	143	139	145	159	159	159	157	183	185	195
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	35
Delirium, dementia, and amnesic and other cognitive disorders	92	93	87	90	82	82	76	72	73	69	63
Developmental disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	34	32	33	32
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	45	44	46	49	51	56	59	50	58	64	65
Mood disorders	172	182	185	194	204	213	220	204	223	232	243
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	60	67	67	70	68	60	58	51	49	54	55
Screening and history of mental health and substance abuse	113	127	131	146	159	165	166	151	171	182	191
Substance-related disorders	N.D.	N.D.	N.D.	31	N.D.	36	39	32	35	40	44
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Carroll County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	478	533	574	602	633	647	629	606	681	707	657
Female	230	258	273	286	293	294	279	271	297	305	283
Male	248	275	301	316	340	353	350	335	384	402	374
White	444	491	525	547	570	586	568	548	614	640	594
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	47%	48%	52%	53%	51%	49%	49%	44%	45%	42%	36%
Female	45%	48%	51%	52%	48%	47%	45%	40%	39%	37%	32%
Male	48%	49%	54%	54%	53%	51%	52%	48%	51%	46%	40%
White	46%	48%	52%	53%	50%	48%	47%	43%	44%	40%	35%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	115	133	131	136	153	144	160	153	152	160	146
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	136	144	164	175	186	189	176	162	181	178	157
Attention-deficit, conduct, and disruptive behavior disorders	233	272	294	312	346	357	347	311	341	344	323
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	77	86	101	114	125	130	133	134	159	166	159
Disorders usually diagnosed in infancy, childhood, adolescence	33	39	44	54	61	56	60	54	59	58	51
Impulse control disorders, NEC	32	34	33	38	40	41	39	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	67	70	68	67	64	62	65	53	57	59	48
Mood disorders	198	214	236	238	230	226	214	194	201	196	163
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	296	318	342	343	341	331	315	289	307	309	283
Substance-related disorders	52	49	45	47	50	48	48	33	37	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Carroll County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	566	617	662	660	694	723	720	686	783	830	810
Female	379	405	443	437	467	487	483	458	529	558	550
Male	187	212	219	223	227	236	237	228	254	272	260
White	519	566	600	596	635	661	666	635	728	772	754
Minority	N.D.	N.D.	31	34	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	63%	67%	67%	70%	69%	70%	65%	68%	67%	61%
Female	58%	63%	68%	67%	71%	70%	69%	65%	67%	66%	61%
Male	56%	64%	66%	66%	67%	68%	70%	67%	71%	69%	62%
White	57%	64%	67%	68%	71%	70%	71%	67%	70%	68%	61%
Minority	N.D.	N.D.	32%	31%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	54	53	58	57	63	65	67	72	85	92	98
Alcohol-related disorders	47	57	58	64	66	71	75	69	87	85	81
Anxiety disorders	267	290	323	343	359	369	382	378	429	449	452
Attention-deficit, conduct, and disruptive behavior disorders	34	37	42	46	44	52	67	64	70	90	94
Delirium, dementia, and amnestic and other cognitive disorders	145	145	155	150	145	148	141	130	143	142	139
Developmental disorders	62	66	70	76	70	78	83	77	86	91	88
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	102	113	130	133	153	158	156	132	143	151	147
Mood disorders	286	315	356	370	392	400	413	399	458	487	469
Personality disorders	N.D.	N.D.	35	39	37	40	42	43	51	48	45
Schizophrenia and other psychotic disorders	153	172	176	177	172	177	172	159	171	174	166
Screening and history of mental health and substance abuse	339	387	426	446	481	498	509	480	556	569	562
Substance-related disorders	70	76	86	82	90	98	107	99	124	122	118
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	34	45	40	52	51	48



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Carter County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,105	2,346	2,409	2,536	2,566	2,564	2,571	2,438	2,484	2,421	2,208
Female	1,047	1,145	1,165	1,231	1,247	1,205	1,216	1,140	1,145	1,113	1,014
Male	1,058	1,201	1,244	1,305	1,319	1,359	1,355	1,298	1,339	1,308	1,194
White	1,979	2,204	2,254	2,364	2,385	2,381	2,370	2,250	2,289	2,228	2,030
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	57%	59%	60%	60%	59%	58%	56%	58%	55%	49%
Female	56%	57%	59%	61%	60%	57%	57%	55%	56%	53%	47%
Male	55%	58%	59%	60%	60%	60%	59%	56%	59%	56%	51%
White	54%	56%	58%	59%	59%	57%	56%	54%	56%	53%	48%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	396	433	446	476	479	477	476	422	433	433	377
Alcohol-related disorders	58	58	53	49	42	36	34	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	697	766	783	820	807	803	809	758	758	738	659
Attention-deficit, conduct, and disruptive behavior disorders	796	898	959	1,020	1,042	1,065	1,094	1,006	1,034	1,016	929
Delirium, dementia, and amnesic and other cognitive disorders	32	34	33	36	N.D.	32	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	691	803	863	930	966	999	1,011	984	1,030	1,025	961
Disorders usually diagnosed in infancy, childhood, adolescence	172	197	208	231	236	244	253	245	256	252	233
Impulse control disorders, NEC	43	51	48	53	52	48	46	43	41	42	36
Miscellaneous disorders	198	217	221	232	243	241	238	226	222	218	199
Mood disorders	700	743	736	742	694	644	614	556	520	475	401
Personality disorders	55	62	59	57	55	48	46	36	32	31	N.D.
Schizophrenia and other psychotic disorders	60	57	63	66	66	61	57	56	55	51	41
Screening and history of mental health and substance abuse	1,588	1,773	1,790	1,871	1,848	1,778	1,768	1,644	1,616	1,542	1,380
Substance-related disorders	236	249	228	224	198	176	164	152	137	129	102
Suicide and intentional self-inflicted injury	49	50	49	53	50	46	49	51	45	41	33



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Carter County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,219	2,437	2,557	2,722	2,811	2,894	2,930	2,670	2,889	2,845	2,711
Female	1,545	1,685	1,744	1,858	1,906	1,951	1,983	1,762	1,888	1,854	1,748
Male	674	752	813	864	905	943	947	908	1,001	991	963
White	2,031	2,252	2,370	2,527	2,620	2,707	2,740	2,489	2,701	2,659	2,517
Minority	48	39	36	36	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	68%	69%	68%	67%	68%	62%	67%	66%	62%
Female	68%	71%	71%	73%	72%	71%	72%	65%	70%	69%	65%
Male	57%	60%	61%	63%	61%	61%	60%	58%	62%	61%	59%
White	64%	67%	67%	69%	67%	67%	67%	61%	67%	66%	62%
Minority	18%	14%	14%	14%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	141	164	165	178	189	210	217	218	237	230	241
Alcohol-related disorders	216	250	272	294	326	319	319	302	333	315	299
Anxiety disorders	1,240	1,397	1,498	1,593	1,658	1,724	1,798	1,668	1,783	1,778	1,709
Attention-deficit, conduct, and disruptive behavior disorders	76	96	103	110	124	142	172	168	186	184	190
Delirium, dementia, and amnesic and other cognitive disorders	516	517	492	492	497	469	442	404	379	354	318
Developmental disorders	268	283	299	315	323	341	334	320	338	333	333
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	32	32	31	34	36	40	43	49	54	54
Impulse control disorders, NEC	N.D.	34	37	34	36	44	50	47	55	54	53
Miscellaneous disorders	382	437	458	487	500	520	518	477	511	494	465
Mood disorders	1,255	1,402	1,491	1,578	1,653	1,732	1,787	1,631	1,772	1,749	1,668
Personality disorders	257	286	304	318	323	317	323	303	312	303	287
Schizophrenia and other psychotic disorders	477	513	518	523	509	504	487	451	470	440	400
Screening and history of mental health and substance abuse	1,318	1,501	1,624	1,755	1,877	1,946	1,996	1,821	2,010	1,998	1,899
Substance-related disorders	315	373	423	460	513	565	603	556	626	614	579
Suicide and intentional self-inflicted injury	61	72	76	83	85	92	106	105	121	116	115



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Casey County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,052	1,177	1,231	1,347	1,415	1,440	1,409	1,337	1,353	1,311	1,234
Female	518	565	582	650	672	679	663	628	629	599	561
Male	534	612	649	697	743	761	746	709	724	712	673
White	1,005	1,121	1,164	1,277	1,344	1,365	1,326	1,251	1,263	1,226	1,148
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	59%	61%	63%	63%	63%	62%	59%	58%	54%	48%
Female	59%	61%	61%	63%	63%	62%	59%	56%	55%	51%	45%
Male	55%	58%	62%	63%	63%	64%	63%	61%	60%	57%	51%
White	57%	59%	61%	63%	63%	62%	61%	58%	57%	53%	47%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	207	226	219	229	235	233	219	204	207	199	173
Alcohol-related disorders	44	41	43	46	38	34	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	460	511	525	583	598	613	598	575	578	552	514
Attention-deficit, conduct, and disruptive behavior disorders	404	458	486	509	530	551	548	516	519	493	457
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	274	315	362	425	460	484	507	482	507	502	489
Disorders usually diagnosed in infancy, childhood, adolescence	73	85	91	96	102	106	114	107	113	103	108
Impulse control disorders, NEC	40	41	41	42	38	38	39	35	N.D.	N.D.	N.D.
Miscellaneous disorders	230	249	252	265	268	268	247	225	218	210	188
Mood disorders	398	409	400	411	384	389	355	309	299	265	234
Personality disorders	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	31	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	495	533	551	584	577	580	547	488	471	452	405
Substance-related disorders	124	122	122	118	106	99	86	73	61	47	36
Suicide and intentional self-inflicted injury	N.D.	N.D.	32	N.D.	N.D.	32	33	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Casey County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,518	1,595	1,669	1,676	1,746	1,759	1,812	1,702	1,770	1,823	1,745
Female	1,030	1,073	1,137	1,124	1,179	1,180	1,229	1,156	1,197	1,231	1,172
Male	488	522	532	552	567	579	583	546	573	592	573
White	1,377	1,451	1,522	1,527	1,602	1,623	1,677	1,574	1,644	1,698	1,618
Minority	50	44	45	41	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	70%	72%	73%	72%	73%	73%	74%	70%	71%	72%	68%
Female	72%	74%	76%	74%	76%	75%	77%	72%	74%	76%	71%
Male	66%	67%	68%	69%	68%	69%	69%	65%	65%	65%	63%
White	69%	71%	73%	72%	73%	73%	74%	69%	70%	72%	67%
Minority	27%	23%	24%	21%	17%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	177	195	221	232	262	271	277	245	277	290	276
Alcohol-related disorders	140	145	153	154	160	159	170	158	164	158	158
Anxiety disorders	965	1,031	1,100	1,135	1,220	1,260	1,323	1,238	1,280	1,318	1,286
Attention-deficit, conduct, and disruptive behavior disorders	63	70	73	77	90	92	100	89	107	128	124
Delirium, dementia, and amnestic and other cognitive disorders	384	393	385	369	355	338	328	311	293	275	247
Developmental disorders	149	157	152	151	161	163	164	163	175	187	174
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	34	33
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	420	457	495	507	542	556	586	541	566	583	554
Mood disorders	934	1,000	1,054	1,074	1,132	1,157	1,220	1,134	1,168	1,200	1,148
Personality disorders	121	126	130	136	144	139	147	123	127	131	118
Schizophrenia and other psychotic disorders	406	427	431	423	429	411	409	386	383	371	340
Screening and history of mental health and substance abuse	691	746	802	829	908	887	923	863	925	954	905
Substance-related disorders	197	200	206	216	247	261	285	273	316	335	321
Suicide and intentional self-inflicted injury	48	52	51	47	51	58	64	68	80	82	78



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Christian County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,516	3,887	4,100	4,374	4,595	4,821	4,804	4,329	4,419	4,550	4,339
Female	1,697	1,885	1,968	2,096	2,200	2,278	2,256	2,008	2,055	2,077	1,968
Male	1,819	2,002	2,132	2,278	2,395	2,543	2,548	2,321	2,364	2,473	2,371
White	1,652	1,877	1,990	2,160	2,298	2,425	2,415	2,182	2,235	2,305	2,212
Minority	1,670	1,800	1,866	1,937	1,995	2,077	2,037	1,802	1,832	1,878	1,780

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	56%	58%	58%	58%	60%	59%	54%	55%	54%	49%
Female	55%	56%	57%	57%	57%	58%	58%	52%	53%	52%	47%
Male	56%	57%	58%	59%	59%	62%	61%	55%	56%	55%	50%
White	58%	58%	60%	60%	59%	61%	59%	54%	54%	53%	47%
Minority	48%	49%	49%	50%	50%	51%	51%	45%	46%	46%	42%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	464	529	555	599	610	622	608	529	537	528	494
Alcohol-related disorders	90	104	97	91	84	84	66	50	45	51	38
Anxiety disorders	521	553	593	628	632	651	623	572	554	515	453
Attention-deficit, conduct, and disruptive behavior disorders	1,810	2,020	2,181	2,315	2,363	2,420	2,375	2,136	2,122	2,146	2,024
Delirium, dementia, and amnesic and other cognitive disorders	45	50	42	48	50	51	49	46	45	49	43
Developmental disorders	645	706	769	866	904	950	978	907	944	992	941
Disorders usually diagnosed in infancy, childhood, adolescence	105	136	155	172	179	187	186	182	188	197	189
Impulse control disorders, NEC	456	477	463	454	428	395	363	307	287	265	224
Miscellaneous disorders	1,077	1,152	1,151	1,162	1,151	1,114	1,052	905	853	818	742
Mood disorders	901	971	979	977	933	893	831	719	671	628	524
Personality disorders	41	40	42	38	35	34	34	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	84	86	90	93	99	83	77	72	66	65	52
Screening and history of mental health and substance abuse	2,116	2,328	2,448	2,572	2,778	2,954	2,984	2,694	2,723	2,720	2,573
Substance-related disorders	305	328	321	300	278	254	220	172	165	162	120
Suicide and intentional self-inflicted injury	51	61	58	60	69	58	58	49	41	40	32



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Christian County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,044	3,265	3,378	3,494	3,612	3,725	3,886	3,605	3,906	4,031	3,767
Female	2,278	2,416	2,499	2,567	2,643	2,723	2,815	2,561	2,805	2,905	2,682
Male	766	849	879	927	969	1,002	1,071	1,044	1,101	1,126	1,085
White	1,841	1,951	2,011	2,088	2,153	2,226	2,317	2,122	2,309	2,403	2,216
Minority	1,082	1,183	1,228	1,260	1,292	1,322	1,374	1,284	1,394	1,434	1,353

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	61%	63%	64%	66%	67%	66%	68%	62%	66%	65%	60%
Female	62%	64%	65%	67%	68%	67%	69%	62%	67%	66%	60%
Male	57%	60%	61%	63%	64%	64%	67%	62%	64%	63%	59%
White	65%	67%	68%	70%	69%	68%	69%	63%	67%	67%	60%
Minority	50%	53%	54%	55%	56%	56%	58%	53%	56%	56%	51%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	287	327	346	374	391	427	449	419	462	482	433
Alcohol-related disorders	244	272	296	299	320	314	344	319	345	353	327
Anxiety disorders	1,219	1,333	1,408	1,464	1,522	1,561	1,676	1,586	1,670	1,660	1,606
Attention-deficit, conduct, and disruptive behavior disorders	190	206	222	265	301	326	357	377	421	484	480
Delirium, dementia, and amnesic and other cognitive disorders	935	954	957	948	910	877	861	792	811	758	689
Developmental disorders	455	460	473	468	462	453	469	465	465	460	459
Disorders usually diagnosed in infancy, childhood, adolescence	36	37	36	42	41	44	53	54	58	58	55
Impulse control disorders, NEC	69	80	90	100	113	130	153	157	168	174	180
Miscellaneous disorders	624	680	699	718	756	772	825	755	818	850	824
Mood disorders	1,413	1,581	1,707	1,799	1,905	2,006	2,126	2,034	2,210	2,248	2,120
Personality disorders	133	141	137	146	160	154	161	156	168	173	159
Schizophrenia and other psychotic disorders	864	889	879	883	853	837	836	800	816	822	786
Screening and history of mental health and substance abuse	1,550	1,720	1,852	1,981	2,080	2,194	2,340	2,141	2,373	2,501	2,340
Substance-related disorders	315	361	404	457	480	514	585	531	596	632	583
Suicide and intentional self-inflicted injury	81	90	88	110	112	117	132	136	167	185	165



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Clark County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,719	1,879	1,958	2,068	2,204	2,233	2,318	2,227	2,402	2,376	2,265
Female	849	922	946	973	1,033	1,031	1,033	980	1,057	1,050	981
Male	870	957	1,012	1,095	1,171	1,202	1,285	1,247	1,345	1,326	1,284
White	1,531	1,646	1,706	1,806	1,919	1,934	1,997	1,923	2,080	2,063	1,968
Minority	118	149	152	158	168	175	189	170	173	167	159

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	59%	61%	61%	61%	61%	61%	57%	58%	55%	52%
Female	57%	59%	61%	58%	60%	59%	58%	54%	55%	52%	47%
Male	56%	59%	61%	63%	62%	62%	63%	60%	60%	59%	56%
White	58%	59%	62%	61%	61%	61%	60%	56%	57%	55%	51%
Minority	31%	37%	36%	37%	37%	36%	37%	34%	35%	32%	31%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	471	531	539	564	575	571	569	501	506	476	430
Alcohol-related disorders	93	95	97	93	86	72	71	63	63	55	52
Anxiety disorders	789	857	916	986	1,041	1,040	1,099	1,065	1,140	1,162	1,116
Attention-deficit, conduct, and disruptive behavior disorders	739	798	846	890	940	936	942	862	880	853	792
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	33	33	32
Developmental disorders	259	309	355	403	472	525	608	621	708	730	707
Disorders usually diagnosed in infancy, childhood, adolescence	195	233	271	295	331	349	380	365	365	365	353
Impulse control disorders, NEC	59	58	61	58	55	56	51	43	36	37	N.D.
Miscellaneous disorders	229	245	251	257	252	249	257	247	257	248	233
Mood disorders	609	610	606	594	594	563	520	450	444	402	356
Personality disorders	45	43	46	45	41	40	34	32	34	31	N.D.
Schizophrenia and other psychotic disorders	90	85	88	87	87	83	82	74	69	66	59
Screening and history of mental health and substance abuse	999	1,034	1,043	1,025	1,036	1,005	981	873	880	811	747
Substance-related disorders	197	191	183	177	165	148	138	116	118	104	94
Suicide and intentional self-inflicted injury	70	72	71	64	60	63	66	60	65	63	55



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Clark County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,730	1,930	2,020	2,140	2,208	2,248	2,444	2,257	2,484	2,543	2,393
Female	1,279	1,421	1,493	1,605	1,664	1,682	1,825	1,656	1,827	1,852	1,721
Male	451	509	527	535	544	566	619	601	657	691	672
White	1,490	1,660	1,742	1,863	1,919	1,951	2,134	1,980	2,181	2,228	2,088
Minority	175	192	190	187	194	197	200	173	201	205	183

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	69%	69%	71%	71%	71%	73%	66%	72%	70%	66%
Female	67%	72%	72%	74%	74%	73%	76%	68%	74%	72%	67%
Male	59%	62%	62%	62%	63%	64%	66%	62%	66%	65%	62%
White	65%	69%	69%	71%	71%	71%	74%	66%	72%	70%	66%
Minority	44%	48%	47%	47%	46%	45%	46%	40%	47%	44%	38%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	224	253	275	317	334	348	379	361	399	415	400
Alcohol-related disorders	174	187	199	207	213	229	247	221	240	239	225
Anxiety disorders	941	1,066	1,139	1,236	1,272	1,330	1,435	1,364	1,491	1,533	1,477
Attention-deficit, conduct, and disruptive behavior disorders	60	80	82	99	120	126	165	175	213	232	245
Delirium, dementia, and amnesic and other cognitive disorders	477	482	458	447	441	423	419	372	357	333	305
Developmental disorders	141	158	157	167	178	176	191	174	191	201	207
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	32	34	37	44	46	48	53	65	68	62
Impulse control disorders, NEC	N.D.	31	31	32	34	35	43	44	50	51	59
Miscellaneous disorders	411	473	510	533	573	613	663	601	646	651	603
Mood disorders	1,007	1,161	1,234	1,345	1,385	1,426	1,567	1,467	1,608	1,657	1,586
Personality disorders	149	153	163	172	179	192	215	205	217	224	216
Schizophrenia and other psychotic disorders	391	415	435	438	444	457	493	446	452	438	421
Screening and history of mental health and substance abuse	999	1,158	1,253	1,371	1,448	1,512	1,685	1,561	1,722	1,761	1,641
Substance-related disorders	299	348	369	405	446	471	519	475	537	561	540
Suicide and intentional self-inflicted injury	55	66	79	81	95	96	111	101	113	117	126



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Clay County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,547	2,649	2,686	2,717	2,691	2,656	2,610	2,462	2,491	2,402	2,158
Female	1,259	1,291	1,286	1,286	1,276	1,260	1,247	1,153	1,169	1,108	991
Male	1,288	1,358	1,400	1,431	1,415	1,396	1,363	1,309	1,322	1,294	1,167
White	2,298	2,372	2,382	2,401	2,374	2,337	2,274	2,135	2,151	2,060	1,847
Minority	N.D.	32	34	37	34	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	60%	62%	61%	61%	60%	60%	57%	59%	57%	51%
Female	61%	63%	64%	63%	61%	60%	61%	56%	57%	54%	49%
Male	56%	57%	60%	60%	61%	61%	60%	59%	60%	59%	53%
White	57%	58%	60%	59%	59%	58%	58%	55%	56%	54%	48%
Minority	N.D.	10%	9%	10%	9%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	288	294	296	300	290	267	259	228	220	193	155
Alcohol-related disorders	166	164	161	151	156	141	131	115	98	88	69
Anxiety disorders	1,159	1,190	1,184	1,173	1,157	1,118	1,089	1,006	977	922	804
Attention-deficit, conduct, and disruptive behavior disorders	725	795	850	877	885	890	891	853	855	823	732
Delirium, dementia, and amnestic and other cognitive disorders	42	38	42	40	38	37	38	36	N.D.	31	N.D.
Developmental disorders	914	986	1,040	1,101	1,119	1,124	1,131	1,084	1,099	1,066	964
Disorders usually diagnosed in infancy, childhood, adolescence	151	164	184	193	208	205	199	188	191	176	160
Impulse control disorders, NEC	59	60	61	56	54	41	40	38	36	N.D.	N.D.
Miscellaneous disorders	626	622	623	603	578	545	522	476	459	421	361
Mood disorders	806	795	752	708	665	581	533	453	411	360	286
Personality disorders	140	138	136	133	133	121	115	110	102	94	68
Schizophrenia and other psychotic disorders	82	84	85	81	80	72	67	65	61	54	43
Screening and history of mental health and substance abuse	1,221	1,217	1,194	1,152	1,092	1,020	971	875	868	812	701
Substance-related disorders	449	427	401	370	327	287	263	228	211	180	134
Suicide and intentional self-inflicted injury	59	60	48	50	43	43	40	35	32	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Clay County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,889	4,070	4,235	4,367	4,467	4,520	4,634	4,426	4,605	4,607	4,565
Female	2,378	2,486	2,585	2,662	2,708	2,726	2,787	2,656	2,749	2,757	2,706
Male	1,511	1,584	1,650	1,705	1,759	1,794	1,847	1,770	1,856	1,850	1,859
White	3,531	3,712	3,870	4,006	4,103	4,156	4,270	4,071	4,253	4,247	4,185
Minority	102	100	95	84	73	68	58	54	55	63	58

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	74%	75%	77%	77%	77%	77%	78%	74%	77%	76%	74%
Female	77%	78%	81%	82%	81%	81%	81%	77%	80%	80%	77%
Male	69%	70%	71%	71%	72%	72%	74%	70%	72%	71%	71%
White	73%	75%	76%	76%	76%	76%	77%	74%	76%	76%	74%
Minority	23%	22%	22%	20%	17%	16%	14%	13%	13%	13%	11%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	383	399	423	450	460	454	495	459	481	500	518
Alcohol-related disorders	434	458	485	520	525	535	541	515	534	529	531
Anxiety disorders	3,010	3,139	3,255	3,412	3,515	3,548	3,676	3,527	3,635	3,645	3,644
Attention-deficit, conduct, and disruptive behavior disorders	160	174	189	208	228	252	264	268	290	339	358
Delirium, dementia, and amnesic and other cognitive disorders	664	665	663	647	630	614	580	526	504	472	424
Developmental disorders	290	295	305	311	326	344	357	349	379	402	440
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	34	39	57	69
Impulse control disorders, NEC	56	63	64	69	76	83	74	69	74	84	86
Miscellaneous disorders	1,566	1,636	1,731	1,839	1,926	1,928	1,991	1,903	1,957	1,966	1,932
Mood disorders	2,480	2,600	2,699	2,841	2,934	3,002	3,077	2,994	3,085	3,092	3,074
Personality disorders	280	295	309	334	358	366	351	332	349	350	367
Schizophrenia and other psychotic disorders	657	673	687	684	667	667	640	604	590	569	548
Screening and history of mental health and substance abuse	2,307	2,466	2,600	2,718	2,794	2,872	2,978	2,846	2,966	2,955	2,897
Substance-related disorders	1,301	1,359	1,459	1,582	1,682	1,724	1,802	1,750	1,791	1,812	1,813
Suicide and intentional self-inflicted injury	122	130	145	147	157	167	184	186	193	193	188



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Clinton County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	852	920	982	1,063	1,084	1,096	1,117	1,080	1,068	1,013	950
Female	419	459	491	527	530	527	537	532	523	497	461
Male	433	461	491	536	554	569	580	548	545	516	489
White	806	862	921	995	1,018	1,032	1,046	1,011	994	948	885
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	54%	58%	62%	63%	63%	62%	59%	59%	57%	53%
Female	54%	56%	60%	62%	64%	63%	62%	60%	58%	56%	52%
Male	51%	53%	56%	61%	62%	62%	62%	59%	59%	58%	55%
White	52%	54%	57%	61%	62%	62%	61%	59%	58%	57%	53%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	120	125	117	117	107	106	101	93	91	80	67
Alcohol-related disorders	37	40	41	42	37	39	39	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	397	424	456	489	512	528	537	543	519	489	465
Attention-deficit, conduct, and disruptive behavior disorders	310	329	338	357	370	377	379	350	330	299	278
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	137	156	185	216	243	263	291	290	291	289	281
Disorders usually diagnosed in infancy, childhood, adolescence	57	55	65	71	80	80	79	71	71	71	68
Impulse control disorders, NEC	58	59	65	69	64	65	60	50	43	36	33
Miscellaneous disorders	110	113	118	113	105	99	89	84	81	70	63
Mood disorders	243	249	244	236	224	209	198	170	153	127	110
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	457	499	560	605	600	590	564	515	509	478	436
Substance-related disorders	105	107	107	96	86	74	65	44	38	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Clinton County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,283	1,346	1,356	1,431	1,446	1,447	1,484	1,388	1,414	1,395	1,296
Female	858	889	886	930	939	952	980	921	928	915	852
Male	425	457	470	501	507	495	504	467	486	480	444
White	1,204	1,256	1,279	1,349	1,359	1,371	1,401	1,304	1,326	1,307	1,206
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	68%	69%	69%	69%	70%	70%	64%	67%	68%	65%
Female	68%	70%	71%	72%	72%	73%	74%	68%	70%	72%	68%
Male	59%	63%	65%	65%	64%	65%	64%	58%	61%	63%	60%
White	65%	67%	69%	69%	69%	70%	70%	64%	66%	68%	64%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	84	93	97	95	99	100	99	99	101	102	96
Alcohol-related disorders	100	113	114	136	136	130	127	120	121	125	119
Anxiety disorders	781	819	847	905	930	953	987	932	941	954	909
Attention-deficit, conduct, and disruptive behavior disorders	38	44	44	47	53	54	56	54	70	73	64
Delirium, dementia, and amnestic and other cognitive disorders	282	294	295	282	267	247	237	227	211	195	173
Developmental disorders	90	89	98	98	99	98	102	91	91	89	79
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	265	282	294	326	330	347	374	344	350	344	310
Mood disorders	766	801	834	876	901	895	913	862	854	855	803
Personality disorders	44	46	50	50	52	47	44	43	37	35	N.D.
Schizophrenia and other psychotic disorders	282	300	300	294	276	266	268	254	237	227	203
Screening and history of mental health and substance abuse	536	581	599	657	674	684	694	658	664	654	597
Substance-related disorders	242	259	267	313	326	322	320	315	326	336	321
Suicide and intentional self-inflicted injury	42	41	47	55	55	54	61	56	57	55	54



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Crittenden County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	437	483	515	532	501	511	516	483	485	474	443
Female	241	266	279	277	260	262	265	241	238	224	208
Male	196	217	236	255	241	249	251	242	247	250	235
White	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	57%	58%	59%	54%	55%	56%	51%	52%	50%	47%
Female	60%	60%	60%	60%	55%	56%	56%	51%	50%	48%	44%
Male	49%	54%	56%	59%	53%	55%	56%	52%	54%	53%	49%
White	54%	56%	58%	59%	53%	55%	55%	51%	52%	50%	46%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	55	58	72	70	71	72	71	67	64	61	55
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	140	142	160	162	154	153	149	135	131	120	104
Attention-deficit, conduct, and disruptive behavior disorders	185	207	225	227	224	237	244	237	225	213	203
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	86	95	110	126	118	122	136	133	142	148	141
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	38	39	37	33
Impulse control disorders, NEC	32	31	N.D.	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	43	48	49	47	46	47	42	36	41	42	37
Mood disorders	190	188	181	177	171	171	162	135	130	114	98
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	243	277	293	288	259	251	237	220	225	211	193
Substance-related disorders	34	34	34	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Crittenden County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	533	578	587	601	621	635	648	560	604	601	567
Female	377	406	421	426	431	450	463	393	418	431	395
Male	156	172	166	175	190	185	185	167	186	170	172
White	N.D.	546	549	562	580	593	606	522	571	562	529
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	61%	65%	65%	65%	64%	67%	67%	59%	64%	65%	63%
Female	63%	67%	68%	68%	67%	70%	71%	62%	68%	70%	66%
Male	56%	61%	57%	59%	58%	59%	57%	52%	57%	56%	56%
White	61%	65%	64%	65%	64%	66%	66%	58%	65%	65%	62%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	N.D.	N.D.	32	31	32	36	39	31	33	37	33
Alcohol-related disorders	38	40	43	47	43	42	50	39	37	41	38
Anxiety disorders	301	327	340	373	382	401	417	367	395	400	371
Attention-deficit, conduct, and disruptive behavior disorders	51	53	59	64	63	62	63	57	69	65	60
Delirium, dementia, and amnestic and other cognitive disorders	132	136	129	123	122	114	109	98	99	90	85
Developmental disorders	45	46	47	45	46	47	46	40	40	36	34
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	86	96	99	108	115	124	126	107	117	130	122
Mood disorders	307	333	369	388	407	420	433	375	415	413	372
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	116	125	132	128	125	123	132	113	113	102	89
Screening and history of mental health and substance abuse	274	308	326	347	360	372	385	300	312	321	294
Substance-related disorders	83	85	92	104	107	106	111	102	107	119	111
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	36	38	41	38	41	47	45



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Cumberland County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	512	558	573	607	614	619	602	583	583	570	537
Female	240	274	260	276	290	298	281	271	274	261	240
Male	272	284	313	331	324	321	321	312	309	309	297
White	459	503	508	547	550	552	535	521	520	508	482
Minority	35	32	35	N.D.	32	31	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	59%	61%	62%	62%	59%	58%	55%	57%	54%	49%
Female	53%	58%	58%	59%	60%	58%	56%	52%	55%	50%	44%
Male	58%	59%	65%	65%	64%	59%	60%	58%	60%	57%	53%
White	55%	58%	60%	61%	60%	57%	56%	54%	56%	52%	47%
Minority	42%	37%	40%	N.D.	39%	36%	N.D.	N.D.	N.D.	N.D.	N.D.


CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	77	81	83	90	89	80	79	73	74	65	60
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	228	240	234	241	250	251	250	238	236	221	206
Attention-deficit, conduct, and disruptive behavior disorders	219	236	245	250	248	242	229	221	210	197	184
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	149	170	198	210	222	228	234	226	231	238	224
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	31	34	35	36	37	37	35	32
Impulse control disorders, NEC	N.D.	N.D.	34	36	36	36	36	31	31	N.D.	N.D.
Miscellaneous disorders	59	67	63	59	64	57	46	46	42	34	N.D.
Mood disorders	216	231	228	221	218	203	189	170	155	135	124
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	267	286	284	302	287	274	262	249	235	214	193
Substance-related disorders	45	44	45	44	42	36	32	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Cumberland County											
ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	846	886	894	904	876	889	893	823	831	829	802
Female	569	596	586	601	570	575	581	528	540	536	516
Male	277	290	308	303	306	314	312	295	291	293	286
White	728	773	783	790	766	787	791	730	746	751	721
Minority	78	71	70	72	68	57	50	42	40	36	38
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	70%	72%	73%	71%	72%	73%	67%	70%	68%	65%
Female	70%	73%	74%	75%	74%	74%	76%	68%	73%	70%	67%
Male	64%	65%	69%	69%	67%	68%	67%	66%	65%	64%	62%
White	67%	70%	72%	72%	70%	71%	72%	66%	69%	67%	64%
Minority	49%	47%	48%	48%	51%	44%	39%	35%	36%	33%	34%
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	71	67	77	82	77	79	85	78	77	81	76
Alcohol-related disorders	91	94	101	94	93	103	102	91	91	91	82
Anxiety disorders	471	510	531	551	543	561	587	554	553	556	545
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	31	34	41	40	51	62	57	62	60	67
Delirium, dementia, and amnestic and other cognitive disorders	359	362	340	328	308	284	262	234	207	196	172
Developmental disorders	78	82	81	91	87	93	95	93	86	89	91
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	173	193	195	206	202	207	218	196	197	192	190
Mood disorders	478	516	538	559	559	581	591	542	553	547	532
Personality disorders	55	60	64	66	62	60	60	54	51	44	46
Schizophrenia and other psychotic disorders	272	285	275	274	276	267	261	245	229	227	209
Screening and history of mental health and substance abuse	297	326	355	368	368	380	385	349	369	374	360
Substance-related disorders	123	133	140	144	133	148	157	135	135	132	136
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	35	33	36	38	31	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Davies County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,935	5,464	5,742	6,048	6,115	6,211	6,289	5,982	6,177	6,173	5,802
Female	2,436	2,664	2,735	2,862	2,856	2,920	2,883	2,712	2,788	2,752	2,561
Male	2,499	2,800	3,007	3,186	3,259	3,291	3,406	3,270	3,389	3,421	3,241
White	3,937	4,350	4,540	4,760	4,806	4,871	4,912	4,674	4,806	4,795	4,488
Minority	643	692	735	778	776	771	762	711	760	747	723

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	62%	63%	63%	62%	61%	61%	57%	56%	53%	48%
Female	61%	62%	62%	62%	60%	59%	58%	54%	52%	49%	44%
Male	60%	62%	64%	64%	64%	62%	64%	60%	60%	58%	52%
White	59%	61%	62%	61%	60%	59%	59%	55%	54%	52%	46%
Minority	43%	43%	43%	42%	41%	39%	38%	35%	35%	33%	31%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,085	1,182	1,225	1,279	1,263	1,277	1,275	1,160	1,127	1,078	1,004
Alcohol-related disorders	210	223	206	198	184	179	163	145	137	121	92
Anxiety disorders	1,493	1,600	1,628	1,745	1,732	1,757	1,768	1,664	1,685	1,608	1,469
Attention-deficit, conduct, and disruptive behavior disorders	2,423	2,679	2,832	3,010	3,068	3,092	3,109	2,914	2,922	2,827	2,614
Delirium, dementia, and amnestic and other cognitive disorders	70	72	75	77	78	77	73	67	70	67	59
Developmental disorders	874	1,045	1,219	1,371	1,513	1,668	1,837	1,900	2,022	2,094	2,053
Disorders usually diagnosed in infancy, childhood, adolescence	351	406	439	461	484	514	548	526	542	539	522
Impulse control disorders, NEC	499	543	564	580	583	579	564	510	487	456	390
Miscellaneous disorders	698	737	736	715	680	671	665	593	586	553	499
Mood disorders	2,114	2,245	2,284	2,297	2,237	2,181	2,113	1,913	1,886	1,787	1,574
Personality disorders	238	239	220	211	202	187	175	159	150	132	102
Schizophrenia and other psychotic disorders	273	282	281	281	273	261	246	236	228	200	167
Screening and history of mental health and substance abuse	3,425	3,725	3,833	3,813	3,675	3,571	3,442	3,140	3,132	3,074	2,755
Substance-related disorders	505	529	503	487	446	422	401	369	356	299	228
Suicide and intentional self-inflicted injury	146	157	141	133	131	124	127	107	95	85	70



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Davies County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,346	4,790	5,146	5,491	5,744	5,983	6,152	5,739	6,320	6,451	6,259
Female	3,240	3,584	3,841	4,102	4,278	4,415	4,490	4,167	4,529	4,561	4,409
Male	1,106	1,206	1,305	1,389	1,466	1,568	1,662	1,572	1,791	1,890	1,850
White	3,745	4,138	4,432	4,707	4,940	5,145	5,291	4,901	5,459	5,545	5,309
Minority	421	455	491	538	539	555	548	530	548	576	587

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	68%	70%	72%	73%	74%	73%	68%	72%	71%	67%
Female	67%	70%	72%	75%	75%	76%	75%	69%	74%	73%	68%
Male	60%	63%	64%	66%	68%	68%	69%	65%	69%	68%	64%
White	65%	69%	70%	72%	74%	74%	74%	68%	72%	71%	67%
Minority	44%	45%	48%	49%	47%	48%	46%	45%	46%	45%	41%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	610	727	808	860	885	883	932	862	927	946	935
Alcohol-related disorders	442	514	542	607	659	691	744	716	774	791	776
Anxiety disorders	2,242	2,512	2,740	2,983	3,146	3,364	3,532	3,421	3,732	3,776	3,700
Attention-deficit, conduct, and disruptive behavior disorders	421	472	546	634	722	796	890	919	1,057	1,138	1,177
Delirium, dementia, and amnesic and other cognitive disorders	944	953	986	978	962	961	923	871	876	847	791
Developmental disorders	623	658	691	721	747	754	777	766	792	827	832
Disorders usually diagnosed in infancy, childhood, adolescence	49	55	61	70	76	79	90	95	106	117	139
Impulse control disorders, NEC	117	141	154	164	178	191	215	204	248	283	292
Miscellaneous disorders	1,028	1,168	1,325	1,450	1,571	1,672	1,704	1,517	1,631	1,620	1,558
Mood disorders	2,594	2,913	3,168	3,489	3,708	3,892	4,091	3,920	4,234	4,286	4,171
Personality disorders	517	564	627	657	682	714	724	695	744	756	727
Schizophrenia and other psychotic disorders	1,213	1,298	1,348	1,403	1,416	1,457	1,515	1,439	1,469	1,446	1,395
Screening and history of mental health and substance abuse	2,522	2,944	3,233	3,590	3,877	4,097	4,258	3,956	4,445	4,537	4,401
Substance-related disorders	730	847	917	1,010	1,101	1,190	1,286	1,239	1,370	1,423	1,374
Suicide and intentional self-inflicted injury	215	248	285	306	332	365	399	403	467	480	474



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Edmonson County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	626	687	738	751	761	768	794	757	802	812	756
Female	309	345	376	379	396	387	394	381	406	390	358
Male	317	342	362	372	365	381	400	376	396	422	398
White	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	758	726	769	777	725
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	51%	53%	55%	54%	54%	54%	56%	53%	55%	52%	49%
Female	52%	54%	56%	54%	56%	55%	56%	56%	58%	53%	49%
Male	49%	52%	54%	54%	52%	52%	55%	50%	51%	52%	49%
White	50%	53%	55%	54%	54%	53%	55%	52%	54%	52%	49%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	100	113	126	126	123	114	116	106	100	103	92
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	182	212	245	265	262	249	260	249	257	260	225
Attention-deficit, conduct, and disruptive behavior disorders	253	278	307	309	313	314	328	316	320	331	307
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	151	163	181	207	217	231	261	257	273	273	264
Disorders usually diagnosed in infancy, childhood, adolescence	46	54	69	69	71	76	78	72	79	83	83
Impulse control disorders, NEC	53	59	67	65	58	53	52	45	45	44	38
Miscellaneous disorders	78	83	87	83	82	77	76	69	73	75	63
Mood disorders	237	255	261	238	223	208	216	197	192	189	158
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	415	430	451	437	417	409	404	388	398	391	360
Substance-related disorders	33	36	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Edmonson County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	810	853	885	923	895	946	964	857	921	916	878
Female	565	591	615	647	620	644	672	591	653	636	603
Male	245	262	270	276	275	302	292	266	268	280	275
White	750	789	822	862	834	885	902	803	864	862	830
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	69%	69%	71%	69%	72%	72%	64%	68%	67%	63%
Female	67%	71%	71%	75%	73%	75%	76%	67%	73%	71%	67%
Male	61%	64%	65%	63%	62%	65%	64%	59%	58%	58%	57%
White	64%	68%	69%	71%	68%	71%	71%	64%	67%	66%	63%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	52	59	61	59	58	63	65	60	67	77	74
Alcohol-related disorders	82	88	98	95	102	101	102	93	94	89	82
Anxiety disorders	470	492	543	571	575	599	618	553	599	593	575
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	31	36	34	34	41	45	60	70	69
Delirium, dementia, and amnesic and other cognitive disorders	212	211	204	206	193	194	190	167	166	159	137
Developmental disorders	34	34	40	43	42	41	46	48	52	62	58
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	141	139	151	156	155	163	173	162	196	186	180
Mood disorders	499	541	573	610	607	646	672	605	649	651	631
Personality disorders	41	43	45	50	49	52	56	53	57	56	52
Schizophrenia and other psychotic disorders	168	180	185	183	178	186	188	166	172	170	154
Screening and history of mental health and substance abuse	465	507	539	574	547	574	607	545	589	576	557
Substance-related disorders	83	90	95	95	101	108	121	104	116	110	103
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	32	34	38



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Elliott County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	640	685	677	738	772	772	750	665	701	696	627
Female	344	361	353	376	391	388	384	333	341	328	300
Male	296	324	324	362	381	384	366	332	360	368	327
White	N.D.	N.D.	645	704	739	737	710	629	663	658	591
Minority	N.D.	N.D.	N.D.	N.D.	#N/A	#N/A	#N/A	#N/A	#N/A	#N/A	#N/A

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	56%	57%	60%	62%	61%	61%	56%	59%	58%	53%
Female	60%	59%	60%	62%	63%	62%	62%	58%	59%	58%	54%
Male	52%	54%	54%	58%	60%	60%	59%	55%	59%	59%	52%
White	55%	56%	56%	59%	61%	60%	59%	56%	58%	57%	52%
Minority	N.D.	N.D.	N.D.	N.D.	#N/A	#N/A	#N/A	#N/A	#N/A	#N/A	#N/A

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	88	95	95	98	108	114	114	98	97	92	83
Alcohol-related disorders	39	42	39	37	35	34	33	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	236	242	232	251	258	259	253	229	233	216	191
Attention-deficit, conduct, and disruptive behavior disorders	199	217	215	233	244	251	260	224	236	238	203
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	121	137	149	169	185	194	191	172	195	202	186
Disorders usually diagnosed in infancy, childhood, adolescence	40	44	50	58	65	65	71	51	59	63	56
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	89	90	86	91	91	88	85	72	74	64	45
Mood disorders	221	223	214	209	213	198	186	145	138	122	97
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	506	540	533	570	590	576	554	480	495	471	421
Substance-related disorders	77	80	67	65	63	55	50	40	31	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Elliott County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	721	795	862	883	903	911	881	813	844	862	824
Female	479	530	577	590	592	610	596	546	564	573	553
Male	242	265	285	293	311	301	285	267	280	289	271
White	667	738	800	824	837	843	811	749	782	797	761
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	65%	66%	67%	68%	68%	72%	67%	70%	70%	67%
Female	68%	70%	72%	73%	73%	75%	77%	72%	74%	74%	71%
Male	56%	56%	57%	57%	60%	59%	63%	59%	63%	63%	60%
White	62%	64%	65%	67%	67%	67%	71%	66%	69%	69%	66%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	49	58	61	68	64	64	66	62	64	70	66
Alcohol-related disorders	70	82	90	99	106	105	100	98	95	88	85
Anxiety disorders	458	499	558	578	597	609	612	542	556	577	540
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	N.D.	34	37	36	44	39	42	47	53
Delirium, dementia, and amnesic and other cognitive disorders	182	188	185	178	169	169	158	145	133	124	119
Developmental disorders	72	74	77	79	81	83	75	70	73	75	79
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	161	188	198	202	210	214	210	190	189	200	192
Mood disorders	417	468	527	551	563	576	571	518	529	548	525
Personality disorders	101	104	107	117	122	119	113	106	104	109	97
Schizophrenia and other psychotic disorders	127	137	145	146	151	143	140	127	119	118	111
Screening and history of mental health and substance abuse	421	479	539	584	609	615	612	570	600	621	596
Substance-related disorders	105	121	144	169	177	191	193	176	184	189	185
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	32	36	33	36	37



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Estill County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,175	1,265	1,311	1,387	1,428	1,424	1,477	1,362	1,428	1,464	1,363
Female	591	602	630	650	674	659	689	635	668	684	641
Male	584	663	681	737	754	765	788	727	760	780	722
White	1,086	1,167	1,195	1,265	1,299	1,289	1,336	1,224	1,293	1,326	1,234
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	59%	62%	63%	64%	65%	64%	66%	61%	62%	60%	55%
Female	61%	62%	63%	63%	62%	61%	63%	58%	59%	57%	53%
Male	57%	62%	64%	66%	67%	67%	69%	64%	64%	62%	57%
White	58%	61%	62%	63%	63%	62%	64%	59%	60%	58%	53%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	268	296	320	339	334	319	333	297	306	299	261
Alcohol-related disorders	51	53	55	59	49	50	45	37	35	33	N.D.
Anxiety disorders	467	503	540	567	577	582	617	559	589	601	562
Attention-deficit, conduct, and disruptive behavior disorders	406	454	488	535	558	566	593	527	542	555	500
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	241	279	324	362	409	442	463	449	490	499	487
Disorders usually diagnosed in infancy, childhood, adolescence	84	97	110	118	130	129	142	135	142	142	131
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	195	205	241	255	265	274	285	262	265	271	266
Mood disorders	362	370	365	352	332	306	301	243	231	218	184
Personality disorders	33	33	36	33	39	37	35	37	37	N.D.	N.D.
Schizophrenia and other psychotic disorders	38	37	38	33	34	32	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	789	822	817	821	811	773	776	683	693	693	626
Substance-related disorders	145	151	142	137	124	116	110	89	95	93	80
Suicide and intentional self-inflicted injury	45	43	42	41	40	39	36	33	32	31	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Estill County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,558	1,653	1,778	1,850	1,850	1,929	1,953	1,870	2,034	2,114	2,058
Female	1,062	1,107	1,191	1,241	1,227	1,266	1,289	1,222	1,325	1,356	1,323
Male	496	546	587	609	623	663	664	648	709	758	735
White	1,442	1,545	1,666	1,734	1,741	1,822	1,840	1,766	1,914	1,988	1,922
Minority	33	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	67%	69%	71%	73%	71%	72%	74%	69%	72%	72%	69%
Female	71%	74%	76%	78%	75%	75%	78%	72%	75%	76%	72%
Male	59%	62%	64%	65%	66%	68%	68%	63%	66%	67%	63%
White	66%	69%	71%	73%	71%	72%	74%	68%	71%	72%	68%
Minority	22%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	211	229	246	267	283	294	304	310	329	338	332
Alcohol-related disorders	129	134	145	142	151	158	168	167	174	183	169
Anxiety disorders	989	1,049	1,141	1,205	1,218	1,264	1,309	1,289	1,365	1,420	1,393
Attention-deficit, conduct, and disruptive behavior disorders	38	44	52	60	66	75	79	92	105	124	146
Delirium, dementia, and amnestic and other cognitive disorders	285	290	302	290	268	270	259	237	240	223	205
Developmental disorders	117	124	128	131	132	139	140	135	138	144	152
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	426	453	491	526	533	564	589	562	582	606	580
Mood disorders	887	949	1,020	1,107	1,120	1,171	1,235	1,191	1,272	1,320	1,311
Personality disorders	97	105	114	117	120	121	125	132	122	125	122
Schizophrenia and other psychotic disorders	276	288	302	300	288	291	297	277	273	269	255
Screening and history of mental health and substance abuse	903	1,007	1,094	1,178	1,221	1,288	1,329	1,284	1,367	1,404	1,342
Substance-related disorders	303	327	340	387	399	430	464	460	500	529	511
Suicide and intentional self-inflicted injury	45	54	53	63	68	70	78	75	81	87	87



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Fayette County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	7,333	8,272	8,945	9,444	9,683	9,926	10,213	9,838	10,551	10,766	10,199
Female	3,699	4,094	4,356	4,473	4,537	4,602	4,667	4,370	4,705	4,810	4,484
Male	3,634	4,178	4,589	4,971	5,146	5,324	5,546	5,468	5,846	5,956	5,715
White	3,810	4,362	4,707	4,981	5,226	5,431	5,577	5,428	5,897	6,036	5,783
Minority	3,108	3,454	3,710	3,884	3,851	3,827	3,889	3,689	3,860	3,910	3,645

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	50%	51%	52%	52%	51%	50%	50%	46%	46%	43%	39%
Female	50%	50%	51%	50%	49%	48%	47%	42%	43%	40%	35%
Male	50%	52%	54%	54%	53%	52%	52%	49%	50%	47%	42%
White	51%	52%	53%	53%	51%	49%	47%	43%	43%	40%	36%
Minority	43%	44%	45%	45%	44%	44%	45%	42%	42%	40%	35%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,962	2,258	2,450	2,592	2,666	2,732	2,725	2,549	2,620	2,557	2,369
Alcohol-related disorders	342	384	374	355	352	360	344	282	276	250	194
Anxiety disorders	2,186	2,414	2,598	2,751	2,869	2,896	2,919	2,760	2,818	2,820	2,625
Attention-deficit, conduct, and disruptive behavior disorders	3,940	4,418	4,772	5,062	5,153	5,265	5,368	5,099	5,217	5,156	4,825
Delirium, dementia, and amnestic and other cognitive disorders	177	203	216	226	244	246	246	228	226	217	190
Developmental disorders	1,531	1,811	2,127	2,371	2,552	2,773	3,048	3,121	3,511	3,703	3,566
Disorders usually diagnosed in infancy, childhood, adolescence	775	887	1,006	1,088	1,165	1,243	1,290	1,249	1,310	1,305	1,251
Impulse control disorders, NEC	377	389	405	420	408	380	368	322	293	266	225
Miscellaneous disorders	1,126	1,215	1,264	1,268	1,286	1,284	1,264	1,175	1,246	1,196	1,080
Mood disorders	2,653	2,845	2,925	2,970	2,949	2,892	2,786	2,508	2,458	2,362	2,113
Personality disorders	258	267	263	256	254	231	223	198	173	165	137
Schizophrenia and other psychotic disorders	412	427	434	452	445	423	426	396	391	364	314
Screening and history of mental health and substance abuse	4,611	5,034	5,291	5,346	5,277	5,138	5,151	4,719	4,751	4,627	4,148
Substance-related disorders	995	1,062	1,070	1,010	976	945	895	763	744	666	553
Suicide and intentional self-inflicted injury	334	351	376	386	397	379	388	349	347	339	286



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Fayette County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	7,887	8,546	8,896	9,248	9,456	9,669	9,978	9,299	10,247	10,625	10,265
Female	5,923	6,409	6,645	6,882	7,044	7,176	7,359	6,813	7,573	7,787	7,412
Male	1,964	2,137	2,251	2,366	2,412	2,493	2,619	2,486	2,674	2,838	2,853
White	5,086	5,516	5,731	5,987	6,117	6,287	6,545	6,061	6,680	6,844	6,571
Minority	2,471	2,673	2,779	2,849	2,904	2,897	2,908	2,710	3,027	3,220	3,062

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	61%	64%	64%	65%	65%	65%	66%	61%	66%	64%	60%
Female	62%	65%	65%	64%	64%	64%	66%	61%	65%	64%	59%
Male	58%	61%	64%	66%	65%	66%	68%	63%	67%	65%	60%
White	64%	67%	68%	69%	69%	69%	70%	64%	68%	66%	61%
Minority	50%	51%	51%	51%	50%	50%	51%	48%	52%	52%	47%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	818	913	961	1,020	1,082	1,128	1,226	1,223	1,377	1,440	1,416
Alcohol-related disorders	974	1,078	1,163	1,240	1,274	1,278	1,372	1,279	1,409	1,456	1,421
Anxiety disorders	3,968	4,359	4,672	4,956	5,153	5,356	5,634	5,384	5,833	5,930	5,743
Attention-deficit, conduct, and disruptive behavior disorders	533	611	686	793	866	960	1,089	1,128	1,311	1,449	1,474
Delirium, dementia, and amnestic and other cognitive disorders	2,345	2,392	2,380	2,305	2,259	2,205	2,135	1,930	1,899	1,815	1,685
Developmental disorders	909	960	1,037	1,110	1,137	1,160	1,205	1,150	1,217	1,256	1,269
Disorders usually diagnosed in infancy, childhood, adolescence	112	129	139	159	177	177	211	216	251	280	300
Impulse control disorders, NEC	153	168	179	190	211	233	261	270	297	318	305
Miscellaneous disorders	2,003	2,262	2,378	2,518	2,657	2,802	2,936	2,721	3,036	3,138	2,978
Mood disorders	4,469	4,905	5,222	5,500	5,696	5,879	6,158	5,871	6,344	6,501	6,311
Personality disorders	806	864	937	966	960	994	1,056	998	1,062	1,072	1,041
Schizophrenia and other psychotic disorders	2,162	2,261	2,302	2,300	2,338	2,328	2,397	2,237	2,253	2,207	2,095
Screening and history of mental health and substance abuse	4,576	5,140	5,526	5,886	6,132	6,288	6,553	6,144	6,796	7,082	6,848
Substance-related disorders	1,643	1,825	1,944	2,155	2,295	2,371	2,580	2,415	2,741	2,915	2,795
Suicide and intentional self-inflicted injury	388	419	441	495	520	577	666	666	748	764	764



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Fleming County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	853	902	960	999	1,012	991	1,002	923	976	1,011	882
Female	430	462	479	494	494	472	485	429	452	471	411
Male	423	440	481	505	518	519	517	494	524	540	471
White	797	836	894	927	937	914	918	842	890	924	804
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	56%	58%	57%	58%	57%	56%	51%	52%	51%	44%
Female	59%	57%	60%	57%	57%	55%	53%	48%	50%	49%	43%
Male	55%	55%	57%	57%	60%	59%	58%	54%	55%	53%	46%
White	56%	55%	58%	57%	57%	56%	54%	50%	51%	50%	43%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	300	318	349	369	382	385	402	373	393	386	347
Alcohol-related disorders	37	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	285	301	320	342	343	327	327	295	311	303	262
Attention-deficit, conduct, and disruptive behavior disorders	333	353	383	402	418	419	416	353	378	355	317
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	233	258	297	323	346	352	368	338	367	372	335
Disorders usually diagnosed in infancy, childhood, adolescence	71	76	83	95	105	104	102	85	94	95	85
Impulse control disorders, NEC	N.D.	32	31	32	N.D.	31	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	128	121	120	123	124	124	119	105	109	107	90
Mood disorders	297	301	304	305	298	286	276	226	234	222	194
Personality disorders	95	93	90	94	92	91	80	62	61	56	50
Schizophrenia and other psychotic disorders	74	65	71	74	75	75	72	60	58	54	50
Screening and history of mental health and substance abuse	568	589	607	617	598	563	549	488	496	484	424
Substance-related disorders	81	82	81	84	85	79	72	56	50	46	34
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Fleming County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	918	990	1,050	1,094	1,128	1,117	1,156	1,101	1,193	1,188	1,151
Female	634	675	713	731	758	757	792	726	802	811	770
Male	284	315	337	363	370	360	364	375	391	377	381
White	835	900	949	993	1,028	1,019	1,060	1,003	1,089	1,086	1,045
Minority	N.D.	31	34	33	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	67%	68%	69%	67%	69%	64%	68%	65%	61%
Female	66%	70%	70%	69%	71%	70%	72%	64%	70%	68%	63%
Male	59%	60%	62%	65%	65%	63%	63%	63%	64%	60%	58%
White	64%	66%	67%	67%	68%	67%	68%	63%	68%	65%	61%
Minority	N.D.	25%	25%	23%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	80	98	100	108	119	125	136	129	137	145	147
Alcohol-related disorders	69	90	94	95	98	105	107	112	113	112	110
Anxiety disorders	490	544	590	630	635	664	708	667	711	704	688
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	36	43	50	62	69	66	76	78	97
Delirium, dementia, and amnestic and other cognitive disorders	228	220	228	216	214	203	203	185	179	162	155
Developmental disorders	92	104	108	112	115	121	128	119	117	122	125
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	210	222	240	259	277	289	307	294	301	300	289
Mood disorders	456	502	553	590	615	631	669	624	664	662	652
Personality disorders	79	92	109	113	123	139	141	134	130	130	129
Schizophrenia and other psychotic disorders	218	220	231	229	229	224	229	208	191	179	171
Screening and history of mental health and substance abuse	498	564	620	658	696	701	711	677	746	745	735
Substance-related disorders	118	144	157	170	177	191	201	197	216	218	213
Suicide and intentional self-inflicted injury	N.D.	N.D.	31	32	38	39	44	38	40	42	43



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Floyd County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,176	4,404	4,495	4,544	4,554	4,491	4,468	4,171	4,216	4,198	3,890
Female	2,136	2,206	2,212	2,220	2,191	2,133	2,097	1,915	1,945	1,945	1,795
Male	2,040	2,198	2,283	2,324	2,363	2,358	2,371	2,256	2,271	2,253	2,095
White	3,846	4,038	4,101	4,121	4,096	4,020	3,981	3,702	3,747	3,714	3,459
Minority	38	37	38	33	33	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	60%	61%	62%	62%	62%	62%	58%	59%	58%	53%
Female	60%	61%	61%	62%	62%	61%	61%	57%	57%	56%	51%
Male	55%	58%	60%	62%	62%	63%	64%	60%	61%	60%	55%
White	57%	58%	59%	60%	60%	60%	60%	56%	57%	56%	51%
Minority	9%	8%	8%	7%	6%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	678	747	758	781	779	760	745	637	602	585	520
Alcohol-related disorders	180	167	162	149	130	115	118	88	79	65	47
Anxiety disorders	1,504	1,549	1,537	1,569	1,612	1,584	1,572	1,469	1,468	1,441	1,313
Attention-deficit, conduct, and disruptive behavior disorders	1,348	1,516	1,637	1,698	1,724	1,713	1,720	1,549	1,541	1,537	1,417
Delirium, dementia, and amnestic and other cognitive disorders	58	54	57	55	49	46	43	45	45	44	42
Developmental disorders	797	915	1,017	1,103	1,178	1,227	1,299	1,295	1,366	1,396	1,320
Disorders usually diagnosed in infancy, childhood, adolescence	241	266	290	311	327	350	365	338	347	352	332
Impulse control disorders, NEC	101	107	105	98	99	88	78	67	59	52	41
Miscellaneous disorders	630	623	604	592	584	539	508	450	434	399	374
Mood disorders	1,224	1,220	1,143	1,102	1,033	943	893	746	692	649	553
Personality disorders	63	61	61	60	56	54	50	40	37	32	N.D.
Schizophrenia and other psychotic disorders	116	117	113	114	110	105	95	77	70	70	60
Screening and history of mental health and substance abuse	2,867	2,980	3,013	2,989	2,921	2,788	2,678	2,418	2,376	2,308	2,060
Substance-related disorders	506	488	440	415	378	322	288	236	222	204	186
Suicide and intentional self-inflicted injury	113	117	111	108	99	96	91	74	68	69	56



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Floyd County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,093	5,494	5,787	6,039	6,124	6,149	6,335	6,049	6,467	6,480	6,407
Female	3,351	3,613	3,787	3,909	3,967	4,012	4,159	3,939	4,233	4,205	4,150
Male	1,742	1,881	2,000	2,130	2,157	2,137	2,176	2,110	2,234	2,275	2,257
White	4,656	5,046	5,324	5,561	5,638	5,675	5,841	5,564	5,971	5,979	5,873
Minority	88	87	88	76	62	44	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	72%	74%	74%	74%	74%	75%	71%	75%	75%	72%
Female	73%	76%	78%	78%	78%	79%	80%	74%	79%	79%	76%
Male	61%	65%	66%	67%	67%	66%	67%	64%	68%	68%	66%
White	67%	71%	73%	73%	73%	73%	74%	70%	74%	74%	72%
Minority	16%	16%	16%	13%	11%	8%	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	544	571	618	655	690	727	750	739	798	808	813
Alcohol-related disorders	511	576	602	648	654	643	677	643	671	681	645
Anxiety disorders	3,521	3,795	4,023	4,221	4,337	4,388	4,534	4,353	4,585	4,561	4,470
Attention-deficit, conduct, and disruptive behavior disorders	112	128	140	160	182	209	248	259	290	350	361
Delirium, dementia, and amnesic and other cognitive disorders	686	712	699	695	693	661	662	608	591	569	516
Developmental disorders	283	299	296	323	329	335	363	359	364	373	381
Disorders usually diagnosed in infancy, childhood, adolescence	50	56	56	58	62	68	69	66	77	82	78
Impulse control disorders, NEC	42	47	50	60	64	72	79	83	88	94	93
Miscellaneous disorders	1,656	1,837	1,914	2,024	2,090	2,130	2,192	2,070	2,118	2,072	1,975
Mood disorders	3,124	3,395	3,565	3,766	3,849	3,919	4,067	3,925	4,108	4,153	4,061
Personality disorders	192	211	221	236	250	252	268	264	257	261	253
Schizophrenia and other psychotic disorders	762	809	828	837	842	835	834	778	777	752	720
Screening and history of mental health and substance abuse	3,462	3,812	4,074	4,297	4,368	4,458	4,661	4,497	4,804	4,858	4,842
Substance-related disorders	1,255	1,354	1,465	1,551	1,618	1,680	1,801	1,760	1,879	1,922	1,889
Suicide and intentional self-inflicted injury	166	187	205	227	231	239	272	269	276	283	280



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Franklin County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,654	1,886	1,981	2,063	2,132	2,123	2,112	2,025	2,178	2,249	2,108
Female	801	923	967	982	999	964	954	924	979	992	927
Male	853	963	1,014	1,081	1,133	1,159	1,158	1,101	1,199	1,257	1,181
White	1,309	1,506	1,559	1,612	1,668	1,650	1,617	1,553	1,673	1,754	1,643
Minority	244	259	285	296	300	297	301	283	298	294	278

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	57%	60%	58%	58%	54%	53%	48%	49%	47%	42%
Female	55%	57%	59%	57%	55%	51%	50%	45%	46%	43%	38%
Male	54%	57%	60%	59%	60%	57%	55%	51%	52%	50%	45%
White	55%	58%	61%	59%	58%	54%	52%	48%	49%	47%	41%
Minority	36%	35%	38%	36%	37%	34%	33%	30%	31%	28%	26%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	365	418	438	439	445	428	420	385	401	416	377
Alcohol-related disorders	70	71	69	64	55	55	46	40	39	38	N.D.
Anxiety disorders	504	564	568	586	586	551	543	511	557	572	509
Attention-deficit, conduct, and disruptive behavior disorders	900	1,025	1,087	1,128	1,165	1,174	1,166	1,080	1,139	1,138	1,076
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	299	362	428	481	500	545	601	604	676	710	682
Disorders usually diagnosed in infancy, childhood, adolescence	164	197	227	255	259	273	294	274	298	309	292
Impulse control disorders, NEC	114	126	125	120	124	116	115	89	95	89	78
Miscellaneous disorders	314	362	377	370	390	387	385	350	375	378	357
Mood disorders	655	723	742	748	738	709	667	613	627	612	531
Personality disorders	53	56	58	50	45	38	36	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	93	102	93	95	94	87	86	66	72	67	56
Screening and history of mental health and substance abuse	1,137	1,229	1,229	1,229	1,201	1,110	1,036	915	940	913	770
Substance-related disorders	192	202	197	170	157	140	132	115	124	118	88
Suicide and intentional self-inflicted injury	57	67	72	67	67	63	67	55	61	63	46



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Franklin County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,711	1,865	1,992	2,099	2,214	2,372	2,450	2,257	2,542	2,705	2,608
Female	1,194	1,288	1,385	1,475	1,561	1,679	1,727	1,584	1,818	1,894	1,787
Male	517	577	607	624	653	693	723	673	724	811	821
White	1,472	1,622	1,726	1,807	1,915	2,045	2,106	1,946	2,195	2,329	2,247
Minority	183	181	199	218	222	238	240	218	254	275	246

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	66%	67%	68%	69%	70%	72%	66%	71%	70%	65%
Female	64%	66%	68%	69%	70%	71%	73%	65%	72%	71%	65%
Male	62%	64%	65%	67%	68%	69%	71%	66%	68%	69%	65%
White	65%	68%	69%	71%	72%	72%	74%	67%	72%	72%	68%
Minority	43%	41%	43%	43%	43%	44%	45%	39%	46%	44%	36%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	178	190	221	241	268	282	291	272	310	334	331
Alcohol-related disorders	240	262	281	298	309	325	348	323	353	356	333
Anxiety disorders	873	971	1,086	1,149	1,236	1,305	1,401	1,334	1,490	1,555	1,517
Attention-deficit, conduct, and disruptive behavior disorders	119	144	176	185	217	259	275	293	340	389	397
Delirium, dementia, and amnesic and other cognitive disorders	373	396	392	388	385	397	396	366	370	382	340
Developmental disorders	237	262	285	312	329	347	380	374	389	423	419
Disorders usually diagnosed in infancy, childhood, adolescence	36	42	48	52	55	61	72	81	90	99	110
Impulse control disorders, NEC	68	80	81	93	109	122	134	142	143	150	151
Miscellaneous disorders	358	403	473	557	625	677	737	652	739	759	729
Mood disorders	991	1,102	1,221	1,308	1,394	1,486	1,561	1,488	1,642	1,695	1,653
Personality disorders	159	176	195	217	226	235	250	250	273	279	276
Schizophrenia and other psychotic disorders	323	366	379	396	417	441	471	448	459	462	432
Screening and history of mental health and substance abuse	1,071	1,191	1,309	1,406	1,517	1,612	1,698	1,580	1,822	1,929	1,884
Substance-related disorders	399	445	502	563	628	652	714	661	759	827	821
Suicide and intentional self-inflicted injury	89	95	102	107	114	128	137	138	153	159	173



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Fulton County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	585	612	638	676	680	646	668	582	580	537	465
Female	284	299	297	315	318	283	309	258	251	238	199
Male	301	313	341	361	362	363	359	324	329	299	266
White	299	294	303	329	330	308	330	287	276	260	235
Minority	250	274	283	292	292	279	275	230	233	212	170

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	53%	56%	57%	57%	55%	55%	50%	50%	49%	42%
Female	51%	50%	51%	56%	56%	53%	54%	46%	46%	45%	38%
Male	53%	56%	60%	59%	58%	57%	57%	54%	54%	52%	46%
White	53%	51%	55%	55%	55%	52%	53%	50%	48%	47%	42%
Minority	44%	48%	48%	50%	49%	48%	47%	39%	40%	39%	32%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	38	37	44	37	37	32	32	N.D.	35	36	34
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	97	99	91	95	95	95	93	75	70	73	58
Attention-deficit, conduct, and disruptive behavior disorders	249	261	267	279	283	285	299	267	270	253	218
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	153	181	211	223	255	265	276	270	277	257	237
Disorders usually diagnosed in infancy, childhood, adolescence	53	52	59	58	58	58	59	60	63	61	55
Impulse control disorders, NEC	N.D.	32	33	33	33	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	47	49	46	56	62	55	53	51	49	46	41
Mood disorders	151	146	133	133	137	128	133	111	109	93	78
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	356	371	371	407	381	335	327	269	272	232	180
Substance-related disorders	35	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Fulton County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	630	692	749	772	787	764	802	742	787	782	750
Female	489	537	574	595	598	587	617	555	587	569	550
Male	141	155	175	177	189	177	185	187	200	213	200
White	347	378	417	441	443	435	473	442	473	467	460
Minority	250	277	296	296	305	286	285	253	271	266	247

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	59%	62%	61%	63%	61%	62%	58%	62%	63%	60%
Female	60%	63%	66%	65%	66%	64%	66%	61%	66%	66%	64%
Male	45%	49%	54%	52%	54%	51%	52%	50%	53%	55%	51%
White	54%	57%	61%	61%	62%	60%	62%	59%	64%	64%	62%
Minority	51%	55%	58%	56%	56%	54%	54%	47%	51%	51%	49%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	N.D.	32	32	35	35	40	41	44	45	45	48
Alcohol-related disorders	74	74	83	77	82	80	79	76	81	69	68
Anxiety disorders	345	376	397	411	413	421	453	419	436	434	416
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	N.D.	31	39	N.D.	39	44	47	56	62
Delirium, dementia, and amnesic and other cognitive disorders	107	122	117	107	105	103	109	104	113	106	97
Developmental disorders	N.D.	32	35	34	33	36	41	50	51	47	42
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	121	139	152	149	161	165	172	166	173	161	153
Mood disorders	349	381	412	433	459	444	469	427	470	459	452
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33	32	N.D.
Schizophrenia and other psychotic disorders	90	94	95	94	97	99	104	96	101	106	96
Screening and history of mental health and substance abuse	327	374	434	454	456	439	451	423	438	444	423
Substance-related disorders	86	100	112	103	113	112	114	110	120	113	100
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	32	35	38	33



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Gallatin County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	399	466	506	527	551	564	593	565	617	633	592
Female	189	204	229	232	251	250	266	260	290	293	275
Male	210	262	277	295	300	314	327	305	327	340	317
White	363	424	461	473	495	502	534	511	563	578	539
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	54%	54%	53%	54%	53%	54%	49%	51%	48%	44%
Female	50%	50%	52%	49%	53%	50%	50%	47%	50%	45%	41%
Male	53%	57%	55%	57%	55%	57%	57%	51%	52%	51%	47%
White	51%	53%	53%	53%	54%	52%	52%	48%	50%	47%	42%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	80	92	94	92	94	92	89	82	89	92	72
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	164	181	191	192	194	185	175	165	167	163	141
Attention-deficit, conduct, and disruptive behavior disorders	205	235	262	278	308	323	337	335	344	355	331
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	55	72	80	90	102	120	131	126	146	142	141
Disorders usually diagnosed in infancy, childhood, adolescence	40	46	49	49	60	56	58	56	61	63	58
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	60	63	68	64	65	53	60	57	58	59	52
Mood disorders	172	180	187	175	181	170	162	141	144	134	126
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	228	264	277	279	269	276	283	255	264	255	239
Substance-related disorders	42	41	44	41	42	37	37	32	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Gallatin County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	485	515	537	516	530	559	600	560	599	607	604
Female	334	352	378	364	363	386	420	383	409	415	414
Male	151	163	159	152	167	173	180	177	190	192	190
White	442	474	491	472	488	516	553	515	552	560	558
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	67%	69%	71%	68%	68%	71%	73%	67%	71%	69%	65%
Female	68%	69%	73%	68%	67%	72%	73%	67%	72%	71%	67%
Male	65%	68%	66%	67%	70%	69%	73%	68%	69%	65%	61%
White	68%	71%	71%	69%	69%	72%	73%	68%	71%	69%	65%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	59	58	59	60	52	61	71	71	69	75	72
Alcohol-related disorders	48	51	53	51	55	61	69	67	68	62	62
Anxiety disorders	272	291	310	312	323	367	403	383	405	397	389
Attention-deficit, conduct, and disruptive behavior disorders	48	55	60	65	67	68	84	88	97	102	97
Delirium, dementia, and amnestic and other cognitive disorders	192	197	194	177	176	170	177	163	147	132	123
Developmental disorders	53	51	58	55	61	57	59	62	64	64	60
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	83	97	110	119	124	136	152	132	136	137	129
Mood disorders	285	303	328	315	319	355	401	389	409	395	391
Personality disorders	33	35	40	45	44	51	57	59	53	53	51
Schizophrenia and other psychotic disorders	147	151	158	142	130	137	148	142	137	136	126
Screening and history of mental health and substance abuse	248	264	308	325	329	359	386	371	404	406	392
Substance-related disorders	79	87	95	94	94	110	130	133	143	145	139
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	35	35	37	36	34



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Garrard County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	744	828	854	929	949	961	980	955	1,010	1,005	955
Female	368	402	408	441	458	468	469	454	488	477	456
Male	376	426	446	488	491	493	511	501	522	528	499
White	670	738	770	840	857	859	878	859	903	902	854
Minority	40	51	40	39	39	49	41	41	44	37	38

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	56%	57%	59%	58%	59%	57%	54%	55%	50%	46%
Female	55%	56%	56%	57%	58%	59%	56%	52%	54%	48%	45%
Male	52%	55%	58%	61%	59%	58%	58%	56%	56%	52%	47%
White	53%	55%	57%	58%	58%	58%	56%	54%	53%	49%	44%
Minority	33%	37%	29%	28%	28%	31%	26%	25%	28%	21%	23%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	229	240	263	293	291	280	282	265	279	266	248
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	291	323	348	386	404	412	406	403	408	404	385
Attention-deficit, conduct, and disruptive behavior disorders	333	369	391	417	423	413	431	403	411	395	372
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	150	173	207	231	242	259	267	273	294	291	280
Disorders usually diagnosed in infancy, childhood, adolescence	75	77	84	98	104	108	106	112	121	126	121
Impulse control disorders, NEC	37	35	36	34	37	35	35	31	31	N.D.	N.D.
Miscellaneous disorders	142	159	162	164	179	174	177	153	153	143	127
Mood disorders	260	267	271	272	267	250	248	221	212	201	182
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	34	32	33	34	37	36	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	419	436	431	440	427	421	421	391	400	389	370
Substance-related disorders	67	75	72	72	68	70	65	55	58	51	49
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	31	33	32	32	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Garrard County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	814	898	933	948	968	1,012	1,082	990	1,072	1,097	1,044
Female	581	651	653	680	695	710	763	679	745	753	707
Male	233	247	280	268	273	302	319	311	327	344	337
White	724	795	836	851	871	909	962	881	957	977	914
Minority	59	68	61	58	55	58	64	57	61	60	61

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	66%	68%	68%	69%	68%	72%	65%	69%	68%	62%
Female	67%	69%	72%	72%	72%	72%	76%	68%	73%	71%	64%
Male	58%	57%	62%	60%	61%	61%	64%	61%	61%	62%	57%
White	64%	65%	68%	68%	69%	68%	72%	64%	68%	67%	61%
Minority	42%	45%	46%	41%	39%	39%	41%	38%	40%	36%	33%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	122	147	166	173	191	204	218	204	215	214	203
Alcohol-related disorders	55	60	67	67	68	77	83	80	83	81	75
Anxiety disorders	466	533	567	579	610	650	694	640	691	703	671
Attention-deficit, conduct, and disruptive behavior disorders	59	70	74	75	90	98	106	101	111	115	108
Delirium, dementia, and amnesic and other cognitive disorders	221	222	229	213	199	196	197	183	176	174	159
Developmental disorders	79	85	90	91	91	94	95	97	98	101	100
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	221	251	270	293	300	313	333	300	314	319	297
Mood disorders	473	545	565	588	611	640	687	626	681	692	658
Personality disorders	86	95	106	114	119	118	128	110	116	120	114
Schizophrenia and other psychotic disorders	251	246	240	226	219	224	235	223	224	209	185
Screening and history of mental health and substance abuse	387	447	475	496	534	571	624	572	631	642	618
Substance-related disorders	130	140	161	163	181	206	234	217	225	239	226
Suicide and intentional self-inflicted injury	37	39	44	46	47	56	71	67	69	71	64



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Grant County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,225	1,488	1,623	1,796	1,827	1,903	1,997	1,887	1,985	2,024	1,901
Female	611	744	803	863	867	918	934	861	920	920	855
Male	614	744	820	933	960	985	1,063	1,026	1,065	1,104	1,046
White	1,165	1,405	1,527	1,679	1,703	1,767	1,836	1,739	1,818	1,850	1,744
Minority	N.D.	N.D.	N.D.	31	32	36	37	31	41	46	40

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	58%	59%	62%	61%	59%	60%	54%	55%	53%	48%
Female	56%	59%	59%	61%	60%	59%	58%	52%	53%	50%	45%
Male	55%	57%	59%	63%	62%	60%	62%	57%	57%	56%	51%
White	55%	57%	59%	61%	60%	58%	59%	53%	53%	52%	47%
Minority	N.D.	N.D.	N.D.	19%	18%	19%	19%	15%	19%	19%	17%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	207	258	285	308	302	289	302	270	279	262	243
Alcohol-related disorders	51	62	63	61	62	60	55	48	43	40	34
Anxiety disorders	527	632	674	751	754	759	759	685	688	651	609
Attention-deficit, conduct, and disruptive behavior disorders	607	749	826	933	976	1,055	1,119	1,068	1,115	1,133	1,085
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	32	31	37	38	N.D.
Developmental disorders	195	245	289	345	373	401	450	429	455	499	488
Disorders usually diagnosed in infancy, childhood, adolescence	121	146	159	189	192	208	215	204	209	201	195
Impulse control disorders, NEC	35	39	47	47	46	46	45	41	36	33	35
Miscellaneous disorders	201	249	252	269	266	269	266	232	229	213	193
Mood disorders	485	553	575	592	562	559	548	477	475	426	392
Personality disorders	73	78	72	68	56	55	50	47	43	39	34
Schizophrenia and other psychotic disorders	43	44	44	50	51	53	56	46	45	43	39
Screening and history of mental health and substance abuse	747	850	913	970	944	931	942	838	837	790	690
Substance-related disorders	126	141	156	147	143	133	128	111	113	95	75
Suicide and intentional self-inflicted injury	54	60	59	52	59	61	62	57	61	59	49



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Grant County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,085	1,210	1,330	1,447	1,469	1,551	1,704	1,583	1,784	1,858	1,765
Female	798	898	970	1,064	1,084	1,148	1,270	1,159	1,311	1,329	1,256
Male	287	312	360	383	385	403	434	424	473	529	509
White	1,028	1,156	1,272	1,386	1,405	1,482	1,628	1,509	1,711	1,779	1,675
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	69%	72%	73%	75%	76%	76%	78%	68%	74%	74%	68%
Female	72%	75%	75%	77%	78%	78%	80%	70%	76%	76%	70%
Male	62%	63%	69%	70%	71%	71%	71%	65%	68%	67%	62%
White	69%	72%	73%	75%	76%	76%	77%	68%	74%	74%	68%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	115	126	141	157	162	176	196	181	197	208	220
Alcohol-related disorders	100	115	142	149	142	147	158	153	165	168	162
Anxiety disorders	689	776	879	978	1,008	1,082	1,199	1,125	1,251	1,267	1,242
Attention-deficit, conduct, and disruptive behavior disorders	105	122	143	158	178	195	224	209	242	270	273
Delirium, dementia, and amnestic and other cognitive disorders	226	244	257	262	252	244	258	236	235	234	211
Developmental disorders	119	124	136	142	143	150	158	151	156	166	174
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	40
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	31	34	N.D.	37	38	42
Miscellaneous disorders	306	344	396	448	460	499	552	475	517	542	487
Mood disorders	713	799	893	992	1,015	1,083	1,200	1,114	1,238	1,292	1,242
Personality disorders	144	167	185	206	207	213	230	202	205	206	189
Schizophrenia and other psychotic disorders	242	269	282	291	289	298	316	287	301	310	304
Screening and history of mental health and substance abuse	684	783	889	998	1,036	1,107	1,229	1,133	1,290	1,350	1,281
Substance-related disorders	183	220	256	293	298	328	362	333	366	400	408
Suicide and intentional self-inflicted injury	62	70	85	100	102	106	121	105	118	136	139



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Graves County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,682	1,836	1,987	2,152	2,204	2,245	2,319	2,144	2,267	2,238	2,118
Female	784	851	920	985	988	1,008	1,048	943	1,001	998	932
Male	898	985	1,067	1,167	1,216	1,237	1,271	1,201	1,266	1,240	1,186
White	1,311	1,422	1,551	1,685	1,734	1,772	1,827	1,680	1,780	1,742	1,653
Minority	296	312	311	319	296	289	283	261	263	260	241

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	50%	52%	53%	53%	52%	50%	50%	47%	48%	45%	41%
Female	49%	51%	51%	50%	48%	46%	47%	42%	43%	41%	37%
Male	52%	54%	55%	56%	56%	54%	53%	51%	52%	48%	45%
White	48%	50%	51%	51%	50%	48%	47%	43%	44%	41%	38%
Minority	47%	46%	45%	43%	39%	38%	37%	36%	35%	33%	31%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	211	232	247	254	255	258	268	240	236	220	201
Alcohol-related disorders	53	50	48	45	44	35	36	31	32	31	N.D.
Anxiety disorders	390	402	412	426	424	397	401	346	346	337	305
Attention-deficit, conduct, and disruptive behavior disorders	927	1,027	1,089	1,211	1,244	1,254	1,266	1,161	1,220	1,172	1,106
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33	32	32	33	33
Developmental disorders	398	478	539	626	701	775	851	846	932	977	934
Disorders usually diagnosed in infancy, childhood, adolescence	107	128	146	167	185	199	203	192	208	203	192
Impulse control disorders, NEC	114	125	123	127	124	127	128	117	125	116	97
Miscellaneous disorders	257	279	300	308	314	302	317	289	312	286	265
Mood disorders	596	612	653	679	657	621	613	527	541	508	446
Personality disorders	39	40	40	46	43	43	40	N.D.	N.D.	31	N.D.
Schizophrenia and other psychotic disorders	52	56	51	56	53	46	44	41	49	50	44
Screening and history of mental health and substance abuse	894	961	1,015	1,048	992	960	963	846	869	839	754
Substance-related disorders	105	113	107	123	123	110	106	82	92	81	61
Suicide and intentional self-inflicted injury	43	49	48	49	44	39	40	31	39	39	32



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Graves County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,025	2,162	2,313	2,404	2,492	2,537	2,583	2,375	2,539	2,528	2,493
Female	1,466	1,561	1,659	1,733	1,787	1,811	1,821	1,636	1,773	1,757	1,705
Male	559	601	654	671	705	726	762	739	766	771	788
White	1,704	1,818	1,962	2,041	2,127	2,169	2,206	2,031	2,175	2,163	2,104
Minority	256	278	280	289	288	281	280	244	259	263	273

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	65%	66%	67%	67%	66%	67%	62%	65%	64%	62%
Female	64%	67%	68%	69%	69%	68%	69%	62%	66%	66%	63%
Male	57%	59%	62%	62%	62%	62%	63%	62%	63%	61%	60%
White	62%	65%	66%	67%	67%	66%	67%	62%	65%	64%	61%
Minority	50%	52%	51%	52%	51%	50%	50%	44%	46%	47%	46%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	164	180	198	218	229	242	263	248	258	266	258
Alcohol-related disorders	166	185	207	219	230	241	245	232	241	241	222
Anxiety disorders	1,011	1,093	1,180	1,263	1,343	1,376	1,401	1,330	1,407	1,432	1,406
Attention-deficit, conduct, and disruptive behavior disorders	149	161	184	202	227	251	281	280	309	340	339
Delirium, dementia, and amnestic and other cognitive disorders	619	623	655	637	619	600	591	543	551	507	477
Developmental disorders	233	246	254	267	272	267	293	279	296	292	307
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	51	56	63	61	68	69	77	70	78	83	86
Miscellaneous disorders	378	416	451	484	516	535	572	547	566	593	579
Mood disorders	1,245	1,336	1,430	1,504	1,584	1,651	1,702	1,594	1,671	1,655	1,653
Personality disorders	133	137	159	159	170	167	188	181	197	194	186
Schizophrenia and other psychotic disorders	535	559	587	607	614	616	618	592	596	568	553
Screening and history of mental health and substance abuse	928	1,038	1,161	1,249	1,314	1,332	1,372	1,287	1,389	1,448	1,450
Substance-related disorders	270	304	342	338	365	386	402	385	394	397	376
Suicide and intentional self-inflicted injury	101	113	121	130	145	152	162	164	175	175	171



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Grayson County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,402	1,547	1,653	1,767	1,841	1,875	1,863	1,791	1,894	1,969	1,854
Female	700	759	778	829	851	857	842	806	837	864	813
Male	702	788	875	938	990	1,018	1,021	985	1,057	1,105	1,041
White	1,311	1,447	1,537	1,638	1,686	1,706	1,685	1,609	1,708	1,774	1,673
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	55%	55%	56%	56%	57%	55%	52%	54%	52%	47%
Female	54%	54%	53%	53%	54%	53%	52%	49%	49%	47%	43%
Male	55%	56%	57%	58%	59%	59%	59%	55%	58%	56%	51%
White	54%	54%	54%	54%	54%	55%	53%	50%	52%	50%	45%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	362	405	438	450	458	462	460	435	468	468	413
Alcohol-related disorders	51	49	50	52	49	47	43	38	36	35	N.D.
Anxiety disorders	470	513	561	588	607	598	606	601	620	631	582
Attention-deficit, conduct, and disruptive behavior disorders	708	819	904	963	1,010	1,053	1,036	1,004	1,044	1,064	992
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	289	322	374	414	472	488	517	517	571	623	600
Disorders usually diagnosed in infancy, childhood, adolescence	99	120	130	140	161	175	189	193	208	216	209
Impulse control disorders, NEC	115	128	136	136	134	131	131	114	117	114	95
Miscellaneous disorders	173	181	181	189	171	170	165	144	145	140	128
Mood disorders	597	638	633	646	645	596	555	511	497	482	414
Personality disorders	96	98	97	98	99	89	90	85	77	71	58
Schizophrenia and other psychotic disorders	43	47	45	43	43	41	39	36	37	36	32
Screening and history of mental health and substance abuse	853	910	921	952	955	945	902	818	835	813	721
Substance-related disorders	150	157	158	153	147	135	123	102	95	81	60
Suicide and intentional self-inflicted injury	31	38	41	38	37	36	36	36	36	33	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Grayson County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,526	1,589	1,701	1,865	1,957	2,022	2,043	1,942	2,138	2,211	2,120
Female	1,088	1,123	1,207	1,304	1,378	1,398	1,427	1,329	1,469	1,494	1,423
Male	438	466	494	561	579	624	616	613	669	717	697
White	1,407	1,457	1,564	1,723	1,814	1,878	1,893	1,787	1,973	2,033	1,945
Minority	N.D.	31	N.D.	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	60%	61%	63%	65%	66%	66%	62%	66%	66%	61%
Female	61%	63%	66%	67%	69%	70%	71%	66%	70%	69%	64%
Male	50%	53%	53%	56%	57%	60%	58%	56%	60%	60%	55%
White	56%	59%	61%	63%	65%	66%	66%	61%	65%	65%	60%
Minority	N.D.	16%	N.D.	16%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	165	176	189	210	217	235	235	232	260	277	280
Alcohol-related disorders	140	147	159	191	190	205	195	180	196	208	191
Anxiety disorders	902	953	1,038	1,153	1,225	1,256	1,273	1,220	1,344	1,394	1,345
Attention-deficit, conduct, and disruptive behavior disorders	60	58	71	92	112	118	141	153	174	205	224
Delirium, dementia, and amnestic and other cognitive disorders	339	320	321	314	320	319	322	297	301	285	278
Developmental disorders	135	139	144	155	156	163	166	166	175	182	186
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	34	33	32	39	38	50
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	36	42	44	42	47	53
Miscellaneous disorders	275	294	324	378	427	435	449	418	452	458	458
Mood disorders	893	966	1,065	1,178	1,248	1,318	1,355	1,290	1,439	1,510	1,453
Personality disorders	149	169	179	198	206	208	219	222	237	234	224
Schizophrenia and other psychotic disorders	305	314	320	329	332	332	349	331	343	346	327
Screening and history of mental health and substance abuse	726	806	888	1,001	1,086	1,162	1,171	1,104	1,207	1,228	1,185
Substance-related disorders	194	216	254	277	301	298	321	303	317	322	307
Suicide and intentional self-inflicted injury	41	44	51	57	64	66	75	73	82	77	79



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Green County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	643	681	705	744	765	761	780	747	802	821	752
Female	283	297	301	306	312	321	321	309	337	355	320
Male	360	384	404	438	453	440	459	438	465	466	432
White	592	629	645	684	692	688	703	669	723	743	672
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	60%	62%	60%	61%	62%	61%	57%	57%	54%	49%
Female	55%	58%	57%	55%	56%	58%	57%	53%	53%	50%	45%
Male	58%	62%	66%	64%	64%	65%	65%	60%	60%	58%	53%
White	56%	60%	61%	60%	59%	61%	60%	55%	55%	53%	48%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	170	180	186	189	186	177	183	158	167	169	153
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	298	312	299	326	330	333	345	328	335	323	289
Attention-deficit, conduct, and disruptive behavior disorders	262	276	290	308	322	313	333	315	334	326	305
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	151	159	177	193	211	223	240	247	268	280	275
Disorders usually diagnosed in infancy, childhood, adolescence	39	47	53	54	60	57	65	64	71	77	75
Impulse control disorders, NEC	56	57	61	61	62	59	61	54	58	58	45
Miscellaneous disorders	91	95	86	84	84	77	88	69	74	76	64
Mood disorders	248	249	238	236	235	218	215	186	183	183	153
Personality disorders	42	37	38	37	36	33	31	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	37	33	31	32	32	33	36	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	377	390	401	399	386	362	361	321	340	339	303
Substance-related disorders	67	61	60	62	55	47	39	34	32	31	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Green County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	894	923	990	970	969	965	1,008	972	1,045	1,054	994
Female	619	633	671	659	658	646	675	646	692	694	652
Male	275	290	319	311	311	319	333	326	353	360	342
White	786	812	873	850	854	846	888	862	933	947	890
Minority	48	49	52	51	46	43	38	33	36	32	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	70%	71%	70%	72%	73%	69%	72%	70%	66%
Female	66%	69%	73%	73%	73%	73%	75%	71%	73%	72%	67%
Male	60%	63%	66%	67%	65%	69%	70%	65%	68%	66%	64%
White	64%	68%	71%	71%	70%	71%	73%	68%	71%	69%	65%
Minority	28%	28%	31%	31%	29%	28%	25%	23%	24%	23%	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	78	84	99	101	114	111	123	126	135	136	138
Alcohol-related disorders	82	93	110	105	102	99	107	94	95	94	93
Anxiety disorders	498	529	575	577	602	601	641	634	690	711	678
Attention-deficit, conduct, and disruptive behavior disorders	40	43	52	51	55	61	66	62	74	81	81
Delirium, dementia, and amnestic and other cognitive disorders	280	279	279	268	252	243	243	223	218	200	178
Developmental disorders	50	47	56	56	63	63	65	72	75	75	76
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33
Miscellaneous disorders	186	210	232	229	234	240	247	234	252	236	225
Mood disorders	532	570	626	608	619	616	669	653	702	722	679
Personality disorders	118	129	142	138	140	140	151	140	142	141	135
Schizophrenia and other psychotic disorders	261	264	280	268	260	255	258	247	242	232	217
Screening and history of mental health and substance abuse	422	453	507	500	500	520	565	541	591	609	585
Substance-related disorders	104	122	133	126	135	142	155	136	154	156	152
Suicide and intentional self-inflicted injury	N.D.	32	37	35	38	41	54	49	49	45	45



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Greenup County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,695	1,922	2,011	2,073	2,146	2,159	2,179	2,050	2,133	2,105	1,896
Female	835	947	999	991	1,030	998	1,005	931	969	947	840
Male	860	975	1,012	1,082	1,116	1,161	1,174	1,119	1,164	1,158	1,056
White	1,611	1,819	1,883	1,932	1,981	1,981	1,981	1,865	1,926	1,895	1,710
Minority	N.D.	N.D.	32	N.D.	N.D.	32	35	N.D.	38	34	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	45%	48%	50%	50%	51%	51%	52%	48%	50%	48%	43%
Female	46%	49%	51%	49%	50%	49%	50%	46%	47%	45%	39%
Male	44%	48%	49%	51%	51%	53%	54%	51%	52%	52%	46%
White	44%	48%	49%	49%	49%	49%	50%	47%	48%	46%	41%
Minority	N.D.	N.D.	18%	N.D.	N.D.	13%	14%	N.D.	14%	12%	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	309	344	373	391	394	397	390	368	372	356	291
Alcohol-related disorders	89	81	79	73	69	62	53	50	45	39	N.D.
Anxiety disorders	514	568	566	561	565	558	547	519	530	523	467
Attention-deficit, conduct, and disruptive behavior disorders	766	866	927	984	1,058	1,082	1,076	1,020	1,043	1,027	925
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	282	324	388	426	464	494	522	522	584	589	551
Disorders usually diagnosed in infancy, childhood, adolescence	124	145	162	176	197	202	208	196	203	194	175
Impulse control disorders, NEC	59	62	59	57	57	54	52	47	43	36	34
Miscellaneous disorders	180	195	208	209	211	207	201	181	189	178	157
Mood disorders	687	729	721	707	695	645	613	540	518	483	391
Personality disorders	80	76	76	69	64	56	51	44	41	35	N.D.
Schizophrenia and other psychotic disorders	74	71	66	60	57	52	50	48	51	43	32
Screening and history of mental health and substance abuse	1,276	1,429	1,484	1,494	1,544	1,538	1,501	1,369	1,381	1,327	1,168
Substance-related disorders	230	241	216	197	194	164	145	128	133	116	99
Suicide and intentional self-inflicted injury	54	56	56	56	51	42	42	43	46	38	34



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Greenup County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,175	2,388	2,514	2,674	2,728	2,715	2,801	2,531	2,671	2,638	2,485
Female	1,531	1,651	1,763	1,861	1,883	1,891	1,947	1,760	1,842	1,801	1,692
Male	644	737	751	813	845	824	854	771	829	837	793
White	1,985	2,185	2,307	2,469	2,530	2,511	2,587	2,329	2,455	2,416	2,268
Minority	58	61	59	52	42	39	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	64%	64%	66%	65%	65%	67%	61%	66%	66%	61%
Female	64%	67%	68%	69%	68%	68%	70%	63%	68%	68%	64%
Male	56%	59%	57%	60%	60%	59%	62%	57%	61%	61%	56%
White	61%	63%	64%	65%	65%	64%	66%	60%	65%	65%	60%
Minority	21%	22%	21%	20%	16%	15%	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	148	152	174	191	201	218	248	213	233	245	229
Alcohol-related disorders	242	267	283	300	314	312	341	292	309	285	280
Anxiety disorders	1,155	1,309	1,390	1,494	1,551	1,572	1,648	1,522	1,611	1,566	1,468
Attention-deficit, conduct, and disruptive behavior disorders	184	204	216	249	258	270	300	270	310	314	302
Delirium, dementia, and amnestic and other cognitive disorders	558	582	589	589	572	562	570	528	511	464	408
Developmental disorders	264	281	296	312	313	321	322	297	304	281	265
Disorders usually diagnosed in infancy, childhood, adolescence	32	32	36	40	41	45	50	51	55	54	65
Impulse control disorders, NEC	43	52	56	60	66	66	70	65	65	65	62
Miscellaneous disorders	399	432	460	498	515	531	573	514	525	504	461
Mood disorders	1,242	1,422	1,531	1,644	1,701	1,743	1,819	1,655	1,748	1,714	1,613
Personality disorders	334	367	384	408	426	424	423	365	364	333	315
Schizophrenia and other psychotic disorders	596	638	649	671	668	655	686	623	623	602	544
Screening and history of mental health and substance abuse	1,250	1,446	1,566	1,729	1,781	1,749	1,818	1,627	1,731	1,725	1,647
Substance-related disorders	358	422	454	510	539	581	641	582	632	604	559
Suicide and intentional self-inflicted injury	86	98	101	114	114	130	150	137	150	134	130



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Hancock County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	343	375	396	417	420	429	448	411	443	450	442
Female	161	179	183	191	188	188	199	174	181	191	188
Male	182	196	213	226	232	241	249	237	262	259	254
White	N.D.	344	363	386	387	N.D.	404	372	399	409	399
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	48%	52%	55%	54%	52%	52%	52%	50%	51%	45%	42%
Female	48%	52%	52%	52%	48%	47%	47%	44%	44%	41%	39%
Male	49%	51%	57%	56%	56%	57%	57%	56%	57%	49%	45%
White	47%	50%	53%	53%	51%	52%	50%	48%	49%	43%	40%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	77	82	91	97	86	82	95	83	88	86	87
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	94	110	113	120	126	135	138	129	137	143	139
Attention-deficit, conduct, and disruptive behavior disorders	137	156	167	185	195	202	210	191	203	202	202
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	90	98	110	125	137	142	160	161	179	178	179
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	35	41	41	47	46	56	59	58
Impulse control disorders, NEC	31	N.D.	N.D.	32	31	N.D.	32	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	46	51	57	60	61	61	65	59	61	63	54
Mood disorders	128	141	147	156	154	157	151	126	126	131	126
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	174	186	196	203	192	172	170	149	141	140	142
Substance-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Hancock County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	306	335	356	396	404	390	433	401	456	463	450
Female	226	233	248	272	279	265	286	260	286	302	289
Male	80	102	108	124	125	125	147	141	170	161	161
White	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	400	N.D.	424	430	416
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	63%	64%	69%	67%	64%	69%	63%	68%	66%	62%
Female	62%	65%	65%	70%	67%	65%	70%	63%	67%	69%	63%
Male	55%	59%	62%	68%	66%	62%	67%	64%	69%	62%	59%
White	60%	62%	63%	69%	67%	64%	68%	63%	67%	65%	61%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	34	38	43	46	53	41	45	36	46	58	46
Alcohol-related disorders	N.D.	31	34	41	44	32	43	41	48	49	43
Anxiety disorders	149	167	172	201	205	201	225	212	243	236	231
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	N.D.	31	40	39	45	43	51	52	53
Delirium, dementia, and amnesic and other cognitive disorders	67	64	72	68	62	67	67	65	66	67	67
Developmental disorders	43	44	42	46	50	49	51	47	53	54	52
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	62	73	75	89	106	97	98	81	98	104	103
Mood disorders	180	193	221	248	250	236	260	249	274	277	270
Personality disorders	32	31	34	40	41	37	38	32	32	33	31
Schizophrenia and other psychotic disorders	86	93	102	109	107	102	116	114	121	115	108
Screening and history of mental health and substance abuse	155	188	194	232	255	241	267	243	288	283	266
Substance-related disorders	51	56	59	67	73	71	74	58	62	72	69
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Hardin County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,454	3,959	4,305	4,510	4,596	4,713	4,905	4,615	4,869	5,028	4,703
Female	1,621	1,843	1,969	2,042	2,069	2,103	2,209	2,077	2,216	2,232	2,084
Male	1,833	2,116	2,336	2,468	2,527	2,610	2,696	2,538	2,653	2,796	2,619
White	2,592	2,971	3,230	3,366	3,425	3,483	3,592	3,368	3,550	3,677	3,423
Minority	633	711	760	779	756	783	813	748	800	826	789

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	47%	49%	51%	52%	52%	51%	53%	49%	49%	47%	43%
Female	45%	47%	49%	50%	49%	48%	49%	45%	46%	43%	39%
Male	48%	51%	53%	54%	55%	55%	56%	52%	53%	51%	46%
White	47%	50%	52%	53%	52%	51%	52%	48%	49%	47%	42%
Minority	33%	34%	34%	34%	32%	32%	33%	30%	31%	29%	27%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	826	963	1,073	1,138	1,166	1,189	1,238	1,148	1,206	1,260	1,190
Alcohol-related disorders	158	173	169	169	157	155	139	117	108	96	67
Anxiety disorders	979	1,101	1,204	1,231	1,244	1,267	1,337	1,269	1,340	1,339	1,239
Attention-deficit, conduct, and disruptive behavior disorders	1,957	2,255	2,451	2,571	2,695	2,756	2,872	2,679	2,762	2,755	2,551
Delirium, dementia, and amnesic and other cognitive disorders	48	53	59	58	59	60	64	56	61	61	55
Developmental disorders	724	882	1,026	1,156	1,248	1,350	1,452	1,467	1,614	1,708	1,651
Disorders usually diagnosed in infancy, childhood, adolescence	340	395	442	470	497	527	555	545	574	585	553
Impulse control disorders, NEC	453	489	497	515	514	499	493	440	416	381	320
Miscellaneous disorders	382	403	412	421	422	423	414	395	395	384	336
Mood disorders	1,291	1,436	1,469	1,458	1,438	1,372	1,360	1,249	1,211	1,135	978
Personality disorders	295	304	314	317	322	305	301	270	255	229	201
Schizophrenia and other psychotic disorders	140	135	155	155	155	147	160	143	141	136	120
Screening and history of mental health and substance abuse	2,302	2,555	2,686	2,684	2,557	2,443	2,413	2,125	2,101	2,083	1,921
Substance-related disorders	337	357	370	368	367	351	326	271	253	227	176
Suicide and intentional self-inflicted injury	74	83	84	83	86	84	85	84	84	91	81



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Hardin County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,895	3,198	3,439	3,656	3,871	3,966	4,087	3,770	4,164	4,278	4,036
Female	2,148	2,393	2,558	2,714	2,892	2,949	3,004	2,739	3,005	3,099	2,899
Male	747	805	881	942	979	1,017	1,083	1,031	1,159	1,179	1,137
White	2,330	2,566	2,748	2,927	3,127	3,210	3,335	3,094	3,416	3,498	3,282
Minority	409	462	510	536	533	530	520	447	502	532	498

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	58%	59%	61%	62%	62%	63%	58%	62%	61%	57%
Female	56%	60%	60%	63%	64%	64%	66%	59%	64%	63%	58%
Male	51%	53%	54%	56%	56%	56%	58%	55%	59%	57%	54%
White	56%	60%	60%	62%	63%	63%	65%	59%	64%	63%	58%
Minority	35%	39%	40%	41%	41%	39%	40%	35%	37%	38%	34%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	254	296	317	344	369	392	429	411	440	464	435
Alcohol-related disorders	242	291	334	351	365	382	407	366	389	395	374
Anxiety disorders	1,368	1,530	1,686	1,832	1,964	2,021	2,167	2,058	2,243	2,278	2,181
Attention-deficit, conduct, and disruptive behavior disorders	200	234	261	313	368	381	432	427	521	591	587
Delirium, dementia, and amnesic and other cognitive disorders	679	698	705	694	700	689	702	658	674	657	596
Developmental disorders	307	327	337	373	385	386	401	402	437	466	486
Disorders usually diagnosed in infancy, childhood, adolescence	55	61	66	76	80	79	95	93	106	125	137
Impulse control disorders, NEC	66	80	97	104	121	132	142	149	177	170	175
Miscellaneous disorders	616	709	778	875	914	963	1,002	921	953	963	906
Mood disorders	1,741	1,975	2,129	2,276	2,428	2,510	2,651	2,478	2,753	2,807	2,648
Personality disorders	357	400	431	463	470	480	496	462	514	531	517
Schizophrenia and other psychotic disorders	698	758	787	798	809	817	833	802	840	823	759
Screening and history of mental health and substance abuse	1,414	1,618	1,823	2,017	2,201	2,216	2,257	2,050	2,230	2,279	2,119
Substance-related disorders	362	428	460	475	511	546	592	532	579	600	567
Suicide and intentional self-inflicted injury	74	87	102	107	120	126	150	144	157	141	140



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Harlan County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,044	3,187	3,260	3,359	3,313	3,275	3,217	3,022	3,053	2,880	2,683
Female	1,605	1,640	1,608	1,630	1,600	1,586	1,556	1,447	1,455	1,345	1,252
Male	1,439	1,547	1,652	1,729	1,713	1,689	1,661	1,575	1,598	1,535	1,431
White	2,819	2,921	2,965	3,046	2,992	2,938	2,865	2,684	2,701	2,526	2,345
Minority	100	113	113	110	98	107	108	107	106	106	111

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	50%	52%	54%	56%	55%	57%	57%	54%	55%	53%	50%
Female	53%	54%	54%	55%	55%	56%	57%	53%	53%	51%	48%
Male	47%	50%	54%	56%	56%	57%	58%	56%	57%	56%	52%
White	49%	51%	53%	54%	54%	55%	55%	52%	53%	51%	48%
Minority	29%	30%	27%	26%	22%	24%	24%	24%	23%	22%	23%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	464	486	502	513	487	480	480	427	424	388	344
Alcohol-related disorders	221	212	204	193	179	171	142	123	100	79	62
Anxiety disorders	1,143	1,171	1,192	1,245	1,236	1,231	1,176	1,081	1,066	999	910
Attention-deficit, conduct, and disruptive behavior disorders	1,030	1,141	1,203	1,271	1,270	1,246	1,261	1,175	1,173	1,106	1,035
Delirium, dementia, and amnestic and other cognitive disorders	36	37	36	37	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	1,083	1,190	1,295	1,396	1,447	1,462	1,470	1,377	1,382	1,308	1,224
Disorders usually diagnosed in infancy, childhood, adolescence	134	149	166	184	191	201	209	195	187	183	168
Impulse control disorders, NEC	95	105	105	104	97	93	82	77	70	57	47
Miscellaneous disorders	383	389	385	377	362	338	329	308	300	285	271
Mood disorders	860	853	842	829	773	725	668	579	520	456	393
Personality disorders	213	214	213	215	218	207	182	157	137	111	97
Schizophrenia and other psychotic disorders	112	107	110	109	102	94	85	77	67	57	45
Screening and history of mental health and substance abuse	1,392	1,401	1,401	1,364	1,287	1,214	1,099	991	982	889	781
Substance-related disorders	506	477	458	449	413	390	342	313	266	243	216
Suicide and intentional self-inflicted injury	123	121	109	109	106	104	90	79	70	59	48



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Harlan County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,846	4,055	4,158	4,266	4,397	4,407	4,452	4,154	4,306	4,281	4,139
Female	2,567	2,687	2,775	2,833	2,925	2,934	2,967	2,759	2,865	2,823	2,734
Male	1,279	1,368	1,383	1,433	1,472	1,473	1,485	1,395	1,441	1,458	1,405
White	3,470	3,669	3,762	3,861	4,005	4,027	4,070	3,793	3,951	3,912	3,769
Minority	177	176	183	176	163	144	131	114	110	117	114

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	65%	65%	66%	67%	68%	69%	65%	68%	68%	66%
Female	67%	70%	70%	70%	71%	72%	73%	69%	72%	72%	70%
Male	56%	58%	58%	59%	60%	60%	62%	59%	62%	62%	60%
White	62%	65%	65%	65%	66%	67%	68%	65%	68%	68%	65%
Minority	32%	32%	34%	33%	31%	29%	26%	24%	23%	23%	23%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	385	423	448	468	498	496	523	491	488	500	484
Alcohol-related disorders	467	505	510	523	533	542	572	531	562	560	548
Anxiety disorders	2,435	2,582	2,692	2,803	2,888	2,904	2,948	2,790	2,874	2,859	2,801
Attention-deficit, conduct, and disruptive behavior disorders	184	201	217	239	267	288	322	326	350	380	378
Delirium, dementia, and amnesic and other cognitive disorders	652	673	658	637	609	600	582	537	524	506	467
Developmental disorders	364	379	414	449	472	477	480	476	486	511	497
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	33	34	35	36	41	49	55	55
Impulse control disorders, NEC	41	47	49	59	62	63	74	65	75	81	81
Miscellaneous disorders	826	857	921	959	995	985	978	915	928	907	885
Mood disorders	2,140	2,279	2,388	2,515	2,598	2,613	2,686	2,513	2,595	2,585	2,521
Personality disorders	400	423	451	486	523	520	529	495	495	498	480
Schizophrenia and other psychotic disorders	711	741	746	755	741	733	750	711	718	695	652
Screening and history of mental health and substance abuse	2,032	2,181	2,300	2,387	2,507	2,562	2,635	2,479	2,580	2,563	2,528
Substance-related disorders	890	948	972	1,047	1,085	1,138	1,189	1,087	1,156	1,157	1,148
Suicide and intentional self-inflicted injury	171	185	200	213	229	248	264	263	279	283	276



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Harrison County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	820	907	971	993	1,010	1,057	1,077	1,047	1,111	1,122	1,064
Female	401	428	451	469	479	491	498	479	509	513	479
Male	419	479	520	524	531	566	579	568	602	609	585
White	725	805	862	882	892	923	938	918	971	986	943
Minority	43	44	42	42	44	51	46	46	47	42	39

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	57%	58%	58%	56%	58%	57%	53%	54%	51%	47%
Female	55%	56%	57%	57%	55%	55%	55%	49%	51%	48%	44%
Male	54%	58%	60%	58%	57%	60%	59%	55%	57%	54%	51%
White	53%	55%	57%	56%	54%	56%	55%	51%	52%	49%	46%
Minority	29%	32%	28%	27%	26%	28%	26%	25%	26%	21%	20%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	179	200	212	231	223	232	222	193	203	202	178
Alcohol-related disorders	35	39	40	34	N.D.	N.D.	32	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	289	314	334	348	367	390	408	398	409	411	387
Attention-deficit, conduct, and disruptive behavior disorders	340	379	419	437	448	454	462	438	447	440	415
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	168	190	205	229	247	271	297	307	325	340	329
Disorders usually diagnosed in infancy, childhood, adolescence	56	64	72	79	88	87	96	94	97	101	96
Impulse control disorders, NEC	32	32	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	126	137	144	144	145	155	155	153	166	163	157
Mood disorders	280	282	299	285	271	263	250	211	222	207	168
Personality disorders	45	41	39	35	N.D.	31	33	31	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	42	46	48	42	36	38	32	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	501	527	562	532	520	503	494	439	447	419	385
Substance-related disorders	102	106	108	99	93	99	88	80	83	61	45
Suicide and intentional self-inflicted injury	32	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Harrison County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	884	983	1,053	1,076	1,091	1,143	1,201	1,162	1,250	1,320	1,304
Female	619	684	753	770	772	813	852	796	857	911	888
Male	265	299	300	306	319	330	349	366	393	409	416
White	784	882	945	964	988	1,036	1,092	1,069	1,153	1,212	1,193
Minority	56	54	58	57	46	45	41	32	37	43	42

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	65%	67%	67%	68%	68%	70%	64%	70%	71%	65%
Female	65%	67%	70%	71%	71%	71%	73%	65%	71%	74%	68%
Male	57%	60%	60%	60%	61%	61%	65%	63%	67%	65%	61%
White	62%	65%	67%	67%	68%	69%	71%	65%	70%	71%	65%
Minority	37%	34%	35%	34%	28%	27%	26%	21%	24%	28%	27%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	85	97	108	121	129	139	153	147	152	163	175
Alcohol-related disorders	81	92	93	103	106	116	133	126	143	149	142
Anxiety disorders	456	526	571	604	625	658	712	687	724	764	779
Attention-deficit, conduct, and disruptive behavior disorders	88	92	102	113	117	136	149	144	150	163	164
Delirium, dementia, and amnestic and other cognitive disorders	273	274	275	266	266	284	302	282	282	273	256
Developmental disorders	88	96	103	98	103	115	127	119	124	127	125
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	158	188	216	240	248	279	300	283	295	301	297
Mood disorders	481	537	583	633	644	696	759	721	769	816	819
Personality disorders	77	85	96	103	108	116	130	126	134	125	127
Schizophrenia and other psychotic disorders	229	238	251	254	260	258	281	273	273	272	262
Screening and history of mental health and substance abuse	447	526	590	622	634	662	697	682	724	776	764
Substance-related disorders	104	127	151	170	184	212	235	229	272	275	291
Suicide and intentional self-inflicted injury	48	52	54	66	72	76	91	82	86	86	94



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Hart County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,032	1,102	1,155	1,176	1,209	1,246	1,254	1,211	1,266	1,281	1,217
Female	495	528	552	562	561	559	557	540	575	588	546
Male	537	574	603	614	648	687	697	671	691	693	671
White	924	979	1,025	1,040	1,072	1,109	1,101	1,065	1,114	1,124	1,064
Minority	62	70	71	76	72	69	69	67	71	69	71

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	51%	51%	53%	54%	55%	56%	56%	52%	53%	49%	44%
Female	51%	52%	53%	55%	54%	55%	54%	50%	52%	45%	41%
Male	51%	50%	53%	54%	55%	58%	58%	54%	55%	52%	48%
White	51%	51%	53%	54%	54%	56%	55%	51%	52%	48%	43%
Minority	29%	31%	32%	33%	33%	32%	31%	30%	30%	26%	27%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	181	194	219	217	213	219	216	200	205	201	187
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	348	372	383	394	406	419	431	415	429	427	405
Attention-deficit, conduct, and disruptive behavior disorders	464	515	549	570	593	615	631	589	603	602	569
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	218	250	286	303	329	345	374	375	403	416	409
Disorders usually diagnosed in infancy, childhood, adolescence	76	90	102	114	121	126	130	132	133	139	135
Impulse control disorders, NEC	63	72	72	70	69	65	64	55	54	48	44
Miscellaneous disorders	190	191	193	206	204	203	191	180	184	176	171
Mood disorders	389	400	402	398	388	364	351	316	310	288	240
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	31	31	34	35	36	36	31	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	556	595	603	582	573	546	530	484	480	476	421
Substance-related disorders	56	63	57	57	59	55	49	44	44	39	31
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Hart County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,339	1,419	1,471	1,506	1,552	1,600	1,668	1,532	1,615	1,640	1,609
Female	917	994	1,023	1,040	1,082	1,097	1,126	1,032	1,086	1,091	1,071
Male	422	425	448	466	470	503	542	500	529	549	538
White	1,129	1,208	1,256	1,280	1,319	1,358	1,425	1,321	1,407	1,424	1,395
Minority	139	136	140	149	148	150	144	123	120	127	124

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	68%	70%	69%	69%	70%	65%	67%	67%	64%
Female	67%	71%	72%	73%	73%	72%	73%	67%	70%	70%	67%
Male	59%	59%	62%	64%	62%	63%	65%	61%	62%	60%	60%
White	65%	68%	69%	70%	69%	69%	70%	65%	68%	67%	64%
Minority	41%	40%	43%	44%	44%	42%	42%	37%	36%	39%	37%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	154	178	178	194	200	209	231	194	204	200	188
Alcohol-related disorders	108	121	134	135	135	146	150	139	147	141	138
Anxiety disorders	726	792	827	855	906	940	1,001	925	983	992	948
Attention-deficit, conduct, and disruptive behavior disorders	53	59	66	73	83	91	113	105	123	143	147
Delirium, dementia, and amnestic and other cognitive disorders	406	404	396	393	384	367	366	344	316	281	273
Developmental disorders	125	132	135	136	140	143	147	133	139	142	137
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	32	36	35	33	42	40
Miscellaneous disorders	321	358	368	391	407	409	446	414	442	454	433
Mood disorders	820	882	929	964	1,016	1,031	1,087	1,007	1,076	1,101	1,082
Personality disorders	99	102	110	109	105	109	116	108	105	103	103
Schizophrenia and other psychotic disorders	326	334	338	345	340	345	363	341	325	313	294
Screening and history of mental health and substance abuse	659	714	775	792	818	863	917	849	915	927	915
Substance-related disorders	161	170	182	193	200	205	230	211	222	245	232
Suicide and intentional self-inflicted injury	41	43	45	44	48	49	55	56	57	66	65



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Henderson County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,425	2,572	2,725	2,889	2,993	3,004	3,031	2,879	2,983	3,031	2,960
Female	1,189	1,250	1,315	1,371	1,396	1,405	1,418	1,338	1,355	1,353	1,308
Male	1,236	1,322	1,410	1,518	1,597	1,599	1,613	1,541	1,628	1,678	1,652
White	1,924	2,021	2,116	2,251	2,329	2,340	2,371	2,244	2,303	2,346	2,287
Minority	349	367	403	415	412	400	376	350	376	379	381

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	62%	64%	64%	64%	62%	62%	57%	58%	54%	50%
Female	60%	60%	62%	62%	61%	59%	59%	54%	54%	50%	45%
Male	61%	63%	65%	66%	66%	65%	65%	60%	61%	59%	55%
White	60%	61%	63%	64%	63%	61%	61%	56%	56%	53%	49%
Minority	42%	42%	42%	42%	41%	39%	37%	34%	35%	32%	32%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	378	408	446	456	460	462	471	430	416	394	369
Alcohol-related disorders	98	98	90	89	74	65	64	63	57	49	45
Anxiety disorders	711	744	783	799	806	804	785	711	690	648	596
Attention-deficit, conduct, and disruptive behavior disorders	1,494	1,618	1,747	1,875	1,995	2,009	2,020	1,880	1,892	1,910	1,857
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	34	37	34	36	33	35	31	31	N.D.	N.D.
Developmental disorders	576	669	755	825	904	932	978	939	1,011	1,063	1,069
Disorders usually diagnosed in infancy, childhood, adolescence	167	184	214	234	258	274	275	257	278	284	281
Impulse control disorders, NEC	211	223	238	245	260	245	243	212	191	183	177
Miscellaneous disorders	368	391	401	418	447	470	467	451	486	490	484
Mood disorders	1,069	1,109	1,144	1,150	1,147	1,102	1,069	947	888	825	757
Personality disorders	89	91	91	84	96	86	87	69	63	56	52
Schizophrenia and other psychotic disorders	106	102	108	107	98	99	100	79	69	64	61
Screening and history of mental health and substance abuse	1,379	1,445	1,502	1,544	1,481	1,417	1,383	1,278	1,264	1,220	1,153
Substance-related disorders	271	280	271	262	232	205	192	168	150	129	115
Suicide and intentional self-inflicted injury	92	88	94	90	91	84	91	74	75	64	65



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Henderson County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,292	2,518	2,651	2,783	2,910	2,860	2,944	2,784	3,078	3,190	2,993
Female	1,698	1,857	1,966	2,067	2,153	2,106	2,177	2,067	2,277	2,358	2,193
Male	594	661	685	716	757	754	767	717	801	832	800
White	1,895	2,078	2,155	2,279	2,381	2,357	2,436	2,301	2,553	2,650	2,449
Minority	317	347	387	388	403	374	367	341	375	380	374

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	68%	69%	69%	70%	69%	69%	64%	69%	68%	63%
Female	67%	70%	71%	71%	71%	71%	71%	66%	72%	71%	66%
Male	59%	64%	63%	64%	65%	64%	65%	59%	63%	63%	58%
White	66%	70%	69%	70%	70%	69%	70%	64%	70%	69%	64%
Minority	46%	49%	53%	50%	52%	49%	49%	45%	47%	45%	43%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	148	159	164	173	193	212	218	213	240	258	242
Alcohol-related disorders	209	215	239	244	256	260	276	272	283	289	275
Anxiety disorders	1,244	1,402	1,476	1,577	1,670	1,644	1,688	1,652	1,822	1,880	1,783
Attention-deficit, conduct, and disruptive behavior disorders	265	312	350	419	478	504	582	600	720	789	767
Delirium, dementia, and amnestic and other cognitive disorders	520	548	544	537	538	528	514	479	487	476	427
Developmental disorders	201	215	218	227	243	256	266	283	291	306	317
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	34	35	44	47	57	63	67	77	80
Impulse control disorders, NEC	45	47	52	60	65	70	78	86	98	108	102
Miscellaneous disorders	492	559	598	639	658	667	701	677	733	774	711
Mood disorders	1,460	1,646	1,742	1,845	1,924	1,901	1,961	1,884	2,067	2,144	2,049
Personality disorders	153	164	177	182	185	188	195	199	223	215	198
Schizophrenia and other psychotic disorders	404	441	464	460	473	463	482	468	494	509	504
Screening and history of mental health and substance abuse	1,130	1,296	1,401	1,489	1,578	1,560	1,620	1,528	1,708	1,741	1,612
Substance-related disorders	384	439	474	522	565	597	625	627	719	726	670
Suicide and intentional self-inflicted injury	84	95	100	108	124	126	148	161	184	200	189



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Henry County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	680	735	801	901	912	898	938	905	911	911	872
Female	314	334	379	417	436	421	430	403	403	399	384
Male	366	401	422	484	476	477	508	502	508	512	488
White	600	646	703	790	789	777	807	782	781	780	747
Minority	46	46	44	48	51	48	45	35	38	39	37

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	57%	57%	60%	60%	58%	58%	54%	54%	50%	45%
Female	55%	54%	55%	57%	58%	56%	55%	51%	50%	46%	42%
Male	56%	59%	60%	64%	61%	60%	61%	57%	57%	53%	49%
White	55%	56%	56%	60%	59%	56%	56%	52%	51%	48%	43%
Minority	34%	33%	29%	26%	27%	28%	26%	21%	22%	20%	19%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	187	212	226	262	272	261	278	264	259	252	226
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	230	254	285	336	333	326	331	318	301	291	267
Attention-deficit, conduct, and disruptive behavior disorders	334	373	405	443	454	462	478	463	452	447	416
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	119	143	163	195	209	219	252	259	276	278	274
Disorders usually diagnosed in infancy, childhood, adolescence	54	55	65	73	75	84	92	93	90	90	93
Impulse control disorders, NEC	33	40	37	38	36	35	37	37	33	33	N.D.
Miscellaneous disorders	55	60	64	73	71	63	67	66	61	56	57
Mood disorders	240	239	254	275	269	259	265	238	235	227	201
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	369	377	401	431	418	397	379	344	307	307	291
Substance-related disorders	51	47	46	48	42	38	36	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Henry County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	568	614	671	679	749	773	799	746	840	844	793
Female	425	450	495	500	547	560	570	534	592	593	536
Male	143	164	176	179	202	213	229	212	248	251	257
White	501	543	590	600	672	688	713	664	750	748	699
Minority	40	42	44	42	41	46	45	44	49	50	48

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	61%	63%	64%	69%	68%	70%	63%	70%	69%	63%
Female	59%	62%	67%	67%	72%	71%	72%	67%	72%	71%	64%
Male	49%	57%	54%	56%	61%	62%	64%	55%	65%	65%	62%
White	56%	62%	64%	64%	70%	68%	70%	63%	70%	68%	62%
Minority	34%	33%	31%	33%	31%	38%	37%	34%	37%	40%	38%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	49	58	70	66	77	86	85	78	107	108	94
Alcohol-related disorders	41	54	55	60	65	72	77	70	80	77	71
Anxiety disorders	277	306	353	364	396	418	448	436	486	489	475
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	35	38	40	42	56	51	80	88	83
Delirium, dementia, and amnestic and other cognitive disorders	138	150	152	133	142	147	144	128	130	113	102
Developmental disorders	62	64	68	63	69	67	77	72	75	75	70
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	92	96	106	113	122	141	123	103	121	116	97
Mood disorders	290	325	370	376	417	433	467	432	483	490	476
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	129	134	146	143	146	145	142	132	119	110	109
Screening and history of mental health and substance abuse	289	321	362	401	438	450	465	425	495	497	453
Substance-related disorders	60	69	78	87	93	92	98	88	101	100	92
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Hickman County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	228	245	260	258	275	283	277	250	244	241	230
Female	112	117	126	124	124	124	126	117	105	98	90
Male	116	128	134	134	151	159	151	133	139	143	140
White	171	183	191	192	207	218	208	182	176	168	164
Minority	45	48	53	48	47	44	47	47	45	48	42

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	48%	53%	54%	51%	52%	52%	53%	48%	48%	48%	42%
Female	47%	52%	50%	47%	47%	46%	48%	45%	41%	37%	33%
Male	50%	55%	58%	56%	57%	58%	57%	52%	55%	59%	52%
White	48%	52%	53%	50%	51%	51%	50%	45%	44%	44%	39%
Minority	40%	43%	45%	41%	39%	37%	42%	40%	41%	40%	36%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	31	36	42	36	42	47	46	39	34	N.D.	N.D.
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	52	52	57	54	62	56	50	43	39	36	35
Attention-deficit, conduct, and disruptive behavior disorders	107	118	127	136	147	153	141	131	127	119	109
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	39	50	63	67	75	82	95	95	99	106	106
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	N.D.	N.D.	N.D.	31	34	35	31	N.D.	N.D.	N.D.	N.D.
Mood disorders	77	81	86	87	91	88	78	71	61	53	47
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	137	143	136	136	132	126	117	101	96	93	79
Substance-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Hickman County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	344	375	374	366	385	400	391	358	377	359	321
Female	249	270	263	266	281	290	277	260	269	255	232
Male	95	105	111	100	104	110	114	98	108	104	89
White	277	300	298	288	299	314	305	274	295	280	249
Minority	58	65	65	68	75	76	73	73	69	65	60

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	66%	69%	70%	68%	70%	63%	70%	70%	62%
Female	65%	69%	67%	70%	71%	71%	71%	65%	71%	71%	63%
Male	61%	61%	64%	65%	67%	63%	67%	59%	66%	70%	62%
White	66%	68%	67%	70%	72%	70%	71%	64%	71%	72%	65%
Minority	49%	53%	52%	57%	55%	56%	54%	54%	56%	55%	46%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Alcohol-related disorders	32	38	42	41	41	37	41	40	42	47	35
Anxiety disorders	170	185	187	186	192	206	210	190	201	195	166
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Delirium, dementia, and amnestic and other cognitive disorders	151	154	141	127	138	141	139	122	113	91	74
Developmental disorders	44	50	49	45	47	48	51	40	38	36	32
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	53	63	70	70	73	68	71	70	70	67	60
Mood disorders	219	237	239	234	244	270	262	237	264	235	213
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	92	97	91	94	96	91	102	86	89	73	63
Screening and history of mental health and substance abuse	152	172	181	180	190	183	196	183	196	190	170
Substance-related disorders	39	45	52	50	54	53	58	58	57	58	52
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Hopkins County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,673	2,873	3,007	3,132	3,218	3,281	3,242	2,942	3,124	3,067	2,855
Female	1,316	1,376	1,388	1,458	1,475	1,498	1,464	1,319	1,388	1,365	1,251
Male	1,357	1,497	1,619	1,674	1,743	1,783	1,778	1,623	1,736	1,702	1,604
White	2,169	2,310	2,392	2,465	2,535	2,555	2,509	2,276	2,421	2,367	2,218
Minority	300	324	339	362	355	369	345	298	326	321	291

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	55%	58%	60%	61%	60%	60%	55%	57%	53%	48%
Female	53%	54%	55%	58%	59%	58%	57%	52%	52%	49%	43%
Male	54%	56%	60%	62%	62%	62%	63%	59%	61%	56%	52%
White	53%	55%	57%	59%	59%	59%	58%	54%	55%	51%	46%
Minority	32%	33%	33%	36%	34%	35%	33%	28%	30%	27%	25%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	510	551	550	560	567	560	530	469	477	468	437
Alcohol-related disorders	68	67	69	61	56	52	40	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	797	814	849	851	841	839	825	707	723	683	609
Attention-deficit, conduct, and disruptive behavior disorders	1,402	1,512	1,613	1,714	1,774	1,832	1,821	1,652	1,683	1,619	1,524
Delirium, dementia, and amnesic and other cognitive disorders	52	54	52	56	55	63	62	56	57	52	42
Developmental disorders	494	619	728	811	910	996	1,073	1,045	1,137	1,157	1,106
Disorders usually diagnosed in infancy, childhood, adolescence	216	245	272	305	320	350	373	348	343	343	332
Impulse control disorders, NEC	265	273	289	282	277	283	255	223	215	206	185
Miscellaneous disorders	549	551	566	592	600	588	573	510	511	496	454
Mood disorders	1,130	1,196	1,192	1,236	1,192	1,154	1,079	930	926	868	771
Personality disorders	55	57	57	51	45	38	37	33	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	92	88	96	94	93	87	83	72	70	69	61
Screening and history of mental health and substance abuse	1,435	1,515	1,538	1,560	1,519	1,491	1,411	1,215	1,233	1,173	1,072
Substance-related disorders	238	231	219	218	200	185	163	132	136	117	101
Suicide and intentional self-inflicted injury	103	98	97	94	89	83	72	58	58	55	44



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Hopkins County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,877	3,069	3,217	3,306	3,357	3,440	3,571	3,239	3,475	3,495	3,384
Female	2,067	2,190	2,310	2,380	2,429	2,459	2,534	2,280	2,462	2,466	2,355
Male	810	879	907	926	928	981	1,037	959	1,013	1,029	1,029
White	2,452	2,606	2,750	2,809	2,863	2,932	3,066	2,759	2,970	2,999	2,868
Minority	316	346	346	363	353	355	340	314	340	325	335

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	66%	68%	70%	70%	70%	73%	66%	70%	69%	65%
Female	66%	69%	70%	72%	72%	73%	75%	68%	73%	72%	67%
Male	58%	61%	62%	64%	65%	65%	68%	63%	65%	63%	60%
White	64%	67%	68%	70%	70%	71%	73%	66%	70%	70%	65%
Minority	45%	48%	48%	51%	49%	48%	48%	43%	47%	43%	41%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	313	347	391	389	404	432	445	419	444	455	444
Alcohol-related disorders	270	273	270	284	295	300	307	288	302	311	295
Anxiety disorders	1,577	1,692	1,830	1,924	1,974	2,037	2,143	1,993	2,112	2,115	2,034
Attention-deficit, conduct, and disruptive behavior disorders	272	313	330	344	370	412	463	471	520	568	569
Delirium, dementia, and amnestic and other cognitive disorders	801	837	811	766	729	710	718	657	644	592	551
Developmental disorders	404	425	437	424	432	427	464	427	408	407	390
Disorders usually diagnosed in infancy, childhood, adolescence	37	41	45	49	50	60	56	59	65	69	67
Impulse control disorders, NEC	85	90	87	102	106	117	147	134	146	158	158
Miscellaneous disorders	676	721	773	793	809	826	875	797	859	860	834
Mood disorders	1,720	1,857	2,012	2,114	2,183	2,275	2,352	2,221	2,368	2,376	2,316
Personality disorders	214	230	245	254	263	270	262	254	257	250	230
Schizophrenia and other psychotic disorders	804	860	865	861	845	845	875	828	814	776	720
Screening and history of mental health and substance abuse	1,544	1,696	1,847	1,953	2,036	2,143	2,258	2,040	2,216	2,244	2,122
Substance-related disorders	394	424	471	481	498	540	576	539	602	656	610
Suicide and intentional self-inflicted injury	152	167	178	183	186	213	228	224	254	266	248



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Jackson County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,112	1,214	1,259	1,306	1,329	1,364	1,406	1,327	1,374	1,376	1,273
Female	553	596	613	631	632	646	646	594	604	619	551
Male	559	618	646	675	697	718	760	733	770	757	722
White	1,050	1,153	1,189	1,229	1,248	1,278	1,310	1,230	1,275	1,278	1,182
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	56%	58%	60%	61%	63%	62%	60%	61%	59%	56%
Female	56%	59%	61%	62%	62%	63%	60%	58%	59%	57%	52%
Male	51%	54%	56%	58%	60%	63%	63%	62%	63%	60%	58%
White	52%	55%	57%	59%	60%	62%	60%	58%	60%	58%	54%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	234	258	267	265	257	257	253	222	229	218	195
Alcohol-related disorders	83	87	89	83	78	69	64	51	48	40	33
Anxiety disorders	559	606	637	668	679	682	687	660	667	668	624
Attention-deficit, conduct, and disruptive behavior disorders	321	359	386	398	419	430	456	432	453	440	407
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	269	330	374	411	458	488	532	505	552	559	539
Disorders usually diagnosed in infancy, childhood, adolescence	64	70	78	77	84	92	98	97	95	96	85
Impulse control disorders, NEC	38	38	41	38	37	36	36	34	33	N.D.	N.D.
Miscellaneous disorders	202	218	216	223	213	210	206	187	190	189	168
Mood disorders	394	377	366	365	341	314	299	260	236	200	166
Personality disorders	76	80	78	75	73	71	70	56	54	46	37
Schizophrenia and other psychotic disorders	31	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	583	601	589	590	575	572	559	514	508	486	422
Substance-related disorders	184	176	174	162	157	142	128	106	105	95	78
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Jackson County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,536	1,662	1,710	1,748	1,805	1,859	1,967	1,840	1,963	2,007	1,981
Female	1,029	1,103	1,120	1,156	1,185	1,201	1,287	1,184	1,281	1,308	1,282
Male	507	559	590	592	620	658	680	656	682	699	699
White	1,393	1,512	1,565	1,598	1,658	1,712	1,808	1,687	1,809	1,848	1,826
Minority	35	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	69%	73%	74%	74%	74%	75%	76%	71%	74%	73%	70%
Female	74%	78%	78%	78%	78%	79%	81%	74%	78%	78%	75%
Male	61%	65%	68%	67%	68%	68%	70%	66%	67%	65%	63%
White	69%	73%	74%	73%	74%	74%	76%	70%	73%	72%	69%
Minority	18%	17%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	164	180	188	214	216	221	250	244	237	248	245
Alcohol-related disorders	171	191	195	212	212	228	231	219	225	222	219
Anxiety disorders	1,014	1,108	1,171	1,228	1,275	1,332	1,423	1,329	1,391	1,416	1,396
Attention-deficit, conduct, and disruptive behavior disorders	46	56	60	71	79	88	95	84	93	103	110
Delirium, dementia, and amnesic and other cognitive disorders	300	309	292	282	278	264	266	234	235	214	194
Developmental disorders	130	134	140	156	162	171	173	167	174	170	177
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	438	470	505	529	553	574	617	594	628	637	621
Mood disorders	985	1,073	1,115	1,165	1,184	1,230	1,294	1,216	1,280	1,307	1,279
Personality disorders	97	110	113	128	136	142	141	129	136	136	140
Schizophrenia and other psychotic disorders	270	277	282	279	273	274	276	251	267	253	238
Screening and history of mental health and substance abuse	861	939	1,005	1,041	1,089	1,141	1,225	1,154	1,246	1,284	1,263
Substance-related disorders	263	293	307	350	369	393	435	415	448	469	458
Suicide and intentional self-inflicted injury	39	41	41	45	44	49	59	64	64	62	67



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Jefferson County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	27,005	29,923	32,167	33,983	35,145	35,410	35,746	33,926	34,849	35,138	33,399
Female	12,855	14,031	14,990	15,689	16,109	16,033	16,048	15,164	15,477	15,462	14,503
Male	14,150	15,892	17,177	18,294	19,036	19,377	19,698	18,762	19,372	19,676	18,896
White	11,662	13,218	14,222	15,210	15,817	16,057	16,241	15,424	15,760	16,025	15,360
Minority	13,733	14,833	15,768	16,352	16,633	16,494	16,432	15,368	15,779	15,645	14,690

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	48%	50%	52%	52%	52%	50%	49%	46%	45%	43%	39%
Female	47%	48%	50%	49%	48%	47%	45%	42%	41%	39%	35%
Male	50%	52%	54%	55%	55%	54%	53%	49%	49%	47%	43%
White	48%	50%	53%	53%	52%	50%	48%	44%	43%	41%	37%
Minority	43%	44%	45%	45%	45%	43%	43%	39%	39%	36%	33%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	8,198	9,197	9,909	10,344	10,588	10,544	10,425	9,548	9,393	8,995	8,275
Alcohol-related disorders	1,056	1,077	1,056	1,012	980	914	849	730	655	574	471
Anxiety disorders	7,205	7,820	8,178	8,443	8,580	8,386	8,222	7,634	7,463	7,170	6,436
Attention-deficit, conduct, and disruptive behavior disorders	13,367	15,095	16,483	17,571	18,326	18,626	18,732	17,608	17,642	17,401	16,345
Delirium, dementia, and amnestic and other cognitive disorders	488	524	549	571	565	554	548	514	519	493	464
Developmental disorders	5,663	6,764	7,718	8,679	9,434	9,918	10,479	10,575	11,173	11,658	11,531
Disorders usually diagnosed in infancy, childhood, adolescence	2,347	2,729	3,012	3,316	3,553	3,730	3,928	3,843	3,978	3,953	3,837
Impulse control disorders, NEC	2,520	2,747	2,876	2,932	2,918	2,810	2,715	2,445	2,332	2,157	1,852
Miscellaneous disorders	2,919	3,049	3,172	3,258	3,278	3,269	3,195	2,962	2,947	2,875	2,660
Mood disorders	9,844	10,536	10,788	10,875	10,820	10,413	10,055	9,066	8,715	8,173	7,203
Personality disorders	791	836	868	878	876	842	810	744	733	667	563
Schizophrenia and other psychotic disorders	1,551	1,636	1,662	1,700	1,716	1,676	1,649	1,547	1,547	1,437	1,250
Screening and history of mental health and substance abuse	16,460	17,732	18,518	18,794	18,684	17,990	17,513	16,008	15,772	15,100	13,668
Substance-related disorders	2,936	3,042	3,010	2,933	2,851	2,700	2,515	2,229	2,166	1,985	1,679
Suicide and intentional self-inflicted injury	923	954	1,002	1,043	1,050	1,023	1,010	951	959	903	792



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Jefferson County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	22,608	24,718	26,599	28,305	29,831	30,885	31,491	30,153	32,964	34,141	33,348
Female	17,078	18,540	19,897	21,176	22,260	22,811	23,167	21,904	23,895	24,432	23,632
Male	5,530	6,178	6,702	7,129	7,571	8,074	8,324	8,249	9,069	9,709	9,716
White	12,720	13,911	14,902	15,855	16,725	17,440	17,892	16,900	18,435	18,886	18,153
Minority	8,884	9,698	10,491	11,150	11,749	12,030	12,093	11,704	12,866	13,498	13,243

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	58%	60%	62%	62%	63%	63%	59%	62%	62%	58%
Female	56%	59%	61%	63%	63%	64%	64%	59%	63%	62%	59%
Male	51%	55%	57%	59%	59%	61%	61%	59%	61%	60%	57%
White	57%	60%	62%	65%	65%	66%	65%	61%	65%	64%	60%
Minority	46%	49%	51%	52%	53%	53%	53%	50%	53%	52%	49%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	2,739	3,040	3,278	3,473	3,748	3,913	4,102	4,023	4,454	4,709	4,714
Alcohol-related disorders	2,667	2,999	3,254	3,484	3,690	3,856	3,974	3,873	4,116	4,177	4,006
Anxiety disorders	9,081	10,109	10,995	11,860	12,560	13,135	13,752	13,570	14,612	15,161	14,772
Attention-deficit, conduct, and disruptive behavior disorders	1,521	1,749	1,997	2,222	2,472	2,777	3,145	3,358	3,914	4,248	4,349
Delirium, dementia, and amnestic and other cognitive disorders	5,914	6,225	6,383	6,358	6,262	6,350	6,389	6,126	6,171	5,936	5,423
Developmental disorders	2,021	2,171	2,320	2,427	2,495	2,576	2,674	2,713	2,897	3,056	3,129
Disorders usually diagnosed in infancy, childhood, adolescence	290	328	360	405	449	495	564	613	696	801	878
Impulse control disorders, NEC	523	580	660	729	808	871	978	1,024	1,173	1,295	1,385
Miscellaneous disorders	3,362	3,843	4,323	4,942	5,513	5,899	6,121	5,641	6,233	6,302	5,940
Mood disorders	12,322	13,531	14,690	15,769	16,710	17,361	17,967	17,508	18,937	19,587	19,281
Personality disorders	1,333	1,476	1,586	1,618	1,686	1,760	1,854	1,817	1,904	1,944	1,878
Schizophrenia and other psychotic disorders	5,753	6,175	6,440	6,599	6,764	6,937	7,033	6,797	6,949	6,932	6,608
Screening and history of mental health and substance abuse	12,370	13,989	15,544	16,957	18,433	19,202	19,608	18,776	20,742	21,489	20,910
Substance-related disorders	3,885	4,436	4,877	5,380	5,783	6,066	6,399	6,229	6,818	7,130	6,889
Suicide and intentional self-inflicted injury	859	975	1,105	1,222	1,326	1,446	1,595	1,599	1,858	1,949	1,950



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Jessamine County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,707	1,911	2,041	2,161	2,244	2,282	2,429	2,333	2,522	2,579	2,448
Female	838	928	974	1,025	1,087	1,082	1,168	1,089	1,183	1,160	1,096
Male	869	983	1,067	1,136	1,157	1,200	1,261	1,244	1,339	1,419	1,352
White	1,501	1,680	1,797	1,895	1,987	2,004	2,099	1,991	2,149	2,204	2,091
Minority	111	123	133	144	133	144	159	164	188	188	181

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	54%	55%	55%	55%	53%	54%	50%	50%	47%	43%
Female	52%	53%	53%	53%	53%	51%	52%	47%	49%	44%	39%
Male	52%	54%	56%	57%	56%	55%	56%	52%	52%	51%	46%
White	51%	53%	54%	54%	54%	53%	52%	48%	48%	46%	42%
Minority	33%	34%	32%	33%	30%	30%	31%	31%	33%	27%	26%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	543	612	667	704	737	714	747	700	711	690	626
Alcohol-related disorders	72	68	70	63	59	56	65	48	51	42	37
Anxiety disorders	543	579	615	658	678	672	739	707	734	719	671
Attention-deficit, conduct, and disruptive behavior disorders	834	948	1,015	1,064	1,120	1,120	1,194	1,163	1,183	1,174	1,100
Delirium, dementia, and amnestic and other cognitive disorders	42	45	46	42	50	55	66	67	65	60	56
Developmental disorders	412	475	536	577	607	652	727	722	788	847	835
Disorders usually diagnosed in infancy, childhood, adolescence	142	165	181	211	227	241	264	256	259	260	252
Impulse control disorders, NEC	55	61	67	70	72	62	68	61	53	48	37
Miscellaneous disorders	245	273	288	278	298	286	297	284	291	279	266
Mood disorders	551	596	609	606	604	569	600	536	537	523	454
Personality disorders	52	57	60	49	50	50	49	36	37	34	32
Schizophrenia and other psychotic disorders	81	87	89	99	100	100	104	94	90	86	76
Screening and history of mental health and substance abuse	1,031	1,121	1,167	1,190	1,158	1,147	1,137	1,045	1,108	1,064	940
Substance-related disorders	219	219	221	204	199	181	186	154	162	141	105
Suicide and intentional self-inflicted injury	63	64	66	66	62	66	65	56	59	62	57



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Jessamine County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,342	1,517	1,633	1,756	1,872	1,906	2,040	1,916	2,173	2,317	2,223
Female	1,017	1,156	1,254	1,343	1,425	1,437	1,527	1,424	1,630	1,718	1,603
Male	325	361	379	413	447	469	513	492	543	599	620
White	1,198	1,363	1,475	1,593	1,701	1,730	1,855	1,738	1,968	2,085	1,985
Minority	91	100	101	103	106	107	102	93	105	132	132

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	68%	67%	68%	70%	70%	71%	65%	70%	71%	66%
Female	66%	70%	69%	70%	72%	70%	71%	64%	71%	72%	66%
Male	60%	63%	61%	64%	65%	67%	70%	65%	69%	66%	64%
White	65%	69%	68%	69%	71%	70%	71%	64%	70%	71%	66%
Minority	39%	41%	38%	38%	40%	40%	39%	36%	38%	38%	36%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	186	206	243	272	286	305	322	299	344	374	355
Alcohol-related disorders	128	154	173	197	209	210	229	214	227	235	228
Anxiety disorders	795	922	1,013	1,117	1,199	1,221	1,337	1,275	1,411	1,476	1,434
Attention-deficit, conduct, and disruptive behavior disorders	114	138	148	175	190	227	259	256	307	347	357
Delirium, dementia, and amnestic and other cognitive disorders	326	349	348	343	336	334	341	307	305	302	278
Developmental disorders	135	152	163	174	182	189	202	212	234	259	259
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	31	31	33	40	42	54	63	60
Impulse control disorders, NEC	N.D.	N.D.	32	35	36	41	42	45	48	51	52
Miscellaneous disorders	322	368	417	462	508	528	584	525	602	644	608
Mood disorders	826	951	1,054	1,139	1,217	1,248	1,355	1,267	1,408	1,485	1,455
Personality disorders	109	131	149	158	179	172	184	180	187	201	190
Schizophrenia and other psychotic disorders	311	335	354	354	361	362	380	349	363	357	329
Screening and history of mental health and substance abuse	760	885	993	1,108	1,195	1,225	1,342	1,258	1,428	1,544	1,502
Substance-related disorders	309	366	419	468	509	518	591	565	652	698	660
Suicide and intentional self-inflicted injury	62	72	88	96	110	104	120	120	133	144	139



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Johnson County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,253	2,354	2,449	2,596	2,606	2,626	2,670	2,508	2,553	2,525	2,364
Female	1,150	1,179	1,188	1,243	1,249	1,243	1,240	1,183	1,174	1,148	1,080
Male	1,103	1,175	1,261	1,353	1,357	1,383	1,430	1,325	1,379	1,377	1,284
White	2,095	2,186	2,266	2,389	2,385	2,401	2,417	2,278	2,308	2,284	2,132
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	59%	63%	64%	63%	64%	65%	62%	63%	63%	59%
Female	60%	61%	63%	63%	63%	62%	64%	61%	61%	60%	57%
Male	56%	58%	63%	64%	64%	65%	66%	63%	65%	66%	61%
White	57%	58%	61%	62%	62%	62%	63%	60%	61%	62%	57%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	331	346	359	364	373	377	364	326	314	312	283
Alcohol-related disorders	86	82	79	71	60	52	45	36	N.D.	N.D.	N.D.
Anxiety disorders	827	868	860	909	919	928	961	901	917	889	835
Attention-deficit, conduct, and disruptive behavior disorders	820	876	976	1,049	1,073	1,109	1,137	1,076	1,083	1,050	977
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	568	623	694	757	777	819	865	821	858	840	802
Disorders usually diagnosed in infancy, childhood, adolescence	133	149	165	186	200	211	223	220	221	219	210
Impulse control disorders, NEC	52	50	55	49	44	43	44	40	N.D.	N.D.	N.D.
Miscellaneous disorders	357	355	362	380	401	423	459	438	446	420	386
Mood disorders	582	570	552	545	519	488	459	388	356	321	264
Personality disorders	37	40	34	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	64	58	62	62	53	53	58	45	43	43	35
Screening and history of mental health and substance abuse	1,516	1,564	1,624	1,728	1,681	1,622	1,576	1,435	1,455	1,422	1,307
Substance-related disorders	241	231	206	191	179	151	139	110	110	100	79
Suicide and intentional self-inflicted injury	60	55	59	60	53	46	42	34	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Johnson County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,524	2,617	2,714	2,819	2,887	2,950	3,140	2,901	3,156	3,164	3,079
Female	1,694	1,761	1,800	1,872	1,901	1,950	2,070	1,901	2,071	2,044	1,968
Male	830	856	914	947	986	1,000	1,070	1,000	1,085	1,120	1,111
White	2,345	2,433	2,520	2,621	2,683	2,742	2,915	2,673	2,925	2,942	2,867
Minority	38	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	69%	72%	72%	72%	72%	75%	70%	74%	72%	69%
Female	72%	73%	75%	76%	75%	75%	79%	74%	77%	76%	73%
Male	62%	63%	66%	66%	67%	66%	69%	63%	67%	66%	64%
White	68%	69%	71%	72%	71%	72%	75%	69%	73%	72%	70%
Minority	16%	14%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	219	226	238	269	277	291	325	322	344	353	341
Alcohol-related disorders	215	229	231	245	263	260	296	261	273	268	255
Anxiety disorders	1,713	1,793	1,863	1,927	1,980	2,046	2,183	2,051	2,161	2,150	2,076
Attention-deficit, conduct, and disruptive behavior disorders	97	100	107	119	124	150	170	160	204	234	255
Delirium, dementia, and amnestic and other cognitive disorders	427	427	418	404	402	403	420	379	389	354	328
Developmental disorders	178	179	181	190	195	202	224	221	233	254	253
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	31	31	32	36	41	37
Impulse control disorders, NEC	39	42	41	46	52	55	63	59	63	67	68
Miscellaneous disorders	690	739	760	820	839	867	917	844	892	875	809
Mood disorders	1,375	1,462	1,519	1,585	1,638	1,678	1,803	1,671	1,808	1,784	1,730
Personality disorders	155	163	169	180	178	187	199	188	201	193	193
Schizophrenia and other psychotic disorders	431	442	437	435	445	428	455	425	429	416	399
Screening and history of mental health and substance abuse	1,678	1,766	1,881	1,988	2,056	2,139	2,280	2,090	2,258	2,295	2,243
Substance-related disorders	495	535	540	587	617	655	715	661	726	736	719
Suicide and intentional self-inflicted injury	93	96	98	109	116	129	143	137	142	145	145



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Kenton County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,505	6,191	6,634	7,040	7,162	7,429	7,800	7,436	8,019	8,225	7,802
Female	2,652	2,966	3,156	3,299	3,298	3,403	3,552	3,352	3,641	3,739	3,493
Male	2,853	3,225	3,478	3,741	3,864	4,026	4,248	4,084	4,378	4,486	4,309
White	4,320	4,872	5,198	5,513	5,588	5,845	6,093	5,801	6,266	6,436	6,111
Minority	941	1,036	1,106	1,141	1,139	1,131	1,174	1,133	1,205	1,241	1,175

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	58%	60%	59%	58%	58%	59%	54%	55%	52%	46%
Female	56%	58%	59%	57%	56%	55%	55%	50%	52%	49%	44%
Male	57%	59%	61%	62%	60%	60%	62%	58%	59%	55%	49%
White	57%	59%	60%	60%	59%	59%	59%	54%	56%	52%	46%
Minority	45%	45%	44%	43%	40%	38%	39%	36%	37%	36%	32%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	1,283	1,410	1,506	1,604	1,608	1,626	1,657	1,529	1,554	1,518	1,373
Alcohol-related disorders	287	306	307	301	305	288	283	257	248	235	200
Anxiety disorders	1,666	1,813	1,870	1,895	1,860	1,880	1,892	1,730	1,760	1,713	1,563
Attention-deficit, conduct, and disruptive behavior disorders	3,117	3,507	3,873	4,217	4,390	4,601	4,875	4,656	4,869	4,890	4,646
Delirium, dementia, and amnestic and other cognitive disorders	126	121	139	147	151	151	161	160	169	153	137
Developmental disorders	1,449	1,676	1,887	2,116	2,310	2,487	2,725	2,661	2,979	3,058	2,947
Disorders usually diagnosed in infancy, childhood, adolescence	704	807	928	1,010	1,074	1,130	1,193	1,128	1,183	1,167	1,103
Impulse control disorders, NEC	319	340	357	355	361	361	347	316	297	286	252
Miscellaneous disorders	1,137	1,258	1,338	1,410	1,444	1,462	1,508	1,379	1,454	1,421	1,328
Mood disorders	2,076	2,256	2,309	2,373	2,303	2,268	2,267	2,030	1,995	1,936	1,719
Personality disorders	354	354	348	331	314	285	257	217	192	166	144
Schizophrenia and other psychotic disorders	268	272	274	293	280	286	292	263	268	275	250
Screening and history of mental health and substance abuse	3,789	4,201	4,384	4,465	4,359	4,338	4,381	4,056	4,173	4,136	3,732
Substance-related disorders	653	680	683	663	659	623	604	533	514	485	413
Suicide and intentional self-inflicted injury	240	240	240	259	264	270	276	278	296	285	245



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Kenton County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,722	5,275	5,698	6,005	6,343	6,545	6,952	6,347	7,147	7,368	6,983
Female	3,617	3,999	4,330	4,533	4,738	4,925	5,203	4,699	5,296	5,417	5,072
Male	1,105	1,276	1,368	1,472	1,605	1,620	1,749	1,648	1,851	1,951	1,911
White	4,032	4,484	4,815	5,103	5,423	5,595	5,946	5,432	6,115	6,296	5,940
Minority	515	609	686	699	704	716	747	670	773	796	739

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	69%	69%	70%	70%	73%	65%	71%	70%	65%
Female	66%	69%	71%	71%	72%	72%	74%	66%	71%	71%	66%
Male	58%	61%	63%	64%	67%	66%	69%	64%	69%	67%	63%
White	65%	68%	69%	70%	72%	72%	74%	66%	72%	72%	66%
Minority	44%	48%	51%	50%	48%	48%	49%	44%	48%	46%	41%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	654	732	831	862	908	950	1,008	957	1,016	1,081	1,042
Alcohol-related disorders	706	784	830	868	931	975	1,021	943	1,022	1,026	964
Anxiety disorders	2,592	2,882	3,143	3,356	3,542	3,681	3,963	3,714	4,043	4,185	3,993
Attention-deficit, conduct, and disruptive behavior disorders	360	424	482	596	698	782	895	898	1,053	1,151	1,118
Delirium, dementia, and amnestic and other cognitive disorders	1,004	1,136	1,180	1,213	1,226	1,253	1,258	1,197	1,227	1,153	1,049
Developmental disorders	367	406	445	455	468	484	545	569	620	640	659
Disorders usually diagnosed in infancy, childhood, adolescence	60	75	82	94	100	113	137	134	154	185	197
Impulse control disorders, NEC	72	88	100	112	122	138	149	145	170	180	180
Miscellaneous disorders	1,130	1,274	1,457	1,642	1,770	1,835	1,962	1,749	1,914	1,946	1,775
Mood disorders	2,910	3,252	3,528	3,833	4,076	4,245	4,571	4,265	4,720	4,883	4,611
Personality disorders	561	638	688	762	834	880	1,018	909	956	921	823
Schizophrenia and other psychotic disorders	1,028	1,151	1,194	1,245	1,290	1,356	1,415	1,360	1,435	1,416	1,332
Screening and history of mental health and substance abuse	3,031	3,433	3,822	4,100	4,346	4,511	4,880	4,434	5,054	5,256	4,963
Substance-related disorders	936	1,074	1,156	1,278	1,370	1,494	1,603	1,474	1,652	1,751	1,674
Suicide and intentional self-inflicted injury	280	310	350	378	398	440	526	501	554	613	576



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Knott County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,659	1,711	1,731	1,751	1,729	1,741	1,753	1,620	1,705	1,681	1,575
Female	815	822	830	828	815	817	822	745	784	760	721
Male	844	889	901	923	914	924	931	875	921	921	854
White	1,560	1,613	1,628	1,647	1,617	1,619	1,617	1,493	1,571	1,547	1,444
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	58%	61%	62%	62%	63%	65%	61%	65%	62%	57%
Female	59%	59%	61%	61%	61%	62%	64%	59%	62%	59%	53%
Male	54%	57%	61%	63%	63%	65%	67%	63%	67%	65%	59%
White	55%	57%	60%	61%	61%	62%	64%	60%	64%	61%	55%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	484	511	518	539	525	523	505	473	473	446	413
Alcohol-related disorders	96	94	84	86	75	64	59	51	45	38	32
Anxiety disorders	570	575	560	561	549	547	562	515	529	502	464
Attention-deficit, conduct, and disruptive behavior disorders	528	559	581	596	601	601	605	545	540	527	488
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	454	505	552	616	661	700	738	714	761	758	738
Disorders usually diagnosed in infancy, childhood, adolescence	80	88	89	98	99	104	109	100	106	103	103
Impulse control disorders, NEC	52	55	53	52	51	48	44	38	N.D.	N.D.	N.D.
Miscellaneous disorders	268	267	268	267	253	254	260	236	228	228	205
Mood disorders	515	490	461	431	385	352	333	289	279	252	215
Personality disorders	89	85	79	83	84	75	71	60	51	46	39
Schizophrenia and other psychotic disorders	60	56	51	52	50	45	45	37	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	1,144	1,161	1,140	1,128	1,089	1,023	990	871	870	825	742
Substance-related disorders	259	248	216	202	181	154	153	126	118	114	99
Suicide and intentional self-inflicted injury	55	48	41	42	37	32	33	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Knott County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,290	2,399	2,442	2,512	2,564	2,589	2,596	2,377	2,513	2,508	2,400
Female	1,440	1,516	1,533	1,559	1,586	1,621	1,621	1,465	1,547	1,566	1,475
Male	850	883	909	953	978	968	975	912	966	942	925
White	2,121	2,227	2,259	2,336	2,378	2,413	2,408	2,198	2,336	2,337	2,226
Minority	42	35	39	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	70%	70%	72%	73%	74%	76%	70%	75%	75%	72%
Female	72%	74%	74%	75%	76%	78%	79%	73%	79%	80%	75%
Male	61%	64%	65%	67%	68%	68%	71%	66%	70%	68%	67%
White	67%	69%	70%	71%	72%	74%	76%	69%	75%	75%	71%
Minority	18%	15%	17%	14%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	264	283	294	325	344	353	375	344	389	384	373
Alcohol-related disorders	276	287	311	316	326	328	328	311	317	314	305
Anxiety disorders	1,514	1,599	1,630	1,728	1,748	1,773	1,827	1,709	1,780	1,761	1,696
Attention-deficit, conduct, and disruptive behavior disorders	59	61	64	85	95	118	127	124	150	159	169
Delirium, dementia, and amnestic and other cognitive disorders	357	362	358	349	336	335	329	292	287	261	242
Developmental disorders	214	213	211	216	228	230	241	229	243	249	251
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	32
Impulse control disorders, NEC	49	52	56	60	63	61	68	65	77	74	74
Miscellaneous disorders	687	751	772	811	859	872	868	799	814	805	758
Mood disorders	1,342	1,419	1,452	1,529	1,558	1,595	1,609	1,503	1,589	1,563	1,511
Personality disorders	173	183	185	208	223	225	226	203	220	208	209
Schizophrenia and other psychotic disorders	576	593	597	612	601	595	595	555	543	524	507
Screening and history of mental health and substance abuse	1,478	1,584	1,677	1,747	1,821	1,850	1,874	1,725	1,867	1,853	1,794
Substance-related disorders	596	651	688	732	764	777	817	760	837	835	801
Suicide and intentional self-inflicted injury	90	91	94	103	107	119	129	124	133	128	131



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Knox County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,418	3,662	3,798	4,010	4,112	4,124	4,210	3,929	4,054	4,021	3,762
Female	1,723	1,831	1,876	1,970	1,989	1,962	2,003	1,864	1,936	1,910	1,783
Male	1,695	1,831	1,922	2,040	2,123	2,162	2,207	2,065	2,118	2,111	1,979
White	3,174	3,398	3,521	3,705	3,788	3,792	3,847	3,586	3,708	3,682	3,443
Minority	50	57	53	53	57	58	64	62	67	60	56

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	59%	61%	63%	65%	65%	66%	66%	62%	62%	60%	56%
Female	61%	63%	64%	65%	66%	65%	65%	60%	61%	58%	54%
Male	57%	60%	62%	64%	65%	66%	67%	63%	63%	62%	57%
White	57%	60%	62%	64%	64%	65%	64%	60%	60%	59%	55%
Minority	17%	18%	16%	14%	14%	14%	16%	15%	16%	13%	13%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	643	671	707	736	747	731	713	638	640	607	536
Alcohol-related disorders	303	302	301	297	297	265	236	195	186	159	133
Anxiety disorders	1,748	1,877	1,987	2,123	2,227	2,241	2,341	2,208	2,260	2,244	2,101
Attention-deficit, conduct, and disruptive behavior disorders	1,300	1,375	1,485	1,567	1,594	1,581	1,566	1,418	1,425	1,363	1,258
Delirium, dementia, and amnesic and other cognitive disorders	39	40	38	36	33	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	859	952	1,007	1,085	1,157	1,170	1,192	1,090	1,108	1,066	985
Disorders usually diagnosed in infancy, childhood, adolescence	231	252	270	292	321	330	338	306	318	307	286
Impulse control disorders, NEC	171	176	167	165	173	165	169	158	154	140	122
Miscellaneous disorders	637	645	622	636	614	598	598	543	536	514	464
Mood disorders	1,107	1,103	1,066	1,027	969	896	840	723	665	599	509
Personality disorders	199	198	198	202	202	188	177	153	143	119	110
Schizophrenia and other psychotic disorders	135	133	127	126	124	118	114	94	96	80	71
Screening and history of mental health and substance abuse	1,739	1,801	1,811	1,819	1,794	1,710	1,672	1,518	1,522	1,449	1,310
Substance-related disorders	593	579	564	554	546	465	420	361	340	297	246
Suicide and intentional self-inflicted injury	109	109	109	100	90	84	84	78	72	58	51



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Knox County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,270	4,510	4,762	4,862	4,918	5,011	5,206	5,006	5,324	5,348	5,240
Female	2,840	2,973	3,132	3,199	3,220	3,261	3,385	3,226	3,451	3,440	3,333
Male	1,430	1,537	1,630	1,663	1,698	1,750	1,821	1,780	1,873	1,908	1,907
White	3,907	4,137	4,370	4,464	4,533	4,629	4,808	4,620	4,931	4,948	4,840
Minority	87	81	80	79	64	53	47	44	42	49	47

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	70%	71%	72%	73%	72%	73%	75%	71%	73%	73%	70%
Female	73%	75%	76%	76%	75%	77%	79%	74%	76%	77%	73%
Male	64%	64%	66%	67%	67%	68%	69%	66%	67%	68%	66%
White	69%	70%	71%	72%	71%	73%	74%	70%	72%	73%	70%
Minority	19%	17%	17%	16%	14%	11%	10%	9%	9%	10%	9%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	422	448	484	489	516	544	580	567	623	646	631
Alcohol-related disorders	476	501	527	557	573	592	626	584	620	617	585
Anxiety disorders	2,931	3,111	3,305	3,404	3,470	3,567	3,725	3,619	3,845	3,859	3,820
Attention-deficit, conduct, and disruptive behavior disorders	130	140	158	182	218	252	279	306	371	388	378
Delirium, dementia, and amnesic and other cognitive disorders	811	819	816	799	783	770	756	687	642	596	544
Developmental disorders	299	315	329	345	343	363	396	400	416	441	448
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	31	37	37	34	42	48	51	57	65	72
Impulse control disorders, NEC	68	75	77	80	83	93	100	96	107	112	111
Miscellaneous disorders	1,318	1,433	1,523	1,608	1,676	1,735	1,810	1,761	1,868	1,840	1,771
Mood disorders	2,321	2,472	2,628	2,728	2,790	2,859	3,034	2,894	3,095	3,081	3,015
Personality disorders	290	303	326	342	353	363	381	371	402	404	387
Schizophrenia and other psychotic disorders	814	836	849	866	851	872	894	854	839	803	752
Screening and history of mental health and substance abuse	2,268	2,454	2,659	2,781	2,858	2,959	3,144	3,052	3,292	3,362	3,289
Substance-related disorders	932	1,017	1,109	1,157	1,211	1,257	1,343	1,291	1,383	1,422	1,382
Suicide and intentional self-inflicted injury	122	132	145	158	175	189	214	213	232	229	222



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Larue County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	625	694	775	862	894	881	920	890	909	910	881
Female	305	338	366	406	402	395	425	406	413	407	393
Male	320	356	409	456	492	486	495	484	496	503	488
White	562	621	691	766	788	773	800	773	793	797	769
Minority	43	45	51	60	66	65	70	65	60	55	54

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	51%	52%	56%	58%	59%	57%	58%	54%	53%	49%	46%
Female	52%	53%	54%	56%	55%	53%	54%	49%	48%	44%	42%
Male	51%	52%	59%	61%	62%	61%	61%	59%	58%	54%	50%
White	52%	51%	56%	58%	57%	56%	56%	52%	52%	48%	45%
Minority	35%	37%	38%	40%	44%	40%	42%	40%	33%	28%	28%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	131	146	176	187	196	187	198	188	185	190	179
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	188	218	229	261	275	284	288	287	282	268	253
Attention-deficit, conduct, and disruptive behavior disorders	296	327	380	422	457	448	463	464	456	436	419
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	175	186	216	256	277	289	295	300	306	316	304
Disorders usually diagnosed in infancy, childhood, adolescence	41	42	47	55	57	62	66	70	66	68	65
Impulse control disorders, NEC	47	53	65	66	63	63	70	63	56	47	39
Miscellaneous disorders	71	77	82	92	91	91	94	83	83	71	60
Mood disorders	226	252	262	279	268	256	261	242	232	207	179
Personality disorders	32	42	40	42	41	38	40	32	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	404	445	483	513	497	468	478	426	415	384	350
Substance-related disorders	45	51	56	55	56	53	52	37	36	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Larue County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	614	666	693	744	766	795	838	819	886	926	866
Female	433	480	501	536	556	567	595	588	626	643	611
Male	181	186	192	208	210	228	243	231	260	283	255
White	541	585	606	657	682	708	745	722	778	827	770
Minority	49	53	58	59	54	55	55	53	60	55	50

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	58%	60%	62%	62%	64%	65%	62%	65%	65%	61%
Female	57%	61%	63%	64%	64%	66%	66%	63%	66%	66%	63%
Male	51%	53%	54%	56%	58%	59%	61%	59%	61%	63%	57%
White	55%	58%	59%	62%	63%	64%	65%	61%	63%	65%	60%
Minority	37%	40%	44%	40%	38%	40%	40%	41%	42%	40%	35%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	50	53	58	65	71	68	82	79	90	97	92
Alcohol-related disorders	43	41	49	53	57	53	56	54	59	63	62
Anxiety disorders	292	316	339	370	388	415	437	428	451	478	435
Attention-deficit, conduct, and disruptive behavior disorders	31	N.D.	31	40	44	56	75	71	90	107	98
Delirium, dementia, and amnestic and other cognitive disorders	175	184	176	182	177	163	178	178	166	165	143
Developmental disorders	59	58	63	62	63	61	64	67	74	78	80
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33	36	37
Miscellaneous disorders	107	116	126	147	152	164	178	170	182	185	170
Mood disorders	358	394	414	457	457	491	529	534	573	595	555
Personality disorders	61	66	75	77	77	78	91	88	91	90	86
Schizophrenia and other psychotic disorders	141	154	149	156	154	154	152	152	152	143	122
Screening and history of mental health and substance abuse	267	297	326	364	388	412	423	385	439	456	439
Substance-related disorders	61	64	66	76	79	88	102	93	99	113	103
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	33	34	32



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Laurel County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,300	4,792	5,060	5,269	5,473	5,572	5,621	5,269	5,521	5,471	5,104
Female	2,195	2,420	2,549	2,613	2,703	2,711	2,707	2,480	2,621	2,562	2,404
Male	2,105	2,372	2,511	2,656	2,770	2,861	2,914	2,789	2,900	2,909	2,700
White	4,032	4,482	4,731	4,917	5,115	5,186	5,202	4,870	5,105	5,037	4,700
Minority	72	83	81	89	84	79	82	83	84	92	82

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	61%	64%	64%	65%	65%	65%	61%	61%	59%	54%
Female	60%	62%	64%	64%	65%	65%	65%	59%	60%	57%	53%
Male	57%	61%	63%	64%	64%	65%	65%	62%	62%	61%	56%
White	57%	60%	63%	63%	64%	64%	64%	60%	60%	58%	53%
Minority	21%	23%	21%	21%	19%	17%	17%	17%	17%	16%	15%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	841	914	959	996	1,037	1,022	988	874	853	836	747
Alcohol-related disorders	305	324	337	331	335	313	289	245	226	203	168
Anxiety disorders	2,227	2,485	2,662	2,784	2,954	3,051	3,056	2,871	2,952	2,884	2,736
Attention-deficit, conduct, and disruptive behavior disorders	1,807	2,019	2,122	2,220	2,317	2,378	2,348	2,194	2,221	2,163	1,992
Delirium, dementia, and amnestic and other cognitive disorders	79	78	70	66	60	59	57	47	54	47	39
Developmental disorders	963	1,126	1,256	1,381	1,528	1,623	1,649	1,536	1,617	1,643	1,558
Disorders usually diagnosed in infancy, childhood, adolescence	242	299	322	356	386	402	416	385	388	383	369
Impulse control disorders, NEC	187	191	193	193	190	179	169	147	144	134	110
Miscellaneous disorders	621	669	669	667	679	656	636	557	573	571	512
Mood disorders	1,379	1,413	1,416	1,360	1,320	1,267	1,177	1,013	981	892	747
Personality disorders	212	220	221	225	226	213	194	170	150	139	113
Schizophrenia and other psychotic disorders	151	151	150	158	151	148	133	125	125	114	94
Screening and history of mental health and substance abuse	2,326	2,533	2,581	2,593	2,552	2,462	2,429	2,177	2,241	2,179	1,903
Substance-related disorders	611	625	612	604	597	551	498	425	396	374	295
Suicide and intentional self-inflicted injury	140	147	147	146	134	127	117	109	108	99	83



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Laurel County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,406	4,770	5,031	5,258	5,386	5,437	5,765	5,375	5,850	5,968	5,740
Female	3,042	3,249	3,429	3,562	3,655	3,668	3,887	3,650	3,969	4,042	3,853
Male	1,364	1,521	1,602	1,696	1,731	1,769	1,878	1,725	1,881	1,926	1,887
White	4,041	4,390	4,654	4,879	5,008	5,068	5,378	5,003	5,474	5,579	5,356
Minority	119	123	108	104	89	73	54	46	48	59	61

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	70%	71%	72%	72%	71%	73%	67%	72%	71%	67%
Female	70%	72%	74%	75%	75%	74%	76%	70%	75%	74%	69%
Male	62%	65%	66%	66%	66%	65%	67%	62%	67%	66%	62%
White	67%	69%	70%	71%	71%	70%	72%	66%	71%	71%	66%
Minority	25%	25%	23%	22%	19%	16%	12%	10%	10%	12%	11%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	539	625	670	715	753	780	854	804	906	916	889
Alcohol-related disorders	399	443	481	507	555	555	623	554	597	615	596
Anxiety disorders	2,923	3,178	3,389	3,556	3,673	3,747	3,986	3,785	4,035	4,112	3,968
Attention-deficit, conduct, and disruptive behavior disorders	218	237	277	316	368	392	439	420	497	544	564
Delirium, dementia, and amnesic and other cognitive disorders	853	875	861	842	814	798	820	738	736	703	643
Developmental disorders	374	407	431	450	483	486	525	502	546	554	570
Disorders usually diagnosed in infancy, childhood, adolescence	54	61	61	71	75	80	90	85	94	105	116
Impulse control disorders, NEC	116	129	142	148	167	171	183	167	189	200	203
Miscellaneous disorders	1,204	1,310	1,447	1,527	1,615	1,669	1,755	1,682	1,800	1,841	1,782
Mood disorders	2,657	2,883	3,081	3,226	3,347	3,380	3,588	3,381	3,618	3,714	3,633
Personality disorders	352	400	419	447	475	491	513	485	516	516	509
Schizophrenia and other psychotic disorders	849	912	933	940	954	965	1,011	931	952	929	871
Screening and history of mental health and substance abuse	2,682	2,980	3,192	3,402	3,559	3,616	3,881	3,647	4,011	4,136	4,025
Substance-related disorders	889	980	1,067	1,147	1,249	1,317	1,437	1,347	1,486	1,550	1,507
Suicide and intentional self-inflicted injury	143	177	191	207	226	242	283	264	300	328	312



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lawrence County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,345	1,420	1,447	1,513	1,499	1,467	1,502	1,396	1,440	1,386	1,301
Female	617	641	668	671	664	641	647	607	643	605	556
Male	728	779	779	842	835	826	855	789	797	781	745
White	1,250	1,312	1,325	1,380	1,361	1,326	1,341	1,253	1,290	1,232	1,152
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	51%	51%	53%	54%	54%	54%	56%	52%	55%	51%	48%
Female	52%	51%	53%	52%	52%	52%	53%	49%	52%	48%	43%
Male	50%	52%	52%	56%	55%	56%	58%	55%	57%	54%	52%
White	49%	50%	51%	52%	52%	52%	53%	50%	53%	49%	45%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	165	185	191	206	214	201	211	192	192	179	165
Alcohol-related disorders	68	62	57	55	53	42	39	31	N.D.	N.D.	N.D.
Anxiety disorders	438	439	452	446	429	390	392	371	360	341	306
Attention-deficit, conduct, and disruptive behavior disorders	554	602	638	712	706	730	762	676	675	639	596
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	386	431	459	508	530	544	571	545	578	574	564
Disorders usually diagnosed in infancy, childhood, adolescence	91	100	108	121	121	119	138	134	140	132	131
Impulse control disorders, NEC	39	40	38	40	37	33	32	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	183	180	182	184	173	164	161	140	142	135	123
Mood disorders	493	493	497	493	459	409	380	326	312	282	251
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	51	54	55	54	46	45	43	32	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	935	984	1,003	1,042	1,005	966	982	864	877	806	749
Substance-related disorders	155	151	136	133	116	91	85	72	73	58	54
Suicide and intentional self-inflicted injury	33	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lawrence County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,711	1,891	1,990	2,073	2,109	2,141	2,236	2,008	2,143	2,145	2,045
Female	1,128	1,250	1,305	1,360	1,389	1,421	1,477	1,314	1,401	1,409	1,312
Male	583	641	685	713	720	720	759	694	742	736	733
White	1,578	1,746	1,845	1,929	1,958	1,987	2,075	1,857	1,984	1,984	1,881
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	66%	66%	67%	67%	68%	69%	63%	67%	66%	63%
Female	66%	69%	70%	72%	71%	73%	75%	67%	72%	71%	68%
Male	58%	60%	60%	61%	60%	60%	61%	56%	59%	58%	57%
White	62%	65%	66%	67%	67%	67%	69%	62%	66%	66%	63%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	84	86	100	113	120	122	134	120	131	135	122
Alcohol-related disorders	212	231	243	263	262	267	279	255	282	304	267
Anxiety disorders	1,010	1,118	1,190	1,238	1,292	1,314	1,374	1,259	1,310	1,333	1,281
Attention-deficit, conduct, and disruptive behavior disorders	42	53	56	59	72	72	96	95	110	120	135
Delirium, dementia, and amnestic and other cognitive disorders	367	382	389	382	373	359	347	310	309	283	248
Developmental disorders	169	178	179	190	198	193	212	200	198	210	212
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	35	42
Impulse control disorders, NEC	41	45	45	52	54	59	62	60	68	66	59
Miscellaneous disorders	406	453	472	479	502	520	543	490	520	500	467
Mood disorders	1,011	1,119	1,183	1,243	1,290	1,319	1,403	1,264	1,342	1,372	1,293
Personality disorders	160	172	177	183	196	196	207	186	189	195	178
Schizophrenia and other psychotic disorders	295	316	321	328	334	329	342	311	310	311	283
Screening and history of mental health and substance abuse	1,025	1,157	1,237	1,309	1,365	1,416	1,516	1,363	1,491	1,498	1,411
Substance-related disorders	330	376	408	431	469	497	534	487	532	536	528
Suicide and intentional self-inflicted injury	56	63	63	65	64	69	74	68	75	65	62



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lee County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	829	890	887	924	923	923	957	892	910	919	847
Female	398	439	423	441	429	440	446	415	417	408	378
Male	431	451	464	483	494	483	511	477	493	511	469
White	790	845	841	870	865	861	883	825	837	841	780
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	65%	65%	67%	68%	69%	71%	67%	70%	65%	60%
Female	65%	68%	67%	69%	69%	71%	71%	69%	70%	62%	58%
Male	61%	62%	63%	65%	67%	68%	72%	66%	70%	68%	62%
White	62%	64%	64%	66%	67%	68%	70%	66%	69%	64%	60%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	147	155	160	169	159	159	166	159	156	161	154
Alcohol-related disorders	67	69	61	59	50	48	47	36	33	N.D.	N.D.
Anxiety disorders	358	378	388	415	407	409	408	371	373	368	353
Attention-deficit, conduct, and disruptive behavior disorders	295	320	331	352	358	357	369	323	316	310	278
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	266	298	319	341	354	357	371	349	363	363	332
Disorders usually diagnosed in infancy, childhood, adolescence	58	67	69	72	76	78	82	73	78	78	72
Impulse control disorders, NEC	32	33	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	118	129	121	119	110	109	111	99	97	91	81
Mood disorders	286	285	274	273	254	232	223	186	170	155	131
Personality disorders	62	63	61	60	58	52	49	41	35	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	580	614	597	613	605	585	585	533	523	511	456
Substance-related disorders	124	125	119	115	103	86	89	77	72	68	58
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lee County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,130	1,188	1,264	1,290	1,312	1,333	1,426	1,345	1,399	1,429	1,399
Female	734	768	810	817	840	866	915	884	904	902	890
Male	396	420	454	473	472	467	511	461	495	527	509
White	1,054	1,114	1,186	1,218	1,241	1,268	1,346	1,267	1,325	1,351	1,311
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	71%	74%	74%	73%	73%	77%	73%	76%	76%	72%
Female	73%	74%	78%	78%	77%	79%	83%	80%	83%	81%	77%
Male	61%	65%	68%	68%	66%	65%	68%	63%	67%	69%	65%
White	68%	71%	74%	74%	73%	74%	77%	73%	77%	76%	72%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	135	146	159	174	168	179	191	187	193	197	192
Alcohol-related disorders	123	129	144	150	155	155	182	179	180	185	169
Anxiety disorders	771	815	870	900	941	950	1,036	1,007	1,028	1,049	1,017
Attention-deficit, conduct, and disruptive behavior disorders	33	34	40	42	44	53	66	72	84	98	105
Delirium, dementia, and amnestic and other cognitive disorders	277	278	282	268	240	219	218	197	191	188	172
Developmental disorders	97	99	108	116	110	106	126	116	119	134	136
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	32
Miscellaneous disorders	338	371	413	438	455	464	493	476	481	478	468
Mood disorders	644	687	748	776	807	824	899	845	857	889	877
Personality disorders	117	124	133	145	149	151	153	144	146	153	142
Schizophrenia and other psychotic disorders	205	222	231	230	228	213	226	217	221	224	212
Screening and history of mental health and substance abuse	672	713	771	804	831	882	963	901	936	954	939
Substance-related disorders	285	306	334	345	356	387	420	414	426	428	412
Suicide and intentional self-inflicted injury	41	48	51	59	61	64	70	73	78	80	73



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Leslie County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,068	1,085	1,108	1,110	1,083	1,089	1,041	956	956	912	814
Female	546	543	544	532	512	507	486	436	429	414	378
Male	522	542	564	578	571	582	555	520	527	498	436
White	1,030	1,047	1,069	1,063	1,034	1,025	968	879	876	833	735
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	49%	50%	51%	51%	50%	52%	51%	49%	51%	49%	44%
Female	53%	53%	54%	53%	52%	53%	52%	47%	48%	46%	42%
Male	46%	47%	49%	50%	49%	51%	51%	50%	53%	52%	46%
White	49%	49%	51%	51%	49%	51%	50%	47%	49%	47%	41%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	199	202	211	209	210	207	203	192	185	179	154
Alcohol-related disorders	57	58	57	47	47	41	38	31	N.D.	N.D.	N.D.
Anxiety disorders	394	387	390	381	371	371	350	321	315	298	265
Attention-deficit, conduct, and disruptive behavior disorders	297	301	328	324	324	335	335	319	316	284	257
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	255	258	279	293	300	308	304	290	289	276	253
Disorders usually diagnosed in infancy, childhood, adolescence	41	48	52	55	56	56	56	50	50	44	40
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	170	167	167	154	151	148	139	105	108	98	93
Mood disorders	327	316	310	280	254	246	215	176	152	140	109
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	43	43	46	43	42	39	35	33	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	592	597	601	599	580	533	497	437	420	395	338
Substance-related disorders	141	131	125	102	95	91	80	67	69	66	58
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Leslie County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,697	1,774	1,844	1,951	1,983	1,970	1,938	1,843	1,953	1,877	1,766
Female	1,106	1,155	1,219	1,267	1,275	1,274	1,245	1,181	1,261	1,187	1,123
Male	591	619	625	684	708	696	693	662	692	690	643
White	1,579	1,659	1,730	1,838	1,867	1,857	1,824	1,733	1,845	1,768	1,660
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	70%	71%	72%	72%	72%	73%	69%	73%	72%	70%
Female	71%	74%	76%	76%	76%	76%	77%	72%	76%	75%	74%
Male	62%	63%	62%	65%	67%	66%	68%	64%	68%	68%	64%
White	67%	69%	70%	72%	72%	72%	73%	68%	73%	72%	70%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	168	180	184	199	210	211	212	199	221	211	204
Alcohol-related disorders	162	180	180	204	214	212	218	207	211	209	206
Anxiety disorders	1,162	1,236	1,287	1,379	1,410	1,406	1,387	1,320	1,369	1,315	1,250
Attention-deficit, conduct, and disruptive behavior disorders	36	37	39	46	55	60	59	60	66	71	74
Delirium, dementia, and amnestic and other cognitive disorders	341	352	338	327	316	307	292	255	242	233	210
Developmental disorders	120	127	130	145	147	145	155	149	144	147	142
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	33	33	34	35	32	N.D.	N.D.	N.D.
Miscellaneous disorders	413	447	476	524	554	557	531	517	533	511	475
Mood disorders	985	1,044	1,091	1,180	1,202	1,206	1,202	1,163	1,226	1,184	1,133
Personality disorders	134	138	146	171	183	178	172	160	156	146	141
Schizophrenia and other psychotic disorders	403	428	418	413	410	401	388	350	344	329	316
Screening and history of mental health and substance abuse	919	982	1,048	1,145	1,198	1,197	1,200	1,136	1,183	1,135	1,079
Substance-related disorders	397	424	439	495	516	536	535	521	557	551	539
Suicide and intentional self-inflicted injury	37	43	44	52	55	62	66	59	63	63	61



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Letcher County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,366	2,425	2,437	2,461	2,437	2,397	2,403	2,277	2,321	2,281	2,152
Female	1,131	1,138	1,136	1,137	1,102	1,081	1,088	1,026	1,050	1,012	950
Male	1,235	1,287	1,301	1,324	1,335	1,316	1,315	1,251	1,271	1,269	1,202
White	2,232	2,281	2,292	2,312	2,285	2,239	2,229	2,112	2,153	2,116	2,009
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	59%	61%	61%	62%	63%	65%	62%	64%	59%	55%
Female	58%	60%	61%	60%	61%	63%	64%	59%	63%	56%	52%
Male	55%	58%	61%	61%	63%	64%	66%	64%	66%	61%	57%
White	55%	58%	60%	60%	61%	62%	64%	61%	63%	58%	54%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	527	547	557	563	550	538	531	499	500	477	430
Alcohol-related disorders	108	107	102	98	91	80	65	56	52	51	36
Anxiety disorders	902	900	905	898	874	869	860	804	790	754	684
Attention-deficit, conduct, and disruptive behavior disorders	734	772	784	812	817	810	800	738	737	717	646
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	31	N.D.	31	33	36	36	N.D.	N.D.	N.D.	N.D.
Developmental disorders	648	708	739	802	844	853	869	840	857	857	807
Disorders usually diagnosed in infancy, childhood, adolescence	110	119	118	122	119	126	133	133	133	129	117
Impulse control disorders, NEC	93	99	95	97	95	86	86	78	63	54	39
Miscellaneous disorders	420	416	413	420	398	400	388	375	377	372	361
Mood disorders	742	714	674	636	580	525	485	420	372	334	274
Personality disorders	99	95	94	86	80	73	65	56	46	41	N.D.
Schizophrenia and other psychotic disorders	64	63	60	57	56	49	50	45	36	34	N.D.
Screening and history of mental health and substance abuse	1,591	1,629	1,618	1,572	1,517	1,430	1,395	1,293	1,271	1,213	1,099
Substance-related disorders	380	348	322	298	264	236	213	188	174	155	128
Suicide and intentional self-inflicted injury	55	53	47	48	43	38	37	33	N.D.	N.D.	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Letcher County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,898	3,069	3,204	3,365	3,472	3,436	3,516	3,244	3,367	3,389	3,294
Female	1,873	2,000	2,072	2,176	2,272	2,245	2,266	2,112	2,190	2,207	2,119
Male	1,025	1,069	1,132	1,189	1,200	1,191	1,250	1,132	1,177	1,182	1,175
White	2,689	2,858	2,994	3,146	3,253	3,211	3,294	3,032	3,161	3,180	3,072
Minority	54	49	42	39	36	33	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	66%	67%	68%	68%	70%	71%	67%	71%	71%	69%
Female	67%	70%	70%	71%	72%	73%	74%	70%	74%	74%	71%
Male	60%	60%	62%	62%	62%	64%	66%	61%	65%	67%	65%
White	64%	66%	67%	67%	68%	69%	71%	66%	70%	71%	68%
Minority	19%	17%	15%	14%	13%	12%	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	263	276	290	320	350	363	388	373	391	409	404
Alcohol-related disorders	363	389	415	448	473	463	466	436	427	426	414
Anxiety disorders	1,792	1,946	2,047	2,193	2,287	2,271	2,338	2,187	2,243	2,242	2,180
Attention-deficit, conduct, and disruptive behavior disorders	77	84	99	117	143	156	174	163	191	219	227
Delirium, dementia, and amnesic and other cognitive disorders	413	439	434	413	414	409	398	359	355	322	310
Developmental disorders	247	252	253	262	289	301	314	293	316	312	316
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	34	38	38	38	41	43	48
Impulse control disorders, NEC	62	66	64	68	76	77	83	73	83	82	93
Miscellaneous disorders	986	1,077	1,147	1,228	1,297	1,297	1,318	1,234	1,229	1,193	1,156
Mood disorders	1,709	1,821	1,912	2,027	2,144	2,129	2,201	2,069	2,119	2,108	2,060
Personality disorders	244	253	275	299	330	318	317	287	292	294	286
Schizophrenia and other psychotic disorders	548	565	572	583	589	573	581	551	547	516	491
Screening and history of mental health and substance abuse	1,939	2,090	2,218	2,345	2,454	2,459	2,531	2,361	2,434	2,444	2,356
Substance-related disorders	785	863	914	991	1,066	1,073	1,140	1,070	1,131	1,133	1,139
Suicide and intentional self-inflicted injury	144	150	156	172	181	191	198	188	197	198	197



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lewis County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,045	1,124	1,189	1,236	1,250	1,226	1,277	1,172	1,191	1,192	1,121
Female	536	565	584	610	610	589	600	551	559	564	525
Male	509	559	605	626	640	637	677	621	632	628	596
White	987	1,056	1,113	1,154	1,166	1,133	1,174	1,079	1,088	1,092	1,028
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	50%	50%	53%	54%	54%	52%	54%	50%	50%	49%	46%
Female	52%	52%	53%	53%	53%	51%	52%	47%	47%	47%	44%
Male	48%	49%	53%	54%	54%	53%	56%	52%	53%	52%	48%
White	49%	49%	52%	52%	53%	51%	52%	48%	49%	48%	45%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	203	201	218	215	215	209	210	207	216	209	184
Alcohol-related disorders	47	49	51	47	44	37	34	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	325	349	372	371	353	332	333	294	299	291	268
Attention-deficit, conduct, and disruptive behavior disorders	407	443	479	502	508	507	533	489	503	505	461
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	241	268	309	338	375	391	429	411	440	445	430
Disorders usually diagnosed in infancy, childhood, adolescence	90	97	113	119	125	122	127	120	128	124	125
Impulse control disorders, NEC	33	34	39	39	38	34	38	32	N.D.	N.D.	N.D.
Miscellaneous disorders	165	173	191	202	191	191	195	174	176	166	156
Mood disorders	350	368	384	369	341	313	304	251	246	234	205
Personality disorders	106	112	121	118	115	112	102	86	83	75	67
Schizophrenia and other psychotic disorders	50	49	48	46	45	44	42	39	37	38	35
Screening and history of mental health and substance abuse	744	784	825	848	843	816	827	728	723	707	659
Substance-related disorders	114	123	118	111	96	77	69	52	45	36	34
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lewis County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,254	1,307	1,367	1,416	1,492	1,506	1,606	1,493	1,609	1,600	1,544
Female	809	841	895	924	980	987	1,058	964	1,052	1,057	1,009
Male	445	466	472	492	512	519	548	529	557	543	535
White	1,151	1,203	1,257	1,301	1,378	1,391	1,480	1,378	1,483	1,475	1,412
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	64%	66%	67%	69%	66%	66%	63%	67%	67%	64%
Female	65%	67%	69%	71%	72%	70%	70%	65%	72%	71%	67%
Male	57%	59%	60%	61%	63%	59%	61%	59%	60%	60%	58%
White	61%	64%	65%	66%	68%	65%	65%	61%	66%	66%	63%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	107	122	131	153	162	166	188	192	204	213	207
Alcohol-related disorders	135	149	153	164	174	182	195	194	194	196	185
Anxiety disorders	793	839	880	934	969	980	1,029	987	1,017	994	948
Attention-deficit, conduct, and disruptive behavior disorders	58	66	71	81	88	100	109	105	121	121	125
Delirium, dementia, and amnestic and other cognitive disorders	296	299	301	286	284	264	253	242	246	226	212
Developmental disorders	134	133	142	150	151	159	166	158	157	159	168
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	40	41	38
Miscellaneous disorders	233	249	276	303	335	337	360	336	356	352	344
Mood disorders	693	737	762	812	850	852	911	859	912	910	875
Personality disorders	147	153	164	188	197	198	208	193	196	191	189
Schizophrenia and other psychotic disorders	270	281	290	289	305	308	324	306	310	299	287
Screening and history of mental health and substance abuse	738	805	876	925	994	992	1,087	1,023	1,115	1,107	1,082
Substance-related disorders	206	216	230	253	291	306	334	314	341	319	314
Suicide and intentional self-inflicted injury	31	33	N.D.	35	38	39	47	52	50	58	58



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lincoln County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,461	1,670	1,749	1,846	1,931	2,009	2,032	1,902	1,963	1,966	1,839
Female	737	838	844	875	899	915	920	857	876	854	783
Male	724	832	905	971	1,032	1,094	1,112	1,045	1,087	1,112	1,056
White	1,345	1,530	1,593	1,666	1,739	1,807	1,823	1,704	1,754	1,754	1,652
Minority	35	41	49	60	57	58	54	52	55	56	53

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	58%	60%	61%	62%	63%	62%	58%	59%	55%	50%
Female	56%	60%	61%	60%	61%	61%	59%	55%	56%	51%	45%
Male	53%	57%	60%	61%	63%	64%	64%	60%	62%	59%	54%
White	54%	57%	59%	59%	61%	61%	60%	57%	57%	54%	48%
Minority	19%	21%	23%	26%	23%	23%	20%	19%	20%	19%	19%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	462	527	550	576	581	563	565	510	521	514	449
Alcohol-related disorders	39	40	38	40	40	40	37	33	31	32	N.D.
Anxiety disorders	633	709	744	793	835	881	912	839	851	826	763
Attention-deficit, conduct, and disruptive behavior disorders	611	692	741	793	833	856	875	820	808	795	730
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	302	371	417	481	541	570	603	570	633	668	640
Disorders usually diagnosed in infancy, childhood, adolescence	133	150	171	191	207	219	227	227	250	255	236
Impulse control disorders, NEC	56	60	71	69	64	62	54	48	53	51	44
Miscellaneous disorders	292	335	338	336	340	355	372	341	338	335	302
Mood disorders	530	551	531	515	510	502	469	417	398	370	309
Personality disorders	51	49	45	40	40	41	34	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	56	57	59	59	60	59	58	58	62	54	40
Screening and history of mental health and substance abuse	801	860	870	902	858	834	796	707	727	706	627
Substance-related disorders	145	143	134	138	135	130	114	107	94	88	68
Suicide and intentional self-inflicted injury	52	52	53	51	46	50	51	49	46	41	32



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lincoln County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,742	1,937	2,026	2,127	2,164	2,221	2,304	2,106	2,246	2,359	2,353
Female	1,181	1,342	1,404	1,460	1,489	1,533	1,583	1,432	1,540	1,594	1,572
Male	561	595	622	667	675	688	721	674	706	765	781
White	1,572	1,760	1,842	1,943	1,982	2,037	2,119	1,937	2,074	2,185	2,170
Minority	89	92	86	81	81	73	62	58	61	60	62

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	69%	70%	72%	70%	70%	73%	66%	69%	68%	65%
Female	69%	73%	74%	76%	75%	74%	77%	69%	73%	73%	69%
Male	58%	61%	63%	64%	62%	62%	66%	60%	61%	60%	58%
White	65%	69%	69%	72%	70%	70%	73%	66%	69%	69%	65%
Minority	35%	37%	35%	32%	32%	29%	25%	23%	23%	23%	22%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	300	350	364	410	425	447	463	423	462	487	501
Alcohol-related disorders	153	167	171	185	184	184	207	184	187	189	185
Anxiety disorders	1,094	1,218	1,269	1,368	1,433	1,466	1,553	1,443	1,531	1,584	1,573
Attention-deficit, conduct, and disruptive behavior disorders	99	109	113	142	145	162	184	183	215	242	262
Delirium, dementia, and amnestic and other cognitive disorders	445	451	458	449	431	421	407	359	360	354	336
Developmental disorders	201	215	208	216	219	222	235	214	224	228	225
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	32	38
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31
Miscellaneous disorders	469	539	576	627	636	679	733	652	676	676	674
Mood disorders	1,043	1,187	1,259	1,374	1,414	1,467	1,536	1,426	1,498	1,565	1,550
Personality disorders	194	229	226	251	264	265	282	254	256	255	248
Schizophrenia and other psychotic disorders	493	506	528	524	513	512	502	454	456	462	445
Screening and history of mental health and substance abuse	846	965	1,051	1,121	1,162	1,195	1,248	1,139	1,232	1,295	1,291
Substance-related disorders	237	272	285	324	339	358	392	366	398	419	432
Suicide and intentional self-inflicted injury	79	88	87	98	104	103	122	120	133	134	135



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Livingston County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	436	471	520	509	469	451	458	443	465	429	395
Female	234	241	254	243	216	199	201	195	209	194	170
Male	202	230	266	266	253	252	257	248	256	235	225
White	N.D.	N.D.	N.D.	474	432	413	414	393	413	381	348
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	#N/A	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	55%	57%	55%	50%	50%	51%	48%	50%	45%	38%
Female	53%	55%	56%	53%	48%	47%	47%	44%	46%	41%	34%
Male	51%	54%	58%	56%	52%	53%	55%	52%	53%	49%	43%
White	52%	54%	56%	54%	49%	49%	49%	45%	47%	42%	36%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	#N/A	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	76	71	72	70	59	59	68	61	63	58	53
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	132	133	136	129	120	110	117	107	108	95	92
Attention-deficit, conduct, and disruptive behavior disorders	218	247	271	275	273	263	274	270	269	233	219
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	43	52	69	76	79	83	93	97	108	110	109
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	35	34	34	37	42	38	45	41	43
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	61	62	62	61	58	51	45	38	36	N.D.	N.D.
Mood disorders	165	170	183	167	158	145	145	123	128	112	91
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	265	276	307	291	250	222	210	196	212	190	175
Substance-related disorders	N.D.	N.D.	34	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Livingston County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	488	522	551	565	564	565	593	511	601	601	565
Female	354	371	384	401	395	396	417	359	419	421	398
Male	134	151	167	164	169	169	176	152	182	180	167
White	447	476	505	520	521	516	539	459	548	544	509
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	63%	64%	65%	66%	64%	69%	60%	69%	68%	63%
Female	64%	67%	66%	68%	68%	67%	72%	62%	70%	70%	64%
Male	50%	54%	59%	58%	60%	58%	64%	56%	66%	64%	59%
White	58%	61%	63%	63%	64%	63%	68%	59%	68%	67%	61%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	39	42	42	45	44	46	50	44	48	50	42
Alcohol-related disorders	38	42	44	43	47	50	55	43	46	48	50
Anxiety disorders	239	258	288	309	326	342	361	309	367	376	353
Attention-deficit, conduct, and disruptive behavior disorders	31	34	45	38	44	58	56	55	76	75	71
Delirium, dementia, and amnesic and other cognitive disorders	124	131	137	124	107	107	113	100	109	95	88
Developmental disorders	35	37	45	46	46	47	47	43	44	46	44
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	107	120	119	125	130	146	145	128	146	149	145
Mood disorders	266	289	312	331	337	350	371	323	385	384	358
Personality disorders	N.D.	N.D.	34	33	37	37	39	35	43	46	40
Schizophrenia and other psychotic disorders	102	107	116	111	103	109	122	114	128	118	109
Screening and history of mental health and substance abuse	262	293	306	331	340	355	377	320	355	360	338
Substance-related disorders	64	68	77	81	90	100	110	91	107	105	99
Suicide and intentional self-inflicted injury	N.D.	N.D.	31	31	35	43	48	42	51	58	49



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Logan County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,034	1,193	1,297	1,405	1,386	1,447	1,448	1,351	1,377	1,388	1,309
Female	520	604	636	683	663	701	697	638	643	639	602
Male	514	589	661	722	723	746	751	713	734	749	707
White	836	973	1,067	1,144	1,126	1,178	1,156	1,087	1,107	1,127	1,069
Minority	166	182	181	204	197	197	199	185	185	176	168

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	45%	46%	47%	48%	47%	47%	48%	44%	45%	44%	40%
Female	45%	46%	46%	47%	44%	46%	46%	42%	43%	42%	37%
Male	44%	47%	49%	50%	49%	48%	49%	46%	47%	46%	42%
White	45%	46%	47%	48%	46%	46%	46%	43%	44%	43%	39%
Minority	37%	40%	38%	41%	38%	38%	37%	34%	35%	33%	31%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	143	165	177	187	185	192	177	149	144	141	128
Alcohol-related disorders	31	35	N.D.	36	33	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	274	320	352	376	379	407	412	383	398	399	357
Attention-deficit, conduct, and disruptive behavior disorders	455	513	575	618	610	620	604	561	568	561	516
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	311	370	429	474	483	540	569	531	539	559	539
Disorders usually diagnosed in infancy, childhood, adolescence	86	104	124	147	162	178	180	166	167	161	159
Impulse control disorders, NEC	81	85	90	89	89	84	75	61	52	49	40
Miscellaneous disorders	151	160	179	201	203	203	203	189	182	175	166
Mood disorders	360	393	420	435	399	377	347	306	296	275	244
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	48	44	45	50	47	44	42	39	35	37	31
Screening and history of mental health and substance abuse	572	619	617	644	590	570	531	464	466	443	393
Substance-related disorders	85	90	86	92	79	79	69	59	57	42	35
Suicide and intentional self-inflicted injury	N.D.	N.D.	31	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Logan County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,331	1,464	1,485	1,531	1,578	1,624	1,659	1,547	1,578	1,599	1,537
Female	973	1,048	1,048	1,094	1,123	1,174	1,195	1,105	1,135	1,130	1,076
Male	358	416	437	437	455	450	464	442	443	469	461
White	1,089	1,202	1,220	1,252	1,294	1,346	1,381	1,277	1,295	1,325	1,270
Minority	174	189	187	199	197	190	179	170	182	179	165

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	63%	64%	64%	65%	65%	67%	63%	64%	63%	58%
Female	62%	65%	66%	66%	66%	67%	69%	64%	67%	65%	60%
Male	54%	60%	60%	61%	62%	61%	62%	60%	59%	59%	55%
White	61%	65%	65%	65%	66%	66%	67%	62%	64%	63%	58%
Minority	39%	41%	43%	44%	43%	42%	42%	41%	43%	42%	37%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	82	90	95	104	109	109	120	121	127	128	137
Alcohol-related disorders	116	130	134	148	163	163	160	148	142	148	139
Anxiety disorders	652	735	778	829	855	890	913	863	867	891	866
Attention-deficit, conduct, and disruptive behavior disorders	50	61	66	85	92	100	132	126	132	149	165
Delirium, dementia, and amnestic and other cognitive disorders	415	413	391	368	354	354	336	298	298	280	247
Developmental disorders	153	172	171	176	186	191	206	200	201	199	204
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	32	34	31
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	39	37	43	47	47
Miscellaneous disorders	233	279	292	312	323	328	372	340	362	378	350
Mood disorders	717	803	828	894	933	970	999	932	978	983	937
Personality disorders	55	60	56	65	67	69	73	65	64	62	56
Schizophrenia and other psychotic disorders	331	354	365	372	369	369	362	334	331	306	280
Screening and history of mental health and substance abuse	622	735	778	823	882	897	928	872	891	925	907
Substance-related disorders	171	195	199	213	231	228	236	205	212	222	220
Suicide and intentional self-inflicted injury	34	38	42	46	58	55	61	54	62	65	67



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lyon County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	238	242	268	269	258	265	280	265	276	297	264
Female	99	107	124	126	122	125	130	126	135	142	123
Male	139	135	144	143	136	140	150	139	141	155	141
White	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	248	N.D.	245	264	N.D.
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	52%	60%	57%	56%	57%	56%	52%	52%	50%	44%
Female	47%	49%	57%	58%	55%	56%	55%	53%	54%	49%	42%
Male	58%	55%	62%	56%	56%	57%	57%	52%	50%	50%	45%
White	52%	52%	59%	55%	54%	55%	55%	51%	50%	48%	42%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	34	39	46	41	38	38	38	35	35	43	35
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	62	66	74	78	75	73	76	70	69	77	72
Attention-deficit, conduct, and disruptive behavior disorders	120	117	128	145	146	161	170	150	144	150	133
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	43	42	49	53	61	65	71	75	77	86	82
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	N.D.	34	31	N.D.	N.D.	N.D.	31	N.D.	N.D.	33	N.D.
Mood disorders	87	90	97	103	93	89	90	80	81	81	72
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	131	128	141	138	126	114	116	107	110	117	94
Substance-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Lyon County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	409	428	427	443	419	423	451	416	435	435	426
Female	305	326	325	328	311	301	325	300	316	312	302
Male	104	102	102	115	108	122	126	116	119	123	124
White	N.D.	397	N.D.	N.D.	N.D.	391	419	379	403	404	392
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	67%	71%	70%	70%	69%	68%	71%	66%	74%	69%	64%
Female	70%	75%	74%	73%	73%	69%	73%	69%	77%	71%	65%
Male	60%	62%	59%	63%	60%	65%	67%	59%	66%	65%	62%
White	67%	72%	70%	70%	70%	68%	71%	66%	74%	70%	64%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	N.D.	N.D.	N.D.	31	N.D.	N.D.	33	35	38	35	31
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	178	193	214	233	219	232	253	243	247	247	246
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	N.D.	31	32	37	46	46	51	57	48
Delirium, dementia, and amnesic and other cognitive disorders	194	189	173	166	165	148	145	140	135	129	113
Developmental disorders	33	35	36	35	36	36	42	40	42	43	45
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	67	74	78	89	83	91	109	99	104	109	104
Mood disorders	193	214	234	258	233	245	270	263	268	278	263
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	87	92	98	101	99	98	106	111	111	109	98
Screening and history of mental health and substance abuse	141	162	172	193	186	201	216	194	213	219	212
Substance-related disorders	N.D.	37	42	46	49	49	61	58	60	57	59
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Madison County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,197	3,602	3,829	4,162	4,303	4,442	4,621	4,501	4,801	4,830	4,509
Female	1,666	1,824	1,887	2,043	2,068	2,119	2,171	2,085	2,225	2,214	2,066
Male	1,531	1,778	1,942	2,119	2,235	2,323	2,450	2,416	2,576	2,616	2,443
White	2,748	3,109	3,287	3,561	3,677	3,805	3,948	3,833	4,075	4,111	3,833
Minority	301	326	344	371	377	368	369	367	398	386	358

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	59%	62%	63%	63%	63%	64%	61%	61%	58%	52%
Female	57%	59%	62%	63%	61%	61%	61%	58%	58%	54%	49%
Male	54%	58%	62%	64%	65%	66%	67%	64%	64%	61%	55%
White	55%	58%	61%	62%	62%	63%	63%	59%	60%	56%	51%
Minority	40%	42%	43%	44%	40%	40%	39%	39%	41%	36%	33%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	881	969	1,041	1,097	1,089	1,102	1,114	1,051	1,033	958	860
Alcohol-related disorders	162	162	164	169	165	167	157	139	129	111	90
Anxiety disorders	1,392	1,576	1,703	1,834	1,903	1,985	2,082	2,025	2,105	2,087	1,953
Attention-deficit, conduct, and disruptive behavior disorders	1,334	1,488	1,612	1,757	1,785	1,823	1,870	1,759	1,835	1,771	1,642
Delirium, dementia, and amnestic and other cognitive disorders	50	54	56	55	56	58	58	51	54	45	34
Developmental disorders	1,014	1,191	1,366	1,540	1,680	1,786	1,896	1,893	2,061	2,069	1,985
Disorders usually diagnosed in infancy, childhood, adolescence	320	393	456	499	539	574	616	591	661	675	644
Impulse control disorders, NEC	123	130	140	148	143	139	130	107	94	77	65
Miscellaneous disorders	632	705	765	844	866	907	940	931	971	981	915
Mood disorders	1,092	1,120	1,146	1,159	1,103	1,071	1,051	969	956	904	766
Personality disorders	126	136	139	138	130	121	113	100	101	87	79
Schizophrenia and other psychotic disorders	150	159	168	185	186	189	180	168	170	150	131
Screening and history of mental health and substance abuse	1,831	1,971	2,018	2,110	2,040	1,982	1,970	1,815	1,810	1,722	1,513
Substance-related disorders	422	434	429	454	418	386	378	342	327	304	254
Suicide and intentional self-inflicted injury	129	129	137	146	140	146	142	145	143	135	123



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Madison County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,377	3,693	3,876	4,019	4,198	4,346	4,570	4,377	4,798	4,974	4,925
Female	2,479	2,674	2,827	2,934	3,070	3,182	3,330	3,125	3,431	3,516	3,463
Male	898	1,019	1,049	1,085	1,128	1,164	1,240	1,252	1,367	1,458	1,462
White	2,948	3,219	3,401	3,528	3,678	3,825	4,035	3,871	4,267	4,433	4,362
Minority	240	274	267	272	284	275	271	234	252	268	263

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	68%	70%	70%	70%	71%	72%	68%	71%	69%	66%
Female	67%	70%	71%	72%	71%	72%	73%	68%	72%	70%	68%
Male	58%	63%	65%	66%	66%	68%	68%	66%	68%	66%	63%
White	64%	68%	70%	71%	71%	71%	72%	68%	71%	70%	67%
Minority	36%	37%	37%	36%	36%	35%	35%	31%	32%	33%	30%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	774	844	928	1,009	1,074	1,128	1,209	1,170	1,311	1,359	1,327
Alcohol-related disorders	284	325	342	375	393	393	435	410	456	452	435
Anxiety disorders	2,043	2,254	2,415	2,563	2,675	2,817	2,976	2,846	3,102	3,180	3,162
Attention-deficit, conduct, and disruptive behavior disorders	210	228	254	287	329	385	439	428	505	569	566
Delirium, dementia, and amnesic and other cognitive disorders	869	912	920	878	900	898	890	842	837	797	726
Developmental disorders	442	493	496	507	532	545	584	583	625	659	659
Disorders usually diagnosed in infancy, childhood, adolescence	54	59	65	67	68	75	91	95	110	134	145
Impulse control disorders, NEC	47	49	53	60	66	69	87	88	107	123	121
Miscellaneous disorders	960	1,047	1,148	1,207	1,318	1,364	1,489	1,436	1,526	1,595	1,562
Mood disorders	2,098	2,289	2,424	2,575	2,719	2,825	3,010	2,861	3,102	3,197	3,165
Personality disorders	273	288	326	345	363	377	418	378	405	418	399
Schizophrenia and other psychotic disorders	747	791	797	799	797	817	865	815	827	834	784
Screening and history of mental health and substance abuse	1,802	2,007	2,167	2,325	2,485	2,585	2,769	2,699	3,007	3,125	3,153
Substance-related disorders	621	691	758	842	914	965	1,064	1,015	1,146	1,180	1,166
Suicide and intentional self-inflicted injury	119	134	153	166	187	201	234	217	242	250	257



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Magoffin County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,516	1,574	1,582	1,652	1,610	1,618	1,635	1,561	1,593	1,592	1,477
Female	767	782	780	807	789	782	779	737	743	733	687
Male	749	792	802	845	821	836	856	824	850	859	790
White	1,422	1,465	1,457	1,521	1,477	1,481	1,482	1,409	1,432	1,423	1,317
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	#N/A	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	60%	62%	63%	61%	62%	62%	60%	63%	59%	55%
Female	61%	63%	65%	66%	63%	62%	62%	60%	62%	58%	54%
Male	55%	57%	60%	61%	60%	62%	62%	61%	64%	60%	55%
White	57%	59%	61%	62%	60%	60%	60%	58%	61%	57%	53%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	#N/A	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	239	259	250	265	255	251	256	230	230	209	195
Alcohol-related disorders	44	45	43	45	42	37	34	31	N.D.	N.D.	N.D.
Anxiety disorders	531	537	548	575	580	589	610	582	582	580	540
Attention-deficit, conduct, and disruptive behavior disorders	472	527	549	591	599	620	648	601	625	626	581
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	272	316	343	387	412	431	453	447	483	498	474
Disorders usually diagnosed in infancy, childhood, adolescence	73	85	87	94	101	109	115	112	116	121	112
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	31	34	33	N.D.	N.D.	N.D.
Miscellaneous disorders	315	324	323	331	324	329	335	309	303	305	279
Mood disorders	405	401	356	338	312	286	293	245	225	225	196
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	1,058	1,093	1,071	1,096	1,039	996	983	903	907	875	781
Substance-related disorders	157	141	123	121	97	89	83	66	59	55	39
Suicide and intentional self-inflicted injury	34	37	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Magoffin County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,130	2,182	2,280	2,332	2,422	2,443	2,501	2,348	2,434	2,424	2,384
Female	1,361	1,394	1,453	1,490	1,534	1,542	1,583	1,470	1,517	1,531	1,491
Male	769	788	827	842	888	901	918	878	917	893	893
White	1,996	2,041	2,130	2,181	2,265	2,292	2,340	2,189	2,274	2,262	2,216
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	73%	76%	76%	76%	75%	75%	77%	72%	76%	74%	71%
Female	77%	79%	80%	80%	80%	79%	81%	76%	80%	80%	76%
Male	68%	70%	70%	69%	68%	69%	70%	66%	70%	66%	65%
White	73%	75%	76%	75%	75%	75%	76%	71%	75%	74%	71%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	188	194	213	215	226	237	240	240	247	253	240
Alcohol-related disorders	195	197	208	213	221	216	228	219	218	207	208
Anxiety disorders	1,444	1,494	1,581	1,616	1,691	1,727	1,789	1,688	1,734	1,727	1,683
Attention-deficit, conduct, and disruptive behavior disorders	69	71	75	78	89	93	104	103	110	124	134
Delirium, dementia, and amnesic and other cognitive disorders	406	407	409	404	403	388	376	346	326	297	262
Developmental disorders	173	188	182	184	179	172	174	166	162	155	153
Disorders usually diagnosed in infancy, childhood, adolescence	41	41	43	47	51	52	54	54	54	52	54
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	792	823	892	912	964	990	1,008	966	969	939	888
Mood disorders	1,197	1,254	1,325	1,342	1,400	1,432	1,477	1,367	1,402	1,374	1,340
Personality disorders	85	91	98	96	105	110	109	102	112	105	104
Schizophrenia and other psychotic disorders	345	343	342	338	342	337	339	301	302	283	255
Screening and history of mental health and substance abuse	1,344	1,392	1,488	1,562	1,643	1,641	1,696	1,621	1,682	1,675	1,652
Substance-related disorders	412	421	467	491	528	522	554	548	580	568	590
Suicide and intentional self-inflicted injury	50	48	55	59	68	70	76	73	79	77	79



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Marion County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	887	972	1,029	1,106	1,161	1,152	1,176	1,084	1,149	1,171	1,113
Female	439	478	498	531	529	523	537	493	518	521	492
Male	448	494	531	575	632	629	639	591	631	650	621
White	710	770	818	876	919	916	916	835	888	910	871
Minority	114	120	115	123	126	121	125	115	126	128	122

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	50%	53%	56%	58%	59%	58%	60%	54%	54%	49%	46%
Female	51%	54%	55%	57%	55%	54%	56%	50%	49%	44%	40%
Male	50%	53%	56%	60%	64%	63%	63%	59%	58%	54%	51%
White	49%	51%	54%	57%	58%	57%	57%	51%	51%	47%	43%
Minority	36%	37%	33%	34%	35%	33%	34%	31%	32%	30%	29%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	215	251	259	291	288	277	270	250	248	246	242
Alcohol-related disorders	59	52	48	55	54	45	41	32	33	N.D.	N.D.
Anxiety disorders	288	303	309	329	330	320	331	299	303	306	272
Attention-deficit, conduct, and disruptive behavior disorders	426	480	497	537	575	556	551	519	538	533	514
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	212	263	320	368	413	426	447	433	463	478	462
Disorders usually diagnosed in infancy, childhood, adolescence	60	66	81	92	92	99	100	97	102	110	110
Impulse control disorders, NEC	62	63	59	62	61	56	55	42	44	44	40
Miscellaneous disorders	159	173	179	194	178	174	172	160	163	153	138
Mood disorders	382	389	374	386	377	353	342	307	296	296	254
Personality disorders	45	46	42	42	39	33	31	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	37	35	33	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	550	595	592	625	604	579	559	501	487	486	454
Substance-related disorders	115	113	110	121	105	89	86	64	67	55	44
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Marion County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,244	1,329	1,399	1,449	1,503	1,530	1,524	1,465	1,572	1,581	1,521
Female	864	916	962	1,009	1,051	1,060	1,079	1,025	1,101	1,094	1,044
Male	380	413	437	440	452	470	445	440	471	487	477
White	1,045	1,119	1,189	1,227	1,261	1,296	1,292	1,238	1,318	1,335	1,275
Minority	140	147	144	154	168	158	151	150	175	170	167

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	64%	67%	69%	69%	70%	71%	67%	70%	69%	67%
Female	64%	66%	68%	71%	72%	72%	74%	68%	71%	71%	69%
Male	58%	59%	64%	64%	64%	67%	64%	64%	66%	65%	63%
White	62%	64%	68%	69%	69%	71%	71%	67%	70%	69%	67%
Minority	43%	43%	42%	46%	49%	45%	43%	44%	48%	47%	44%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	107	121	136	143	147	148	167	152	168	175	169
Alcohol-related disorders	153	167	178	184	195	203	202	189	194	188	183
Anxiety disorders	701	751	808	863	909	913	940	905	981	987	965
Attention-deficit, conduct, and disruptive behavior disorders	61	71	83	99	118	123	126	121	145	148	144
Delirium, dementia, and amnestic and other cognitive disorders	304	312	321	322	316	321	308	296	293	273	251
Developmental disorders	136	142	151	155	164	172	168	171	172	180	182
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	31	33	38	39	44	46	49	45	49	49	47
Miscellaneous disorders	286	299	331	350	364	380	394	379	387	409	386
Mood disorders	766	818	869	904	957	973	1,003	950	994	1,001	968
Personality disorders	179	185	203	213	216	215	219	190	207	205	200
Schizophrenia and other psychotic disorders	301	308	333	347	344	349	336	307	316	302	276
Screening and history of mental health and substance abuse	601	660	715	759	830	841	856	811	889	872	847
Substance-related disorders	171	192	213	223	250	257	257	239	260	266	246
Suicide and intentional self-inflicted injury	65	63	74	76	84	86	94	88	94	93	88



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Marshall County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,024	1,222	1,307	1,400	1,476	1,492	1,509	1,425	1,485	1,501	1,398
Female	492	580	601	643	667	672	673	635	656	668	615
Male	532	642	706	757	809	820	836	790	829	833	783
White	973	1,152	1,230	1,322	1,381	1,388	1,394	1,314	1,369	1,389	1,293
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	49%	55%	54%	54%	55%	54%	55%	50%	51%	50%	45%
Female	47%	53%	53%	52%	52%	51%	51%	47%	47%	46%	40%
Male	50%	56%	56%	57%	57%	56%	58%	53%	55%	54%	50%
White	48%	54%	53%	54%	54%	52%	53%	48%	50%	49%	44%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	129	146	151	158	158	158	155	146	139	140	137
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	342	384	379	412	415	424	418	366	356	327	276
Attention-deficit, conduct, and disruptive behavior disorders	519	641	685	732	783	791	813	765	766	763	716
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	161	205	243	287	338	386	412	422	471	504	499
Disorders usually diagnosed in infancy, childhood, adolescence	64	78	93	101	120	133	141	134	137	132	128
Impulse control disorders, NEC	70	79	84	88	84	92	91	80	69	64	55
Miscellaneous disorders	153	178	187	201	209	211	203	182	175	151	130
Mood disorders	375	438	446	458	463	441	435	384	379	350	306
Personality disorders	32	31	32	34	32	31	34	N.D.	31	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	36	35	32	36	38	38	31	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	540	609	650	671	676	666	661	600	618	610	535
Substance-related disorders	80	84	81	85	88	81	77	60	46	51	45
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Marshall County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,253	1,348	1,388	1,478	1,502	1,535	1,579	1,453	1,635	1,678	1,599
Female	963	1,026	1,063	1,128	1,149	1,151	1,193	1,084	1,215	1,217	1,146
Male	290	322	325	350	353	384	386	369	420	461	453
White	1,181	1,260	1,299	1,385	1,411	1,445	1,481	1,357	1,529	1,569	1,483
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	65%	66%	67%	66%	66%	68%	61%	68%	67%	61%
Female	64%	68%	69%	68%	69%	69%	72%	64%	70%	69%	63%
Male	56%	57%	58%	61%	58%	59%	60%	54%	61%	61%	57%
White	62%	64%	65%	66%	65%	66%	68%	60%	67%	66%	60%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	85	90	84	96	96	106	103	99	116	122	125
Alcohol-related disorders	63	77	82	89	96	102	110	93	112	118	111
Anxiety disorders	705	784	823	889	937	963	1,006	911	1,055	1,102	1,049
Attention-deficit, conduct, and disruptive behavior disorders	43	52	54	76	81	96	102	99	129	162	167
Delirium, dementia, and amnestic and other cognitive disorders	387	388	387	380	347	324	333	302	312	308	270
Developmental disorders	84	92	95	101	103	102	112	109	118	129	129
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	35	40	40
Miscellaneous disorders	230	253	284	321	338	345	370	333	366	364	328
Mood disorders	718	790	842	928	952	982	1,015	938	1,055	1,093	1,042
Personality disorders	70	79	84	103	109	103	106	101	114	125	108
Schizophrenia and other psychotic disorders	319	345	350	353	330	310	331	307	319	316	280
Screening and history of mental health and substance abuse	506	599	629	695	716	764	790	747	848	856	807
Substance-related disorders	130	153	173	188	204	219	234	210	230	238	208
Suicide and intentional self-inflicted injury	42	40	44	56	60	68	87	89	102	112	106



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Martin County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,399	1,448	1,449	1,507	1,489	1,445	1,418	1,335	1,397	1,352	1,242
Female	673	681	677	696	690	650	631	588	616	583	535
Male	726	767	772	811	799	795	787	747	781	769	707
White	1,301	1,340	1,332	1,380	1,367	1,322	1,284	1,204	1,257	1,212	1,119
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	56%	58%	59%	59%	59%	59%	56%	59%	58%	54%
Female	55%	57%	58%	58%	58%	57%	57%	53%	56%	54%	50%
Male	52%	55%	58%	59%	59%	60%	60%	59%	61%	62%	57%
White	52%	54%	56%	57%	57%	57%	56%	54%	56%	56%	52%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	209	221	232	233	236	238	235	215	217	206	187
Alcohol-related disorders	87	77	67	63	57	52	50	46	48	46	39
Anxiety disorders	441	418	405	425	426	390	384	340	347	320	291
Attention-deficit, conduct, and disruptive behavior disorders	476	525	543	576	597	585	578	533	549	533	498
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	302	341	358	385	403	417	428	416	438	444	418
Disorders usually diagnosed in infancy, childhood, adolescence	60	72	77	83	86	90	93	89	96	88	82
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	179	171	163	169	155	148	139	126	128	122	110
Mood disorders	505	485	459	457	434	385	348	303	290	258	215
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	35	38	33	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	939	968	975	998	972	919	894	831	857	806	714
Substance-related disorders	207	185	167	162	146	126	110	95	97	91	76
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Martin County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,791	1,913	1,972	2,018	2,062	2,074	2,116	1,944	2,073	2,072	2,065
Female	1,178	1,254	1,292	1,322	1,338	1,338	1,377	1,256	1,331	1,305	1,319
Male	613	659	680	696	724	736	739	688	742	767	746
White	1,644	1,759	1,819	1,862	1,901	1,914	1,945	1,780	1,909	1,914	1,898
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	70%	72%	73%	74%	74%	74%	75%	70%	74%	76%	74%
Female	76%	77%	78%	80%	80%	79%	81%	74%	78%	79%	78%
Male	61%	64%	65%	66%	66%	66%	67%	64%	69%	70%	67%
White	69%	71%	72%	74%	74%	73%	74%	69%	73%	75%	73%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	184	196	209	239	247	250	265	232	236	232	232
Alcohol-related disorders	207	225	239	253	258	259	266	246	258	259	246
Anxiety disorders	1,256	1,337	1,381	1,412	1,427	1,435	1,454	1,348	1,398	1,403	1,389
Attention-deficit, conduct, and disruptive behavior disorders	49	53	66	64	68	79	93	86	98	109	126
Delirium, dementia, and amnestic and other cognitive disorders	230	242	245	250	237	225	221	207	193	192	173
Developmental disorders	80	88	100	105	108	114	122	101	112	110	128
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	39	47	56	54	51	59	64	56	57	57	55
Miscellaneous disorders	458	516	544	557	573	587	600	559	552	538	525
Mood disorders	1,231	1,307	1,368	1,397	1,426	1,448	1,491	1,373	1,427	1,431	1,407
Personality disorders	124	129	137	151	147	135	145	146	152	149	154
Schizophrenia and other psychotic disorders	268	276	289	277	274	266	275	244	243	244	238
Screening and history of mental health and substance abuse	1,168	1,249	1,342	1,404	1,425	1,423	1,466	1,362	1,468	1,460	1,458
Substance-related disorders	422	451	476	513	531	535	561	527	559	565	566
Suicide and intentional self-inflicted injury	64	69	70	74	81	77	79	76	77	76	74



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Mason County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	943	1,026	1,123	1,130	1,162	1,177	1,212	1,150	1,245	1,246	1,149
Female	465	490	540	535	552	558	571	544	578	571	506
Male	478	536	583	595	610	619	641	606	667	675	643
White	787	859	943	943	968	980	999	941	1,019	1,020	940
Minority	108	111	112	117	117	119	124	122	141	138	133

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	57%	60%	59%	59%	56%	56%	53%	54%	49%	44%
Female	58%	58%	60%	57%	56%	55%	53%	52%	53%	47%	40%
Male	56%	56%	61%	60%	61%	58%	59%	53%	55%	51%	47%
White	57%	58%	61%	58%	58%	56%	56%	51%	52%	48%	43%
Minority	38%	36%	35%	38%	37%	35%	34%	34%	38%	33%	31%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	204	219	241	236	242	255	267	274	302	291	282
Alcohol-related disorders	78	77	75	64	65	53	51	37	39	38	N.D.
Anxiety disorders	310	320	351	352	354	355	353	330	332	319	285
Attention-deficit, conduct, and disruptive behavior disorders	457	512	554	566	564	565	582	526	551	546	494
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	299	337	384	403	440	465	487	471	522	530	502
Disorders usually diagnosed in infancy, childhood, adolescence	84	89	109	115	120	123	126	122	126	130	117
Impulse control disorders, NEC	N.D.	N.D.	32	33	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	128	135	140	129	129	121	124	112	129	132	115
Mood disorders	341	346	388	369	355	341	330	293	288	260	215
Personality disorders	71	82	86	90	96	96	88	67	54	46	34
Schizophrenia and other psychotic disorders	50	48	47	51	51	54	55	47	45	45	37
Screening and history of mental health and substance abuse	699	748	820	813	816	789	790	727	750	724	652
Substance-related disorders	130	126	130	121	114	106	97	76	73	65	47
Suicide and intentional self-inflicted injury	N.D.	N.D.	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Mason County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,017	1,093	1,142	1,230	1,269	1,323	1,391	1,276	1,414	1,483	1,383
Female	770	813	841	892	914	960	1,012	913	1,007	1,049	955
Male	247	280	301	338	355	363	379	363	407	434	428
White	891	952	1,010	1,075	1,113	1,175	1,214	1,121	1,245	1,310	1,205
Minority	92	105	96	116	114	106	123	106	118	116	116
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	64%	63%	66%	68%	67%	68%	62%	68%	68%	63%
Female	67%	68%	68%	70%	71%	71%	72%	65%	71%	71%	64%
Male	52%	55%	54%	57%	61%	60%	60%	56%	62%	62%	60%
White	64%	65%	65%	67%	68%	68%	68%	63%	68%	69%	64%
Minority	38%	42%	40%	44%	48%	45%	46%	42%	49%	40%	38%
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	94	98	112	132	139	150	166	148	173	187	176
Alcohol-related disorders	95	99	111	142	138	140	153	137	144	162	160
Anxiety disorders	548	584	647	685	704	724	779	722	779	823	789
Attention-deficit, conduct, and disruptive behavior disorders	86	88	101	127	142	154	160	163	180	191	185
Delirium, dementia, and amnestic and other cognitive disorders	206	206	198	201	204	201	195	189	190	174	162
Developmental disorders	102	113	117	117	134	137	143	139	149	151	154
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	33	N.D.	34	34	35	38	35
Miscellaneous disorders	186	207	227	242	259	275	276	269	293	310	291
Mood disorders	547	581	632	685	707	730	769	712	781	819	793
Personality disorders	127	146	161	175	177	187	199	180	195	210	197
Schizophrenia and other psychotic disorders	195	212	220	235	238	228	234	226	226	235	220
Screening and history of mental health and substance abuse	656	725	770	860	931	974	1,006	914	1,037	1,100	1,014
Substance-related disorders	140	147	172	202	217	240	253	229	282	283	279
Suicide and intentional self-inflicted injury	N.D.	33	33	36	41	45	51	46	48	53	53



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

McCracken County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,300	3,593	3,840	3,902	3,952	3,901	3,783	3,438	3,578	3,528	3,218
Female	1,594	1,728	1,841	1,846	1,829	1,779	1,721	1,562	1,630	1,590	1,437
Male	1,706	1,865	1,999	2,056	2,123	2,122	2,062	1,876	1,948	1,938	1,781
White	2,136	2,358	2,501	2,542	2,588	2,521	2,448	2,221	2,323	2,324	2,107
Minority	974	1,014	1,083	1,078	1,054	1,045	982	903	919	883	794

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	58%	59%	59%	58%	56%	54%	50%	51%	49%	43%
Female	53%	56%	57%	57%	54%	52%	50%	46%	47%	44%	40%
Male	55%	60%	61%	61%	61%	59%	58%	54%	55%	54%	47%
White	56%	59%	60%	59%	58%	55%	54%	50%	50%	49%	42%
Minority	44%	47%	47%	46%	44%	43%	40%	38%	38%	37%	33%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	348	388	395	375	372	370	378	342	348	327	273
Alcohol-related disorders	103	99	100	89	89	80	87	71	69	55	45
Anxiety disorders	714	734	763	743	758	729	708	622	602	579	514
Attention-deficit, conduct, and disruptive behavior disorders	1,774	1,969	2,113	2,178	2,307	2,292	2,245	2,011	2,049	1,995	1,821
Delirium, dementia, and amnestic and other cognitive disorders	45	40	42	50	45	45	43	44	48	46	41
Developmental disorders	689	804	927	1,022	1,076	1,129	1,175	1,090	1,183	1,208	1,154
Disorders usually diagnosed in infancy, childhood, adolescence	202	226	241	281	305	317	332	297	312	313	301
Impulse control disorders, NEC	235	244	248	239	257	245	238	216	199	175	156
Miscellaneous disorders	407	419	417	394	400	390	368	334	335	320	262
Mood disorders	1,156	1,191	1,224	1,186	1,220	1,159	1,124	986	969	937	832
Personality disorders	89	83	78	71	68	60	60	52	53	42	33
Schizophrenia and other psychotic disorders	117	106	111	107	110	110	108	95	102	91	79
Screening and history of mental health and substance abuse	2,111	2,299	2,401	2,342	2,213	2,060	1,951	1,731	1,731	1,642	1,444
Substance-related disorders	255	255	250	238	216	196	196	163	156	133	121
Suicide and intentional self-inflicted injury	107	108	107	98	100	91	88	79	74	75	68



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

McCracken County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,353	3,660	3,875	3,997	4,025	4,107	4,171	3,804	4,089	4,103	3,895
Female	2,579	2,787	2,928	3,021	3,046	3,102	3,147	2,826	3,009	3,005	2,808
Male	774	873	947	976	979	1,005	1,024	978	1,080	1,098	1,087
White	2,380	2,615	2,763	2,877	2,897	2,953	3,012	2,740	2,944	2,971	2,750
Minority	825	878	934	932	932	947	928	831	901	888	879

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	66%	68%	69%	69%	69%	70%	66%	71%	69%	64%
Female	63%	67%	69%	70%	70%	71%	73%	67%	72%	71%	65%
Male	57%	60%	63%	65%	64%	63%	63%	61%	66%	65%	63%
White	62%	66%	68%	70%	69%	69%	71%	65%	71%	70%	64%
Minority	51%	55%	56%	56%	55%	56%	55%	51%	55%	52%	50%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	197	216	243	248	257	272	283	268	276	298	278
Alcohol-related disorders	358	400	441	485	491	492	492	455	487	481	465
Anxiety disorders	1,711	1,888	2,004	2,149	2,204	2,270	2,349	2,214	2,312	2,339	2,217
Attention-deficit, conduct, and disruptive behavior disorders	150	170	195	236	249	294	334	321	381	462	480
Delirium, dementia, and amnestic and other cognitive disorders	767	794	793	751	721	718	723	682	706	674	618
Developmental disorders	257	271	275	286	277	288	304	299	320	338	324
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	32	35	40	40	45	51	57
Impulse control disorders, NEC	48	49	59	66	68	83	98	90	85	117	115
Miscellaneous disorders	662	757	809	874	922	961	1,013	920	984	1,011	948
Mood disorders	1,887	2,098	2,244	2,393	2,439	2,532	2,622	2,421	2,602	2,640	2,514
Personality disorders	254	278	308	320	333	338	360	349	379	401	397
Schizophrenia and other psychotic disorders	592	628	641	661	674	682	689	648	672	688	663
Screening and history of mental health and substance abuse	1,910	2,182	2,402	2,510	2,559	2,616	2,636	2,370	2,523	2,530	2,407
Substance-related disorders	467	536	562	632	654	683	713	650	703	713	698
Suicide and intentional self-inflicted injury	144	169	183	206	230	247	292	273	332	341	327



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

McCreary County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,085	2,272	2,301	2,394	2,401	2,377	2,404	2,225	2,210	2,098	1,890
Female	1,080	1,162	1,164	1,211	1,195	1,168	1,165	1,076	1,069	1,005	886
Male	1,005	1,110	1,137	1,183	1,206	1,209	1,239	1,149	1,141	1,093	1,004
White	1,967	2,127	2,152	2,236	2,237	2,211	2,229	2,051	2,039	1,938	1,739
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	59%	60%	62%	63%	63%	64%	64%	61%	62%	60%	55%
Female	62%	63%	64%	65%	64%	64%	64%	60%	61%	59%	52%
Male	56%	58%	60%	62%	63%	64%	64%	62%	63%	62%	57%
White	58%	59%	61%	62%	62%	62%	63%	60%	61%	59%	53%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	521	541	541	569	552	545	542	482	458	412	344
Alcohol-related disorders	97	95	92	86	81	68	63	52	44	N.D.	N.D.
Anxiety disorders	1,096	1,183	1,192	1,262	1,290	1,273	1,283	1,179	1,170	1,107	999
Attention-deficit, conduct, and disruptive behavior disorders	876	960	968	1,001	1,018	1,005	1,006	878	842	792	715
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	310	368	391	434	460	470	520	516	559	557	525
Disorders usually diagnosed in infancy, childhood, adolescence	122	131	138	151	158	155	154	149	146	135	126
Impulse control disorders, NEC	79	78	76	68	67	65	63	52	49	47	38
Miscellaneous disorders	259	272	262	258	249	235	237	208	199	171	142
Mood disorders	775	789	746	741	708	655	609	491	447	382	327
Personality disorders	47	44	45	44	43	38	36	32	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	75	75	77	77	72	66	62	52	52	48	33
Screening and history of mental health and substance abuse	1,109	1,164	1,166	1,190	1,162	1,104	1,063	903	859	779	689
Substance-related disorders	299	305	278	263	238	204	188	150	144	124	96
Suicide and intentional self-inflicted injury	85	90	88	86	80	66	61	44	42	34	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

McCreary County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,482	2,661	2,727	2,833	2,872	2,833	2,878	2,733	2,886	2,954	2,876
Female	1,642	1,784	1,817	1,857	1,886	1,867	1,893	1,767	1,885	1,901	1,846
Male	840	877	910	976	986	966	985	966	1,001	1,053	1,030
White	2,258	2,425	2,495	2,598	2,631	2,599	2,636	2,498	2,643	2,714	2,620
Minority	34	36	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	71%	72%	74%	73%	72%	73%	70%	72%	73%	70%
Female	73%	76%	78%	78%	78%	78%	78%	73%	77%	77%	74%
Male	60%	62%	62%	66%	66%	63%	65%	64%	65%	67%	64%
White	67%	70%	71%	73%	72%	71%	72%	69%	71%	72%	69%
Minority	12%	13%	11%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	303	318	328	347	360	376	386	389	436	449	451
Alcohol-related disorders	285	303	307	316	319	301	318	299	303	316	313
Anxiety disorders	1,708	1,851	1,915	1,997	2,050	2,036	2,084	2,010	2,117	2,163	2,129
Attention-deficit, conduct, and disruptive behavior disorders	68	74	92	101	120	127	149	183	196	235	269
Delirium, dementia, and amnestic and other cognitive disorders	372	388	381	370	368	350	340	309	307	284	259
Developmental disorders	221	226	237	239	245	257	269	273	278	285	292
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	34	37	42	46
Impulse control disorders, NEC	32	N.D.	36	41	44	45	50	52	50	57	69
Miscellaneous disorders	599	647	660	686	708	718	726	712	717	750	718
Mood disorders	1,614	1,743	1,798	1,886	1,931	1,906	1,953	1,869	1,944	1,976	1,932
Personality disorders	156	163	170	173	175	166	160	151	157	166	173
Schizophrenia and other psychotic disorders	376	395	406	416	429	421	438	423	414	413	396
Screening and history of mental health and substance abuse	1,558	1,650	1,731	1,816	1,861	1,832	1,871	1,803	1,874	1,937	1,856
Substance-related disorders	514	562	588	626	618	619	641	628	656	722	737
Suicide and intentional self-inflicted injury	116	127	132	144	148	155	161	153	164	183	195



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

McLean County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	490	529	563	597	613	626	634	621	649	647	605
Female	270	284	305	317	325	313	315	302	314	318	296
Male	220	245	258	280	288	313	319	319	335	329	309
White	459	493	521	552	565	577	588	571	594	595	557
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	56%	58%	59%	59%	58%	59%	57%	58%	53%	48%
Female	57%	59%	60%	59%	58%	55%	57%	53%	56%	50%	46%
Male	51%	53%	56%	58%	60%	62%	62%	60%	60%	55%	50%
White	53%	55%	57%	57%	58%	57%	58%	55%	56%	52%	47%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	97	115	125	126	128	123	117	109	116	112	108
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	155	173	180	201	207	207	194	180	180	172	155
Attention-deficit, conduct, and disruptive behavior disorders	215	240	256	264	272	281	272	260	275	257	240
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	95	111	131	146	161	176	193	201	212	213	197
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	31	31	35	38	39	44	47	44	39	37
Impulse control disorders, NEC	36	39	43	44	46	47	42	43	40	31	N.D.
Miscellaneous disorders	64	69	70	74	71	68	63	58	63	58	53
Mood disorders	204	222	229	225	225	220	206	189	182	161	147
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	290	315	327	329	322	314	306	294	305	295	271
Substance-related disorders	38	45	42	39	37	35	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

McLean County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	489	527	548	573	592	635	641	560	620	654	633
Female	354	380	394	414	415	450	460	383	416	441	428
Male	135	147	154	159	177	185	181	177	204	213	205
White	450	488	508	532	556	594	598	521	580	609	586
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	67%	68%	69%	69%	71%	70%	62%	69%	67%	62%
Female	64%	69%	70%	71%	71%	73%	73%	62%	68%	67%	63%
Male	55%	62%	63%	64%	66%	67%	64%	62%	69%	68%	61%
White	61%	66%	68%	68%	69%	71%	70%	61%	68%	67%	62%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	47	48	45	52	50	53	55	43	55	62	53
Alcohol-related disorders	40	39	44	51	55	53	64	59	64	72	72
Anxiety disorders	222	256	272	294	290	316	339	295	312	318	311
Attention-deficit, conduct, and disruptive behavior disorders	62	66	71	75	88	94	108	104	110	127	128
Delirium, dementia, and amnesic and other cognitive disorders	142	140	149	140	140	134	129	132	135	143	137
Developmental disorders	68	68	67	68	68	61	60	66	67	78	81
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	75	87	93	101	114	130	129	109	125	130	119
Mood disorders	266	302	325	341	349	389	407	357	383	403	394
Personality disorders	41	43	46	46	45	48	51	50	53	56	56
Schizophrenia and other psychotic disorders	155	155	166	160	164	155	158	161	164	169	158
Screening and history of mental health and substance abuse	272	297	320	355	391	418	441	372	419	432	426
Substance-related disorders	61	69	83	97	103	103	121	107	119	119	115
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33	N.D.	36	38	37



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Meade County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	906	1,023	1,157	1,254	1,299	1,305	1,295	1,241	1,310	1,339	1,288
Female	413	481	530	582	623	612	603	576	596	599	576
Male	493	542	627	672	676	693	692	665	714	740	712
White	840	948	1,067	1,162	1,199	1,214	1,191	1,143	1,203	1,223	1,183
Minority	N.D.	32	38	34	38	N.D.	N.D.	N.D.	N.D.	33	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	46%	47%	51%	52%	51%	51%	52%	48%	49%	48%	44%
Female	43%	46%	49%	51%	50%	49%	51%	45%	47%	45%	41%
Male	48%	48%	53%	54%	52%	53%	53%	50%	52%	50%	47%
White	46%	47%	51%	52%	51%	51%	51%	47%	48%	47%	44%
Minority	N.D.	21%	23%	21%	21%	N.D.	N.D.	N.D.	N.D.	17%	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	243	292	331	360	391	405	404	383	403	406	384
Alcohol-related disorders	33	39	41	41	37	33	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	228	276	298	315	345	332	324	307	333	323	311
Attention-deficit, conduct, and disruptive behavior disorders	454	491	556	609	645	657	668	628	663	675	639
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	140	163	209	247	272	290	309	311	328	378	375
Disorders usually diagnosed in infancy, childhood, adolescence	68	82	93	96	104	108	115	104	115	123	120
Impulse control disorders, NEC	78	80	86	86	91	80	81	72	72	70	62
Miscellaneous disorders	63	76	85	86	84	84	89	82	90	87	84
Mood disorders	348	367	391	407	409	374	360	313	323	322	285
Personality disorders	47	48	49	56	60	52	46	45	40	41	34
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	535	597	655	692	666	638	602	553	545	519	463
Substance-related disorders	75	82	80	85	85	82	70	60	64	62	52
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Meade County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	527	572	651	734	858	853	913	878	934	906	837
Female	401	431	487	556	636	640	672	656	698	646	603
Male	126	141	164	178	222	213	241	222	236	260	234
White	469	510	590	676	791	788	845	817	869	844	770
Minority	N.D.	31	32	N.D.	34	32	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	49%	51%	56%	58%	62%	58%	60%	56%	59%	59%	54%
Female	51%	54%	60%	62%	66%	62%	64%	60%	63%	61%	57%
Male	42%	44%	47%	49%	53%	48%	51%	47%	50%	53%	48%
White	48%	51%	56%	58%	62%	58%	60%	56%	59%	58%	54%
Minority	N.D.	28%	31%	N.D.	32%	29%	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	43	52	63	66	74	77	86	81	91	92	89
Alcohol-related disorders	43	47	59	68	83	85	92	80	81	84	74
Anxiety disorders	278	303	366	409	453	463	524	495	508	493	454
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	33	36	45	55	57	72	70	77	85	95
Delirium, dementia, and amnestic and other cognitive disorders	116	119	118	117	124	118	120	110	112	102	97
Developmental disorders	41	49	50	52	55	61	69	68	65	65	67
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	82	96	107	134	157	158	162	153	153	151	133
Mood disorders	302	343	402	454	533	539	587	570	608	583	536
Personality disorders	54	62	69	74	81	84	88	74	73	73	65
Schizophrenia and other psychotic disorders	105	115	116	119	123	121	130	117	125	124	110
Screening and history of mental health and substance abuse	250	289	349	412	493	481	520	497	525	519	469
Substance-related disorders	59	77	87	101	114	121	125	107	112	119	104
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Menifee County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	645	697	721	750	758	780	769	726	732	698	653
Female	333	352	358	378	376	376	367	335	346	329	305
Male	312	345	363	372	382	404	402	391	386	369	348
White	605	650	665	694	698	717	705	672	679	637	593
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	66%	66%	66%	66%	67%	66%	64%	61%	62%	57%
Female	69%	71%	71%	69%	69%	66%	65%	60%	59%	58%	53%
Male	57%	61%	63%	64%	63%	68%	67%	67%	64%	66%	61%
White	62%	65%	65%	65%	64%	66%	64%	63%	61%	60%	55%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	131	140	149	161	163	163	165	137	134	122	111
Alcohol-related disorders	43	43	44	42	42	43	42	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	249	274	281	302	314	313	322	297	308	289	275
Attention-deficit, conduct, and disruptive behavior disorders	217	242	250	262	267	272	279	241	240	224	204
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	166	190	213	240	266	281	276	271	274	264	261
Disorders usually diagnosed in infancy, childhood, adolescence	50	54	58	63	65	68	73	69	73	67	65
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	106	109	107	109	113	113	116	104	102	104	99
Mood disorders	224	232	218	208	208	193	183	145	138	116	101
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	500	521	521	532	510	514	503	448	440	410	372
Substance-related disorders	75	75	75	71	67	62	55	45	42	37	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Menifee County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	650	713	764	814	818	848	852	826	878	880	837
Female	440	477	512	552	549	557	575	551	585	583	552
Male	210	236	252	262	269	291	277	275	293	297	285
White	610	667	719	765	772	800	804	780	831	830	783
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	66%	67%	68%	69%	70%	72%	71%	67%	70%	71%	67%
Female	70%	74%	73%	75%	75%	76%	78%	71%	75%	76%	73%
Male	58%	57%	59%	58%	61%	65%	60%	61%	62%	63%	58%
White	65%	67%	68%	68%	70%	71%	71%	66%	69%	71%	67%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	49	58	65	73	72	90	95	98	95	91	87
Alcohol-related disorders	69	76	84	87	96	101	105	107	105	101	92
Anxiety disorders	368	410	462	490	500	528	525	516	539	542	522
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	N.D.	33	33	44	46	57	59	54	52
Delirium, dementia, and amnestic and other cognitive disorders	154	157	147	150	156	149	144	141	144	133	123
Developmental disorders	55	62	71	68	72	69	73	75	82	77	77
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	164	189	206	222	232	241	252	248	255	268	260
Mood disorders	369	406	460	496	507	544	554	529	557	555	532
Personality disorders	59	59	69	74	77	78	86	80	80	85	78
Schizophrenia and other psychotic disorders	133	139	142	147	151	151	149	147	139	133	127
Screening and history of mental health and substance abuse	413	460	518	557	562	600	608	591	629	635	608
Substance-related disorders	94	116	129	141	150	166	180	175	184	189	176
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	31	N.D.	33	31	31	33



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Mercer County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	860	990	1,040	1,151	1,214	1,263	1,272	1,164	1,216	1,247	1,164
Female	430	485	488	535	547	575	583	531	545	559	513
Male	430	505	552	616	667	688	689	633	671	688	651
White	738	854	901	1,002	1,062	1,109	1,101	1,000	1,059	1,083	1,024
Minority	79	91	88	89	87	85	96	88	80	83	73

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	59%	62%	62%	62%	63%	62%	56%	56%	53%	47%
Female	56%	59%	62%	62%	60%	61%	61%	54%	52%	49%	43%
Male	54%	59%	62%	63%	64%	64%	63%	57%	60%	56%	50%
White	53%	58%	61%	61%	61%	62%	61%	54%	55%	52%	46%
Minority	42%	46%	44%	42%	39%	38%	40%	35%	32%	31%	29%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	327	375	397	428	448	462	437	398	421	413	376
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	368	420	448	494	521	540	540	495	541	539	518
Attention-deficit, conduct, and disruptive behavior disorders	426	492	519	587	616	622	610	555	566	564	515
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	179	211	249	291	318	346	368	360	377	390	367
Disorders usually diagnosed in infancy, childhood, adolescence	76	83	98	115	124	132	136	132	132	131	127
Impulse control disorders, NEC	31	41	37	44	39	35	34	31	N.D.	N.D.	N.D.
Miscellaneous disorders	167	180	199	211	220	216	213	194	208	208	185
Mood disorders	270	293	280	303	295	281	272	236	227	217	188
Personality disorders	36	39	40	39	36	32	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	38	38	33	35	33	33	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	466	520	530	567	570	555	522	446	446	417	372
Substance-related disorders	80	81	82	84	79	65	59	49	43	35	N.D.
Suicide and intentional self-inflicted injury	37	38	37	38	38	34	31	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Mercer County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	856	943	1,031	1,126	1,204	1,195	1,244	1,153	1,211	1,245	1,173
Female	617	680	731	798	863	851	889	806	850	883	820
Male	239	263	300	328	341	344	355	347	361	362	353
White	749	833	918	1,016	1,083	1,079	1,126	1,042	1,092	1,121	1,047
Minority	75	75	77	75	83	75	69	60	69	72	71

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	61%	65%	65%	65%	67%	66%	68%	64%	67%	65%	59%
Female	64%	68%	67%	68%	70%	69%	72%	65%	68%	67%	61%
Male	55%	58%	60%	60%	60%	59%	60%	60%	63%	62%	56%
White	61%	65%	65%	66%	67%	66%	68%	63%	67%	65%	58%
Minority	44%	42%	44%	43%	48%	42%	40%	35%	39%	40%	38%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	124	134	157	166	193	214	238	211	243	257	243
Alcohol-related disorders	70	79	91	96	111	112	119	116	117	121	116
Anxiety disorders	460	517	589	643	694	710	767	721	759	785	723
Attention-deficit, conduct, and disruptive behavior disorders	32	40	48	54	71	79	95	95	109	117	120
Delirium, dementia, and amnestic and other cognitive disorders	253	250	250	247	250	238	242	217	209	197	169
Developmental disorders	60	66	64	72	71	78	91	90	93	101	99
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	179	200	235	264	281	297	332	294	311	305	288
Mood disorders	491	567	627	678	726	743	793	729	777	777	724
Personality disorders	109	118	138	154	156	157	170	151	164	171	159
Schizophrenia and other psychotic disorders	225	235	245	246	247	257	274	243	240	230	212
Screening and history of mental health and substance abuse	399	453	548	610	668	658	683	638	682	725	678
Substance-related disorders	131	148	167	184	206	217	226	208	236	248	221
Suicide and intentional self-inflicted injury	N.D.	N.D.	35	37	42	50	57	56	60	64	64



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Metcalfe County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	615	704	764	798	825	838	884	826	882	892	834
Female	303	338	369	378	388	395	418	391	421	432	409
Male	312	366	395	420	437	443	466	435	461	460	425
White	574	656	707	737	763	771	811	763	815	824	769
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	56%	57%	60%	62%	62%	62%	64%	59%	61%	60%	54%
Female	57%	59%	61%	61%	62%	63%	63%	57%	61%	59%	54%
Male	54%	56%	59%	62%	61%	62%	65%	62%	62%	61%	54%
White	55%	56%	58%	60%	60%	61%	63%	59%	60%	59%	53%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	161	180	200	207	210	203	213	193	200	190	175
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	270	303	327	348	367	381	400	391	418	416	397
Attention-deficit, conduct, and disruptive behavior disorders	278	318	354	370	396	385	401	364	384	370	347
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	93	120	145	162	180	194	220	210	230	254	245
Disorders usually diagnosed in infancy, childhood, adolescence	51	62	70	72	76	77	85	80	89	86	71
Impulse control disorders, NEC	52	56	63	61	58	52	51	42	42	36	N.D.
Miscellaneous disorders	71	82	86	86	84	74	77	70	73	68	63
Mood disorders	245	266	274	267	260	251	245	222	222	201	174
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	449	506	529	528	514	489	488	431	436	413	366
Substance-related disorders	52	54	59	56	47	37	36	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Metcalfe County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	837	879	923	925	933	956	980	925	982	1,010	965
Female	553	588	614	608	612	625	652	607	653	667	641
Male	284	291	309	317	321	331	328	318	329	343	324
White	758	794	836	844	849	871	893	838	896	925	886
Minority	38	38	35	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	67%	68%	71%	71%	72%	72%	73%	68%	69%	69%	65%
Female	69%	71%	73%	74%	74%	73%	75%	71%	74%	71%	67%
Male	64%	64%	66%	66%	67%	70%	70%	63%	62%	65%	62%
White	67%	68%	70%	71%	71%	71%	73%	67%	69%	69%	65%
Minority	35%	34%	32%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	76	89	98	106	100	114	112	117	131	131	124
Alcohol-related disorders	62	69	73	72	74	77	85	81	84	86	76
Anxiety disorders	494	536	571	575	588	614	648	626	659	682	639
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	31	N.D.	35	42	53	54	73	79	92	82
Delirium, dementia, and amnestic and other cognitive disorders	212	223	226	214	209	196	205	190	191	172	156
Developmental disorders	104	109	113	109	116	115	120	118	126	115	115
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	31	31	N.D.
Miscellaneous disorders	166	180	199	193	197	203	219	198	207	218	200
Mood disorders	531	578	606	615	631	648	679	638	672	692	658
Personality disorders	57	62	66	70	68	66	64	56	61	63	57
Schizophrenia and other psychotic disorders	236	252	255	252	254	254	255	241	245	220	198
Screening and history of mental health and substance abuse	460	496	534	539	554	578	585	572	593	618	583
Substance-related disorders	81	94	93	93	101	113	114	103	111	120	103
Suicide and intentional self-inflicted injury	N.D.	32	34	N.D.	31	36	47	45	52	55	47



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Monroe County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	752	787	818	827	840	846	862	814	834	788	737
Female	378	382	382	370	374	381	379	367	378	340	316
Male	374	405	436	457	466	465	483	447	456	448	421
White	685	720	744	751	757	766	772	723	736	694	648
Minority	41	39	39	37	40	35	39	40	41	36	36

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	55%	57%	56%	56%	57%	58%	52%	52%	47%	44%
Female	56%	56%	57%	55%	55%	56%	55%	51%	49%	43%	41%
Male	52%	54%	57%	58%	57%	58%	61%	54%	53%	50%	46%
White	53%	55%	57%	56%	55%	56%	57%	51%	50%	45%	42%
Minority	35%	34%	31%	28%	31%	28%	30%	29%	30%	25%	25%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	137	145	158	166	166	172	171	161	154	147	134
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	272	284	288	299	304	305	317	309	309	295	268
Attention-deficit, conduct, and disruptive behavior disorders	252	276	288	303	316	312	322	291	309	300	270
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	158	179	208	227	245	262	271	273	280	273	256
Disorders usually diagnosed in infancy, childhood, adolescence	35	43	44	54	56	54	58	54	57	52	46
Impulse control disorders, NEC	31	35	33	35	40	35	33	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	125	130	125	122	115	111	107	97	91	90	76
Mood disorders	279	281	268	257	248	238	233	200	190	175	150
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	404	408	413	406	393	379	374	330	329	302	277
Substance-related disorders	68	66	61	58	63	55	58	46	44	37	36
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Monroe County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,199	1,249	1,296	1,314	1,317	1,307	1,291	1,208	1,246	1,279	1,216
Female	813	849	868	879	880	871	872	823	828	852	797
Male	386	400	428	435	437	436	419	385	418	427	419
White	1,018	1,071	1,111	1,133	1,142	1,124	1,107	1,040	1,081	1,101	1,049
Minority	109	104	106	95	86	86	85	72	72	83	76

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	70%	72%	74%	73%	73%	72%	73%	67%	69%	71%	66%
Female	72%	75%	77%	77%	76%	76%	76%	70%	73%	73%	68%
Male	66%	66%	68%	67%	67%	66%	66%	61%	64%	66%	62%
White	69%	71%	73%	72%	72%	71%	71%	65%	68%	69%	65%
Minority	46%	44%	44%	42%	38%	38%	38%	33%	36%	37%	32%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	90	103	106	112	115	110	113	105	108	118	117
Alcohol-related disorders	104	113	123	126	126	123	126	118	120	116	119
Anxiety disorders	785	829	871	871	888	892	898	854	872	892	863
Attention-deficit, conduct, and disruptive behavior disorders	34	40	49	48	48	56	59	58	63	72	75
Delirium, dementia, and amnestic and other cognitive disorders	347	355	337	328	312	310	293	274	265	241	216
Developmental disorders	53	55	65	65	61	59	62	59	61	66	71
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	306	320	336	359	358	367	372	339	353	350	345
Mood disorders	749	799	842	872	885	882	882	833	855	892	850
Personality disorders	43	46	49	49	44	46	44	42	40	38	39
Schizophrenia and other psychotic disorders	250	260	268	264	246	252	244	229	221	212	198
Screening and history of mental health and substance abuse	491	528	578	603	600	601	601	575	576	598	590
Substance-related disorders	155	171	196	209	196	191	191	184	180	189	193
Suicide and intentional self-inflicted injury	39	44	49	49	48	51	52	48	47	53	52



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Montgomery County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,544	1,666	1,788	1,821	1,861	1,903	1,985	1,869	1,988	1,990	1,827
Female	800	833	901	898	888	902	929	867	910	903	818
Male	744	833	887	923	973	1,001	1,056	1,002	1,078	1,087	1,009
White	1,420	1,525	1,636	1,660	1,685	1,729	1,786	1,673	1,778	1,781	1,634
Minority	48	58	58	50	53	51	52	49	52	57	50

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	60%	62%	63%	63%	62%	62%	57%	58%	55%	48%
Female	59%	60%	62%	63%	62%	60%	60%	55%	55%	53%	46%
Male	54%	61%	62%	63%	64%	62%	64%	59%	60%	56%	50%
White	56%	59%	61%	62%	62%	60%	61%	55%	57%	53%	47%
Minority	24%	29%	27%	22%	22%	22%	20%	18%	18%	19%	17%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	327	343	385	410	421	442	443	387	397	394	349
Alcohol-related disorders	88	78	83	79	73	66	60	44	43	35	N.D.
Anxiety disorders	636	692	766	765	797	799	835	796	844	840	788
Attention-deficit, conduct, and disruptive behavior disorders	566	620	672	715	733	741	778	708	734	714	639
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	302	334	374	406	451	478	530	528	570	576	544
Disorders usually diagnosed in infancy, childhood, adolescence	160	175	201	224	238	241	258	247	258	254	234
Impulse control disorders, NEC	46	50	57	55	55	52	55	45	43	38	N.D.
Miscellaneous disorders	225	236	253	243	250	256	276	261	266	266	240
Mood disorders	626	634	662	650	646	616	611	526	522	487	420
Personality disorders	56	57	51	50	47	42	44	34	33	32	N.D.
Schizophrenia and other psychotic disorders	67	63	67	70	72	66	70	63	63	59	49
Screening and history of mental health and substance abuse	1,193	1,267	1,354	1,356	1,342	1,326	1,349	1,199	1,232	1,205	1,059
Substance-related disorders	221	208	215	201	193	173	180	137	131	111	87
Suicide and intentional self-inflicted injury	67	66	68	66	67	64	64	56	60	59	46



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Montgomery County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,586	1,713	1,818	1,869	1,920	1,943	1,980	1,895	2,100	2,087	2,048
Female	1,159	1,250	1,313	1,356	1,404	1,420	1,468	1,367	1,534	1,505	1,464
Male	427	463	505	513	516	523	512	528	566	582	584
White	1,425	1,543	1,645	1,690	1,737	1,761	1,791	1,720	1,920	1,912	1,863
Minority	78	83	79	77	76	72	68	58	61	56	58

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	69%	71%	71%	69%	70%	66%	70%	68%	65%
Female	68%	71%	72%	74%	74%	72%	74%	68%	73%	71%	67%
Male	57%	58%	61%	63%	64%	63%	62%	62%	64%	61%	60%
White	65%	67%	69%	71%	71%	70%	70%	66%	71%	68%	65%
Minority	31%	33%	31%	29%	29%	26%	26%	23%	23%	21%	21%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	149	164	177	189	208	212	228	221	246	234	250
Alcohol-related disorders	184	202	213	220	232	219	233	223	238	227	219
Anxiety disorders	917	1,010	1,056	1,120	1,169	1,192	1,249	1,180	1,288	1,296	1,284
Attention-deficit, conduct, and disruptive behavior disorders	59	71	75	88	101	108	130	137	148	173	182
Delirium, dementia, and amnestic and other cognitive disorders	350	344	332	313	305	305	296	275	287	264	225
Developmental disorders	141	157	171	169	179	187	204	203	205	218	224
Disorders usually diagnosed in infancy, childhood, adolescence	33	39	38	35	34	35	46	48	52	58	58
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	36	38	40	38	35	38	41
Miscellaneous disorders	409	443	469	496	519	554	575	539	580	594	580
Mood disorders	924	1,021	1,095	1,148	1,190	1,214	1,276	1,201	1,316	1,307	1,302
Personality disorders	145	174	189	196	205	198	214	193	195	192	194
Schizophrenia and other psychotic disorders	331	326	344	335	343	348	360	339	326	307	296
Screening and history of mental health and substance abuse	1,003	1,133	1,259	1,301	1,352	1,368	1,408	1,355	1,527	1,531	1,515
Substance-related disorders	307	352	370	404	430	450	480	449	505	511	528
Suicide and intentional self-inflicted injury	66	77	80	86	91	91	106	106	113	114	113



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Morgan County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,135	1,211	1,263	1,297	1,277	1,270	1,295	1,245	1,240	1,221	1,131
Female	578	619	628	645	618	623	618	585	578	565	535
Male	557	592	635	652	659	647	677	660	662	656	596
White	1,063	1,131	1,178	1,210	1,188	1,181	1,188	1,144	1,136	1,119	1,037
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	60%	62%	62%	62%	64%	65%	62%	62%	61%	56%
Female	60%	62%	63%	63%	62%	64%	65%	62%	61%	59%	55%
Male	56%	58%	61%	62%	62%	63%	65%	63%	64%	62%	57%
White	57%	59%	61%	61%	60%	62%	63%	60%	60%	59%	54%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	195	203	213	219	224	223	222	208	197	194	181
Alcohol-related disorders	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	458	484	504	512	512	524	534	503	488	463	438
Attention-deficit, conduct, and disruptive behavior disorders	293	310	350	385	396	383	416	376	377	362	332
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	394	437	480	517	540	548	576	559	572	561	520
Disorders usually diagnosed in infancy, childhood, adolescence	66	70	78	90	96	97	101	90	87	83	83
Impulse control disorders, NEC	34	36	35	39	39	35	32	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	166	164	163	154	143	127	127	116	106	103	94
Mood disorders	383	378	377	371	357	340	329	304	293	268	235
Personality disorders	33	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	32	N.D.	N.D.	N.D.	N.D.	N.D.	31	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	768	809	833	841	822	796	785	733	712	688	637
Substance-related disorders	94	94	93	87	81	74	69	57	51	45	37
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Morgan County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,264	1,347	1,398	1,473	1,512	1,529	1,604	1,523	1,581	1,590	1,567
Female	826	875	904	948	978	999	1,041	985	1,014	1,024	994
Male	438	472	494	525	534	530	563	538	567	566	573
White	1,154	1,239	1,288	1,364	1,399	1,422	1,491	1,414	1,474	1,484	1,453
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	65%	66%	67%	68%	68%	70%	67%	69%	70%	68%
Female	68%	70%	70%	72%	73%	74%	76%	71%	75%	74%	72%
Male	57%	58%	60%	61%	61%	59%	62%	60%	61%	63%	61%
White	63%	65%	65%	67%	67%	67%	70%	66%	69%	69%	67%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	67	74	75	87	94	100	112	114	125	123	121
Alcohol-related disorders	105	113	115	125	130	127	154	154	153	156	158
Anxiety disorders	761	825	849	920	967	989	1,064	1,008	1,049	1,037	1,008
Attention-deficit, conduct, and disruptive behavior disorders	48	55	60	66	69	74	75	75	92	86	83
Delirium, dementia, and amnestic and other cognitive disorders	268	266	278	268	254	241	230	212	208	198	172
Developmental disorders	87	94	97	108	109	105	112	111	118	130	135
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	275	293	301	327	345	351	369	331	337	331	325
Mood disorders	685	738	773	843	872	874	951	886	921	920	890
Personality disorders	129	141	146	160	169	164	164	153	153	146	141
Schizophrenia and other psychotic disorders	229	241	248	265	264	257	279	260	258	249	233
Screening and history of mental health and substance abuse	781	853	917	984	1,011	1,035	1,110	1,062	1,123	1,146	1,128
Substance-related disorders	155	164	183	210	229	237	275	256	276	280	283
Suicide and intentional self-inflicted injury	46	51	52	60	62	68	78	74	84	80	74



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Muhlenberg County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,916	2,116	2,216	2,347	2,363	2,349	2,368	2,205	2,182	2,118	2,004
Female	942	1,059	1,070	1,118	1,102	1,089	1,095	1,015	1,009	976	904
Male	974	1,057	1,146	1,229	1,261	1,260	1,273	1,190	1,173	1,142	1,100
White	1,704	1,868	1,951	2,056	2,060	2,027	2,028	1,879	1,857	1,794	1,701
Minority	85	100	99	105	102	100	106	98	99	94	86

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	52%	56%	60%	59%	58%	58%	56%	56%	53%	49%
Female	55%	55%	56%	58%	56%	56%	55%	53%	53%	51%	45%
Male	49%	49%	57%	61%	61%	61%	61%	58%	58%	55%	51%
White	54%	54%	56%	59%	58%	57%	56%	54%	54%	51%	46%
Minority	16%	16%	21%	25%	23%	22%	23%	21%	21%	20%	19%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	399	434	455	474	477	461	471	417	403	378	355
Alcohol-related disorders	60	63	66	62	53	50	46	39	33	N.D.	N.D.
Anxiety disorders	491	538	564	590	588	582	590	539	522	498	451
Attention-deficit, conduct, and disruptive behavior disorders	869	967	1,026	1,122	1,155	1,166	1,173	1,082	1,069	1,021	961
Delirium, dementia, and amnestic and other cognitive disorders	32	35	35	31	34	34	34	33	31	N.D.	N.D.
Developmental disorders	423	504	564	640	692	731	777	785	806	807	780
Disorders usually diagnosed in infancy, childhood, adolescence	87	109	120	136	155	156	171	166	170	177	167
Impulse control disorders, NEC	107	110	118	120	122	117	118	103	98	86	71
Miscellaneous disorders	238	249	271	267	272	270	257	224	225	202	186
Mood disorders	712	764	758	763	750	714	685	605	561	496	429
Personality disorders	N.D.	N.D.	N.D.	31	33	31	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	64	65	66	64	62	62	60	52	49	42	34
Screening and history of mental health and substance abuse	1,149	1,221	1,251	1,266	1,200	1,131	1,106	987	915	862	787
Substance-related disorders	155	168	167	170	158	146	145	130	127	121	110
Suicide and intentional self-inflicted injury	38	39	37	38	35	32	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Muhlenberg County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,826	1,954	2,160	2,246	2,296	2,330	2,446	2,218	2,459	2,451	2,353
Female	1,320	1,393	1,533	1,593	1,628	1,645	1,726	1,537	1,701	1,708	1,625
Male	506	561	627	653	668	685	720	681	758	743	728
White	1,622	1,726	1,923	2,004	2,043	2,084	2,193	1,978	2,218	2,199	2,104
Minority	110	128	129	127	123	110	104	89	93	99	94

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	63%	64%	69%	70%	71%	71%	71%	65%	71%	70%	66%
Female	66%	69%	73%	73%	74%	73%	74%	67%	74%	73%	68%
Male	55%	55%	60%	64%	65%	66%	66%	61%	67%	64%	60%
White	63%	65%	69%	69%	71%	71%	71%	64%	71%	69%	65%
Minority	31%	33%	36%	39%	37%	32%	29%	27%	28%	30%	27%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	206	210	243	252	264	273	298	267	299	292	290
Alcohol-related disorders	162	185	208	230	229	226	232	206	222	217	202
Anxiety disorders	901	965	1,076	1,166	1,220	1,248	1,324	1,248	1,358	1,359	1,304
Attention-deficit, conduct, and disruptive behavior disorders	93	103	114	136	150	170	205	197	244	248	264
Delirium, dementia, and amnesic and other cognitive disorders	393	387	407	401	373	348	354	337	361	332	305
Developmental disorders	128	135	138	140	130	140	151	145	149	146	152
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	31	35	33	31	34	39	39	46	47	54
Miscellaneous disorders	313	354	386	418	449	463	498	479	524	524	491
Mood disorders	1,012	1,101	1,250	1,333	1,403	1,437	1,530	1,433	1,572	1,590	1,549
Personality disorders	58	65	68	69	74	77	79	77	81	76	73
Schizophrenia and other psychotic disorders	374	394	410	414	419	417	417	389	399	382	359
Screening and history of mental health and substance abuse	1,062	1,159	1,328	1,419	1,492	1,528	1,604	1,434	1,593	1,609	1,534
Substance-related disorders	227	258	295	311	331	354	364	336	382	384	355
Suicide and intentional self-inflicted injury	47	59	71	76	86	87	99	89	102	112	109



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Nelson County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,653	1,873	2,031	2,128	2,232	2,298	2,309	2,199	2,326	2,401	2,251
Female	816	905	962	1,006	1,040	1,056	1,059	992	1,033	1,056	976
Male	837	968	1,069	1,122	1,192	1,242	1,250	1,207	1,293	1,345	1,275
White	1,362	1,550	1,665	1,740	1,821	1,873	1,872	1,777	1,894	1,960	1,840
Minority	190	200	216	225	236	246	247	236	233	236	223

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	55%	57%	57%	58%	58%	57%	52%	53%	50%	45%
Female	53%	55%	56%	55%	55%	56%	55%	49%	49%	45%	40%
Male	52%	55%	59%	59%	60%	60%	59%	56%	57%	54%	49%
White	52%	54%	56%	56%	56%	57%	56%	51%	51%	48%	44%
Minority	37%	37%	37%	36%	36%	36%	36%	33%	33%	30%	28%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	405	456	525	550	562	575	573	535	570	573	534
Alcohol-related disorders	71	72	81	81	75	72	64	56	51	42	32
Anxiety disorders	521	583	607	622	632	640	641	606	597	599	544
Attention-deficit, conduct, and disruptive behavior disorders	868	987	1,101	1,188	1,256	1,270	1,256	1,199	1,243	1,262	1,192
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	32	N.D.	N.D.	31	33	35	35	35	34	34
Developmental disorders	333	405	484	529	592	631	665	643	709	736	706
Disorders usually diagnosed in infancy, childhood, adolescence	190	224	253	286	306	320	326	306	318	324	311
Impulse control disorders, NEC	118	133	136	135	128	125	112	105	106	101	87
Miscellaneous disorders	309	338	365	388	426	429	435	417	435	431	394
Mood disorders	667	737	771	783	776	742	709	646	636	623	550
Personality disorders	75	86	78	80	77	70	64	64	57	53	46
Schizophrenia and other psychotic disorders	77	84	86	95	88	86	84	78	74	68	55
Screening and history of mental health and substance abuse	1,019	1,150	1,209	1,205	1,159	1,121	1,078	976	957	913	838
Substance-related disorders	185	192	204	203	189	173	157	145	123	115	91
Suicide and intentional self-inflicted injury	55	55	55	58	58	62	59	53	47	47	40



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Nelson County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,456	1,571	1,726	1,868	2,004	2,097	2,186	2,036	2,283	2,402	2,275
Female	1,072	1,163	1,284	1,387	1,492	1,568	1,649	1,504	1,687	1,757	1,660
Male	384	408	442	481	512	529	537	532	596	645	615
White	1,252	1,358	1,500	1,609	1,732	1,816	1,886	1,759	1,978	2,095	1,956
Minority	139	141	156	177	187	189	201	177	199	202	212

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	60%	62%	65%	66%	68%	68%	63%	69%	68%	62%
Female	60%	63%	64%	67%	69%	71%	71%	64%	70%	70%	64%
Male	53%	52%	55%	59%	59%	63%	62%	60%	64%	63%	57%
White	58%	61%	63%	65%	67%	69%	69%	63%	69%	69%	61%
Minority	37%	37%	38%	42%	45%	44%	44%	40%	43%	42%	42%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	152	163	191	195	212	258	261	254	273	288	287
Alcohol-related disorders	156	162	184	210	223	226	249	229	235	235	224
Anxiety disorders	732	811	909	979	1,063	1,146	1,220	1,170	1,265	1,326	1,276
Attention-deficit, conduct, and disruptive behavior disorders	98	120	135	160	179	193	211	198	246	289	302
Delirium, dementia, and amnestic and other cognitive disorders	325	348	359	357	355	363	367	338	334	321	301
Developmental disorders	151	164	185	189	199	209	215	208	223	240	250
Disorders usually diagnosed in infancy, childhood, adolescence	44	47	50	54	62	64	70	64	72	79	89
Impulse control disorders, NEC	44	49	59	65	76	74	77	77	90	99	97
Miscellaneous disorders	223	261	291	308	357	402	418	388	454	480	458
Mood disorders	838	938	1,044	1,135	1,214	1,282	1,356	1,294	1,397	1,475	1,429
Personality disorders	152	160	176	191	199	200	209	216	225	233	235
Schizophrenia and other psychotic disorders	278	294	313	319	334	343	355	330	342	339	322
Screening and history of mental health and substance abuse	825	916	1,032	1,159	1,284	1,357	1,441	1,323	1,520	1,608	1,502
Substance-related disorders	235	255	289	343	380	390	433	404	445	482	450
Suicide and intentional self-inflicted injury	47	43	57	67	73	79	93	91	110	121	134



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Nicholas County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	554	603	663	657	679	690	723	674	732	717	646
Female	214	241	252	247	258	259	265	242	254	247	229
Male	340	362	411	410	421	431	458	432	478	470	417
White	494	532	583	579	596	607	631	589	645	628	563
Minority	38	42	50	46	48	47	53	50	52	53	42

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	63%	67%	66%	65%	65%	66%	61%	65%	60%	55%
Female	58%	61%	64%	61%	61%	61%	61%	55%	56%	54%	49%
Male	65%	64%	69%	69%	68%	67%	70%	65%	71%	64%	58%
White	60%	61%	65%	63%	62%	62%	64%	59%	63%	58%	52%
Minority	51%	46%	53%	53%	55%	49%	54%	51%	54%	51%	44%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	135	152	165	166	171	168	181	161	171	154	128
Alcohol-related disorders	51	52	57	56	56	58	64	66	61	52	42
Anxiety disorders	203	222	252	262	273	281	290	269	288	272	248
Attention-deficit, conduct, and disruptive behavior disorders	283	313	346	339	357	367	401	376	407	386	331
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	87	104	127	131	135	139	154	152	172	173	163
Disorders usually diagnosed in infancy, childhood, adolescence	42	53	68	70	74	75	76	72	73	78	69
Impulse control disorders, NEC	44	48	56	58	54	52	52	49	54	45	31
Miscellaneous disorders	93	103	107	101	101	109	110	107	113	107	95
Mood disorders	215	238	249	250	248	242	240	228	235	221	176
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	385	419	436	419	414	417	435	413	434	427	359
Substance-related disorders	126	122	131	118	122	129	137	133	136	124	93
Suicide and intentional self-inflicted injury	34	32	33	34	33	N.D.	N.D.	32	31	33	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Nicholas County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	542	593	604	619	640	648	685	598	652	687	655
Female	393	432	433	436	447	452	471	418	452	470	446
Male	149	161	171	183	193	196	214	180	200	217	209
White	508	559	568	584	607	614	651	N.D.	N.D.	N.D.	624
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	67%	70%	67%	70%	69%	70%	73%	63%	68%	66%	62%
Female	70%	73%	70%	72%	73%	73%	74%	66%	72%	71%	66%
Male	60%	62%	60%	65%	63%	65%	69%	58%	60%	58%	55%
White	66%	71%	67%	70%	70%	70%	73%	64%	69%	66%	62%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	35	36	41	47	52	50	54	45	58	64	61
Alcohol-related disorders	46	46	55	56	60	60	65	56	64	67	62
Anxiety disorders	288	312	320	336	354	354	369	338	363	372	359
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	37	39	52	58	55
Delirium, dementia, and amnesic and other cognitive disorders	161	169	161	158	152	157	157	138	127	125	114
Developmental disorders	39	41	36	40	46	47	54	47	53	56	56
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	102	104	109	116	123	117	134	119	131	142	139
Mood disorders	252	281	298	318	332	329	364	331	353	367	363
Personality disorders	49	55	60	58	66	64	70	60	63	67	66
Schizophrenia and other psychotic disorders	136	141	146	140	134	137	144	127	120	118	114
Screening and history of mental health and substance abuse	274	307	331	363	382	381	405	354	398	417	396
Substance-related disorders	100	107	120	129	152	159	173	157	184	184	175
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Ohio County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,357	1,453	1,534	1,616	1,669	1,730	1,780	1,671	1,763	1,744	1,580
Female	688	742	757	787	805	839	857	814	857	815	750
Male	669	711	777	829	864	891	923	857	906	929	830
White	1,261	1,344	1,417	1,483	1,525	1,577	1,620	1,517	1,598	1,579	1,428
Minority	N.D.	N.D.	N.D.	N.D.	31	33	32	N.D.	33	31	31

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	50%	51%	52%	52%	51%	52%	52%	50%	51%	49%	44%
Female	52%	54%	54%	53%	51%	52%	52%	49%	50%	47%	42%
Male	47%	48%	51%	51%	51%	52%	53%	51%	52%	51%	46%
White	49%	50%	51%	51%	50%	51%	51%	48%	49%	48%	43%
Minority	N.D.	N.D.	N.D.	N.D.	15%	15%	15%	N.D.	16%	14%	14%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	435	458	489	517	526	514	503	456	471	455	405
Alcohol-related disorders	35	32	32	33	N.D.	33	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	407	442	466	501	506	519	523	476	475	450	395
Attention-deficit, conduct, and disruptive behavior disorders	612	648	690	738	753	767	751	666	693	660	594
Delirium, dementia, and amnesic and other cognitive disorders	32	N.D.	31	32	35	36	33	N.D.	N.D.	N.D.	N.D.
Developmental disorders	173	202	236	279	297	317	347	329	354	374	343
Disorders usually diagnosed in infancy, childhood, adolescence	76	79	82	101	112	114	119	118	128	125	112
Impulse control disorders, NEC	90	93	95	102	99	99	95	84	84	75	68
Miscellaneous disorders	247	266	259	250	241	229	214	185	176	164	140
Mood disorders	536	557	572	570	541	535	516	447	452	421	369
Personality disorders	39	44	44	38	36	37	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	49	45	47	48	47	47	44	37	35	N.D.	N.D.
Screening and history of mental health and substance abuse	745	793	822	814	822	863	899	859	918	877	785
Substance-related disorders	132	122	123	120	105	99	85	70	66	54	43
Suicide and intentional self-inflicted injury	38	34	38	36	34	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Ohio County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,484	1,612	1,692	1,759	1,828	1,918	2,021	1,824	1,984	1,948	1,871
Female	1,032	1,126	1,189	1,233	1,276	1,346	1,420	1,254	1,376	1,369	1,281
Male	452	486	503	526	552	572	601	570	608	579	590
White	1,373	1,494	1,562	1,626	1,690	1,784	1,879	1,686	1,852	1,798	1,715
Minority	44	46	50	51	41	36	N.D.	31	N.D.	32	31
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	61%	65%	66%	66%	66%	67%	69%	63%	67%	66%	61%
Female	64%	68%	70%	70%	70%	71%	73%	65%	71%	70%	64%
Male	55%	57%	57%	58%	59%	60%	62%	58%	60%	58%	56%
White	61%	64%	65%	65%	66%	67%	69%	62%	67%	66%	61%
Minority	27%	27%	27%	27%	21%	18%	N.D.	17%	N.D.	15%	14%
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	167	203	238	253	242	258	280	254	263	264	251
Alcohol-related disorders	110	119	123	135	143	140	160	138	152	149	145
Anxiety disorders	671	737	787	858	930	979	1,087	1,026	1,098	1,083	1,061
Attention-deficit, conduct, and disruptive behavior disorders	89	100	105	126	141	167	193	195	220	228	248
Delirium, dementia, and amnestic and other cognitive disorders	435	449	449	421	404	397	400	366	369	342	316
Developmental disorders	111	112	128	134	147	153	158	153	156	149	153
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	31	N.D.	32
Impulse control disorders, NEC	N.D.	31	N.D.	31	38	39	50	48	52	47	51
Miscellaneous disorders	363	397	439	476	517	557	591	513	544	531	529
Mood disorders	795	895	948	1,017	1,090	1,145	1,237	1,139	1,225	1,234	1,195
Personality disorders	80	89	92	97	102	113	135	121	122	133	114
Schizophrenia and other psychotic disorders	315	323	333	350	355	366	382	341	344	334	327
Screening and history of mental health and substance abuse	674	744	822	901	963	1,038	1,127	1,031	1,132	1,125	1,063
Substance-related disorders	214	239	276	304	333	342	373	327	355	346	316
Suicide and intentional self-inflicted injury	33	35	43	45	56	63	75	74	80	82	79



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Oldham County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	650	773	837	950	954	1,000	1,082	1,046	1,116	1,173	1,162
Female	309	375	401	450	446	464	500	487	526	542	526
Male	341	398	436	500	508	536	582	559	590	631	636
White	537	652	699	797	798	840	904	878	939	996	982
Minority	80	85	97	99	98	99	104	95	98	96	98

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	50%	52%	53%	54%	51%	50%	50%	45%	46%	42%	39%
Female	48%	50%	51%	52%	48%	47%	47%	43%	44%	40%	36%
Male	52%	54%	55%	56%	54%	52%	53%	47%	47%	45%	41%
White	49%	51%	52%	54%	51%	50%	49%	44%	44%	41%	38%
Minority	39%	38%	39%	37%	32%	31%	35%	32%	31%	28%	26%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	199	231	260	292	299	319	339	331	342	332	321
Alcohol-related disorders	N.D.	N.D.	N.D.	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	204	236	246	276	274	279	291	284	288	291	276
Attention-deficit, conduct, and disruptive behavior disorders	337	393	426	494	503	508	551	533	554	573	555
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	151	183	219	252	264	297	339	346	377	407	422
Disorders usually diagnosed in infancy, childhood, adolescence	77	90	105	116	115	124	139	132	133	146	144
Impulse control disorders, NEC	63	64	64	64	62	58	61	56	55	52	44
Miscellaneous disorders	57	76	75	87	86	89	96	97	103	109	97
Mood disorders	252	294	300	314	296	283	272	248	259	251	232
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	32	35	32	34	39	36	37	35	32
Screening and history of mental health and substance abuse	365	407	417	448	426	415	438	406	422	420	396
Substance-related disorders	64	72	73	77	70	71	67	58	52	58	51
Suicide and intentional self-inflicted injury	36	36	33	36	35	37	36	35	40	40	32



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Oldham County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	699	730	808	850	876	930	1,007	922	1,010	985	932
Female	478	484	542	582	619	648	695	626	698	683	634
Male	221	246	266	268	257	282	312	296	312	302	298
White	614	637	712	751	768	811	870	811	893	881	816
Minority	56	62	66	68	73	83	96	68	73	58	65

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	62%	64%	67%	67%	67%	70%	64%	68%	63%	58%
Female	62%	60%	63%	67%	67%	67%	70%	62%	67%	62%	56%
Male	62%	66%	67%	67%	66%	67%	70%	70%	72%	65%	61%
White	62%	63%	66%	69%	69%	68%	70%	64%	69%	64%	58%
Minority	38%	38%	37%	37%	39%	41%	47%	38%	39%	29%	32%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	65	77	86	79	80	88	87	86	102	94	103
Alcohol-related disorders	59	59	66	66	69	74	87	82	81	74	67
Anxiety disorders	317	344	388	411	450	463	512	494	523	495	454
Attention-deficit, conduct, and disruptive behavior disorders	47	54	61	66	77	86	116	118	135	129	134
Delirium, dementia, and amnesic and other cognitive disorders	237	246	269	278	261	267	294	281	286	279	266
Developmental disorders	141	163	172	171	174	193	228	209	221	230	218
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	31	32	33	34	41	46	45	47	51	50
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	31	35	46	50	47	49	48
Miscellaneous disorders	98	99	122	128	143	152	158	135	144	154	149
Mood disorders	360	395	459	478	505	514	559	526	564	545	502
Personality disorders	47	49	53	53	56	57	66	67	63	57	46
Schizophrenia and other psychotic disorders	248	266	285	283	276	261	276	263	259	257	229
Screening and history of mental health and substance abuse	276	300	347	389	419	441	490	436	486	446	436
Substance-related disorders	103	112	120	132	138	141	155	144	147	138	134
Suicide and intentional self-inflicted injury	N.D.	N.D.	34	34	39	44	55	52	55	51	45



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Owen County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	439	521	574	589	620	667	678	637	659	687	623
Female	210	236	259	271	289	310	310	296	301	310	292
Male	229	285	315	318	331	357	368	341	358	377	331
White	N.D.	N.D.	N.D.	558	585	629	635	594	613	636	576
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	47%	47%	50%	51%	52%	54%	54%	49%	50%	50%	44%
Female	49%	47%	49%	52%	52%	53%	51%	47%	47%	47%	42%
Male	45%	48%	51%	51%	52%	55%	56%	52%	53%	53%	46%
White	47%	48%	50%	51%	51%	53%	53%	48%	48%	48%	43%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	79	92	100	107	106	113	118	106	100	102	87
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	177	182	205	209	217	222	226	198	191	210	184
Attention-deficit, conduct, and disruptive behavior disorders	193	230	266	283	316	331	335	315	324	328	317
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	63	86	98	104	110	122	138	141	140	147	138
Disorders usually diagnosed in infancy, childhood, adolescence	40	46	45	51	54	56	57	53	54	48	44
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	73	78	84	91	88	89	87	71	72	73	61
Mood disorders	148	158	167	166	180	182	173	151	147	144	116
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	253	289	298	302	299	315	306	268	276	266	228
Substance-related disorders	42	44	48	48	48	52	47	38	34	33	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Owen County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	486	569	610	615	613	653	704	641	683	731	665
Female	333	388	407	421	423	451	485	433	474	500	445
Male	153	181	203	194	190	202	219	208	209	231	220
White	452	527	566	566	563	602	656	592	631	680	614
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	71%	70%	70%	70%	71%	75%	67%	72%	72%	65%
Female	69%	75%	73%	74%	74%	75%	76%	67%	74%	76%	67%
Male	58%	62%	65%	64%	63%	63%	71%	69%	68%	66%	61%
White	65%	71%	70%	71%	70%	70%	75%	67%	72%	72%	64%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	39	45	50	49	50	52	69	59	73	83	78
Alcohol-related disorders	31	42	52	52	50	54	70	65	61	63	56
Anxiety disorders	303	348	382	387	406	419	478	448	482	509	469
Attention-deficit, conduct, and disruptive behavior disorders	35	39	43	43	50	56	60	61	72	81	73
Delirium, dementia, and amnestic and other cognitive disorders	161	187	185	177	160	158	159	156	152	146	127
Developmental disorders	54	54	59	62	63	67	70	68	74	73	68
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	87	106	130	146	150	155	160	147	164	171	150
Mood disorders	270	325	361	370	374	390	426	396	426	455	421
Personality disorders	N.D.	35	41	40	40	40	42	46	46	42	39
Schizophrenia and other psychotic disorders	138	149	152	159	154	155	157	152	151	153	133
Screening and history of mental health and substance abuse	260	313	346	362	378	419	455	405	441	477	436
Substance-related disorders	74	94	108	119	120	124	140	139	158	170	161
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33	33	46	49	45



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Owsley County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	674	701	691	681	682	683	681	618	632	610	551
Female	340	349	351	345	331	333	321	283	283	266	236
Male	334	352	340	336	351	350	360	335	349	344	315
White	632	658	643	628	630	625	613	554	567	548	496
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	67%	68%	69%	69%	69%	70%	65%	67%	61%	56%
Female	69%	71%	73%	72%	69%	69%	69%	63%	64%	58%	53%
Male	61%	64%	64%	66%	69%	70%	71%	67%	70%	64%	58%
White	64%	67%	67%	68%	68%	68%	68%	63%	65%	60%	54%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	92	98	98	95	93	82	76	72	72	67	58
Alcohol-related disorders	50	52	49	45	43	40	33	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	349	361	367	363	361	353	344	309	304	285	255
Attention-deficit, conduct, and disruptive behavior disorders	206	222	223	231	237	222	219	199	198	178	159
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	134	141	146	146	149	149	155	146	145	143	133
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	33	34	37	37	41	40	40	39	39	37
Impulse control disorders, NEC	N.D.	33	35	33	34	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	176	177	164	147	134	124	115	94	83	70	60
Mood disorders	231	230	220	210	189	172	160	135	131	125	98
Personality disorders	31	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	438	445	440	441	445	427	408	360	361	355	315
Substance-related disorders	124	119	112	102	91	77	68	59	51	44	34
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Owsley County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,173	1,213	1,269	1,277	1,265	1,284	1,348	1,238	1,288	1,282	1,241
Female	716	736	757	758	757	772	822	741	765	773	740
Male	457	477	512	519	508	512	526	497	523	509	501
White	1,081	1,125	1,182	1,195	1,184	1,202	1,261	1,153	1,205	1,198	1,154
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	72%	75%	79%	77%	77%	77%	81%	76%	79%	79%	75%
Female	79%	80%	84%	82%	83%	82%	86%	79%	83%	84%	80%
Male	64%	68%	72%	71%	70%	70%	73%	71%	73%	73%	69%
White	72%	75%	78%	77%	77%	77%	80%	75%	79%	78%	74%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	112	115	120	126	124	126	147	124	129	134	137
Alcohol-related disorders	147	153	155	155	154	165	177	165	161	169	165
Anxiety disorders	800	846	888	909	913	939	1,003	943	981	989	957
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	32	38	39	38	51	62	57	68	73	76
Delirium, dementia, and amnesic and other cognitive disorders	285	281	271	259	236	229	233	213	207	185	161
Developmental disorders	96	105	112	114	109	112	124	118	124	115	115
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	457	480	528	542	548	562	588	549	562	556	537
Mood disorders	686	704	739	758	768	791	831	779	796	805	785
Personality disorders	86	93	98	106	113	109	116	103	103	104	101
Schizophrenia and other psychotic disorders	194	199	200	202	198	199	204	191	200	196	172
Screening and history of mental health and substance abuse	709	762	807	830	833	865	914	827	874	871	850
Substance-related disorders	331	344	365	385	387	407	440	411	428	428	416
Suicide and intentional self-inflicted injury	49	52	58	57	58	60	75	63	67	61	64



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Pendleton County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	638	737	801	833	828	841	857	809	901	930	880
Female	308	368	394	391	394	399	407	390	435	434	406
Male	330	369	407	442	434	442	450	419	466	496	474
White	599	691	747	771	760	768	774	736	820	848	799
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	59%	61%	61%	58%	58%	58%	54%	57%	54%	48%
Female	54%	59%	61%	59%	57%	57%	57%	53%	57%	53%	47%
Male	55%	58%	61%	63%	59%	60%	59%	55%	58%	55%	50%
White	54%	58%	60%	60%	57%	57%	56%	53%	56%	53%	47%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	126	158	167	158	154	156	150	137	143	137	119
Alcohol-related disorders	N.D.	33	34	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	192	224	241	235	239	235	237	224	246	233	206
Attention-deficit, conduct, and disruptive behavior disorders	297	348	393	407	426	445	456	419	482	498	484
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	115	143	186	203	230	245	266	254	296	340	325
Disorders usually diagnosed in infancy, childhood, adolescence	33	41	46	51	53	51	54	53	56	61	53
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	115	127	133	128	130	118	111	99	107	101	87
Mood disorders	233	252	264	252	251	239	226	200	201	193	162
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	451	505	523	530	502	480	477	427	462	447	405
Substance-related disorders	74	84	82	75	70	68	62	56	61	48	36
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.




Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Pendleton County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	596	658	692	749	817	819	896	826	945	972	933
Female	416	465	483	537	577	570	618	569	665	683	642
Male	180	193	209	212	240	249	278	257	280	289	291
White	562	624	657	705	776	781	849	780	899	924	877
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	69%	71%	72%	73%	71%	75%	68%	73%	72%	67%
Female	68%	72%	72%	76%	76%	75%	78%	70%	77%	75%	69%
Male	58%	62%	67%	64%	66%	64%	69%	63%	66%	67%	63%
White	64%	68%	71%	71%	73%	71%	75%	67%	73%	72%	67%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	47	54	52	61	71	81	99	88	91	105	108
Alcohol-related disorders	56	63	71	92	96	101	118	110	124	116	109
Anxiety disorders	327	371	387	426	482	477	534	508	569	578	557
Attention-deficit, conduct, and disruptive behavior disorders	47	53	61	63	80	95	114	108	110	124	120
Delirium, dementia, and amnesic and other cognitive disorders	135	141	145	147	154	157	163	153	155	136	128
Developmental disorders	68	72	73	75	77	85	94	88	91	87	91
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	126	125	145	180	196	208	244	216	242	248	239
Mood disorders	388	437	454	498	548	562	628	593	675	676	645
Personality disorders	51	56	68	71	81	84	97	78	87	83	86
Schizophrenia and other psychotic disorders	148	151	152	157	173	178	190	186	196	178	170
Screening and history of mental health and substance abuse	372	409	457	508	552	563	626	573	681	704	671
Substance-related disorders	86	96	112	125	145	162	196	183	215	227	207
Suicide and intentional self-inflicted injury	42	42	50	55	59	61	72	69	82	84	80



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Perry County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,008	3,090	3,125	3,189	3,191	3,179	3,177	3,022	3,080	3,059	2,795
Female	1,533	1,539	1,525	1,540	1,513	1,500	1,520	1,426	1,433	1,407	1,286
Male	1,475	1,551	1,600	1,649	1,678	1,679	1,657	1,596	1,647	1,652	1,509
White	2,807	2,870	2,882	2,946	2,938	2,909	2,873	2,733	2,782	2,763	2,521
Minority	78	81	87	84	81	84	86	84	89	82	73

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	59%	61%	61%	62%	63%	65%	61%	62%	60%	54%
Female	61%	61%	61%	61%	61%	62%	63%	60%	60%	57%	52%
Male	55%	58%	60%	62%	64%	64%	66%	63%	64%	62%	56%
White	57%	58%	59%	60%	61%	62%	63%	60%	61%	58%	53%
Minority	29%	28%	27%	26%	25%	25%	25%	24%	25%	22%	19%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	652	683	727	760	769	802	795	758	765	742	672
Alcohol-related disorders	237	235	218	222	206	192	178	166	159	143	113
Anxiety disorders	1,071	1,098	1,093	1,100	1,096	1,061	1,041	967	947	915	824
Attention-deficit, conduct, and disruptive behavior disorders	1,113	1,178	1,229	1,300	1,307	1,319	1,309	1,261	1,265	1,229	1,120
Delirium, dementia, and amnestic and other cognitive disorders	37	38	39	44	46	44	52	53	52	45	37
Developmental disorders	788	875	954	1,022	1,075	1,114	1,162	1,146	1,185	1,170	1,110
Disorders usually diagnosed in infancy, childhood, adolescence	171	188	202	209	221	226	227	217	230	224	206
Impulse control disorders, NEC	179	189	196	200	194	183	175	156	144	133	110
Miscellaneous disorders	409	438	449	462	468	474	472	444	471	470	442
Mood disorders	922	896	881	846	790	751	687	604	573	519	428
Personality disorders	92	94	97	100	94	86	78	70	68	66	52
Schizophrenia and other psychotic disorders	103	102	100	97	91	79	73	65	61	56	45
Screening and history of mental health and substance abuse	2,102	2,145	2,118	2,158	2,141	2,081	2,034	1,885	1,871	1,815	1,621
Substance-related disorders	481	475	441	433	402	386	359	313	309	292	261
Suicide and intentional self-inflicted injury	89	91	81	76	69	60	59	52	46	48	31



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Perry County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,005	4,291	4,394	4,471	4,471	4,430	4,527	4,284	4,551	4,574	4,392
Female	2,578	2,748	2,824	2,834	2,829	2,819	2,897	2,720	2,915	2,925	2,785
Male	1,427	1,543	1,570	1,637	1,642	1,611	1,630	1,564	1,636	1,649	1,607
White	3,693	3,952	4,041	4,113	4,118	4,087	4,181	3,946	4,221	4,250	4,061
Minority	113	127	130	120	111	99	76	74	74	74	76

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	70%	73%	73%	73%	73%	75%	71%	75%	73%	70%
Female	72%	74%	76%	77%	77%	77%	80%	74%	79%	77%	73%
Male	62%	65%	67%	68%	68%	67%	69%	66%	68%	67%	65%
White	68%	70%	72%	73%	73%	72%	75%	71%	75%	73%	69%
Minority	27%	29%	29%	27%	25%	22%	18%	17%	18%	17%	17%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	372	414	434	453	457	468	495	467	505	539	542
Alcohol-related disorders	623	663	708	745	763	768	772	728	752	740	719
Anxiety disorders	2,570	2,747	2,851	2,930	2,955	2,976	3,044	2,939	3,070	3,080	2,963
Attention-deficit, conduct, and disruptive behavior disorders	127	144	164	182	213	225	270	267	307	339	347
Delirium, dementia, and amnestic and other cognitive disorders	655	664	650	623	598	563	570	517	513	487	457
Developmental disorders	367	386	410	430	448	457	490	466	501	504	507
Disorders usually diagnosed in infancy, childhood, adolescence	38	42	45	53	59	62	63	70	76	78	85
Impulse control disorders, NEC	130	134	140	145	151	155	168	160	172	185	191
Miscellaneous disorders	1,061	1,172	1,240	1,292	1,345	1,348	1,368	1,277	1,307	1,287	1,225
Mood disorders	2,313	2,477	2,588	2,667	2,689	2,706	2,805	2,691	2,819	2,817	2,758
Personality disorders	294	312	332	349	369	366	365	325	337	339	322
Schizophrenia and other psychotic disorders	782	822	850	858	858	834	851	770	779	781	731
Screening and history of mental health and substance abuse	2,660	2,887	3,045	3,118	3,178	3,200	3,283	3,110	3,309	3,316	3,164
Substance-related disorders	1,236	1,337	1,433	1,479	1,523	1,577	1,631	1,539	1,612	1,641	1,599
Suicide and intentional self-inflicted injury	168	181	198	205	225	228	257	247	268	268	263



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Pike County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,987	5,241	5,329	5,378	5,407	5,354	5,294	4,922	4,980	4,896	4,532
Female	2,454	2,547	2,574	2,590	2,592	2,530	2,468	2,314	2,316	2,261	2,071
Male	2,533	2,694	2,755	2,788	2,815	2,824	2,826	2,608	2,664	2,635	2,461
White	4,667	4,895	4,954	4,981	4,993	4,922	4,835	4,475	4,527	4,428	4,098
Minority	73	72	70	61	59	53	49	47	45	46	47

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	50%	52%	54%	54%	54%	54%	55%	52%	53%	52%	48%
Female	51%	53%	54%	53%	53%	53%	53%	50%	50%	49%	45%
Male	49%	52%	54%	54%	55%	55%	57%	53%	55%	54%	50%
White	49%	51%	53%	53%	53%	53%	53%	50%	51%	50%	46%
Minority	18%	17%	15%	12%	11%	10%	9%	8%	8%	7%	7%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	611	650	670	704	691	684	681	622	614	588	525
Alcohol-related disorders	196	184	178	161	147	132	117	85	66	53	42
Anxiety disorders	1,611	1,648	1,676	1,698	1,712	1,687	1,685	1,549	1,517	1,456	1,329
Attention-deficit, conduct, and disruptive behavior disorders	1,751	1,879	1,975	2,033	2,063	2,044	2,071	1,901	1,882	1,828	1,682
Delirium, dementia, and amnestic and other cognitive disorders	51	50	51	47	45	50	49	45	41	42	39
Developmental disorders	1,010	1,137	1,248	1,329	1,433	1,501	1,547	1,507	1,562	1,550	1,491
Disorders usually diagnosed in infancy, childhood, adolescence	279	304	322	342	366	379	401	387	379	370	350
Impulse control disorders, NEC	83	86	88	86	81	80	78	66	64	60	47
Miscellaneous disorders	1,011	1,032	1,028	1,040	1,032	1,014	990	911	894	859	783
Mood disorders	1,661	1,635	1,594	1,533	1,449	1,363	1,272	1,110	1,025	949	807
Personality disorders	61	64	70	65	57	60	53	42	41	39	31
Schizophrenia and other psychotic disorders	149	154	159	154	144	139	133	106	101	97	80
Screening and history of mental health and substance abuse	3,162	3,274	3,326	3,289	3,208	3,108	3,030	2,739	2,697	2,564	2,305
Substance-related disorders	607	557	522	479	435	397	356	335	313	295	263
Suicide and intentional self-inflicted injury	124	113	117	114	105	105	96	87	75	68	54



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Pike County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	6,535	6,898	7,253	7,467	7,681	7,767	7,872	7,390	7,894	7,972	7,732
Female	4,348	4,579	4,783	4,935	5,071	5,161	5,208	4,897	5,230	5,263	5,083
Male	2,187	2,319	2,470	2,532	2,610	2,606	2,664	2,493	2,664	2,709	2,649
White	6,010	6,358	6,692	6,904	7,109	7,186	7,288	6,805	7,295	7,390	7,149
Minority	142	135	132	116	95	76	46	36	41	40	41

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	65%	67%	70%	71%	72%	72%	73%	68%	72%	72%	69%
Female	68%	70%	73%	74%	75%	75%	76%	71%	76%	75%	72%
Male	60%	62%	65%	66%	67%	65%	67%	63%	66%	66%	64%
White	64%	66%	69%	71%	71%	71%	72%	67%	71%	72%	69%
Minority	21%	19%	19%	17%	14%	11%	7%	5%	6%	5%	5%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	343	374	399	406	433	447	465	426	465	474	460
Alcohol-related disorders	718	783	826	856	872	878	921	903	946	928	857
Anxiety disorders	4,217	4,522	4,765	4,985	5,127	5,195	5,313	5,046	5,289	5,298	5,112
Attention-deficit, conduct, and disruptive behavior disorders	154	188	204	223	256	283	322	319	402	439	468
Delirium, dementia, and amnestic and other cognitive disorders	956	959	961	939	916	904	881	802	805	782	708
Developmental disorders	390	410	423	437	445	454	463	470	499	511	510
Disorders usually diagnosed in infancy, childhood, adolescence	35	42	42	43	46	45	46	51	64	71	73
Impulse control disorders, NEC	46	48	53	60	64	68	69	68	71	73	79
Miscellaneous disorders	2,113	2,284	2,465	2,667	2,787	2,819	2,850	2,652	2,720	2,697	2,565
Mood disorders	4,246	4,558	4,887	5,083	5,216	5,297	5,401	5,159	5,421	5,402	5,234
Personality disorders	254	286	293	307	314	312	321	307	306	302	303
Schizophrenia and other psychotic disorders	994	1,056	1,076	1,088	1,083	1,088	1,076	997	993	974	918
Screening and history of mental health and substance abuse	4,256	4,582	4,960	5,243	5,458	5,527	5,736	5,532	5,952	5,996	5,829
Substance-related disorders	1,490	1,641	1,752	1,869	1,998	2,082	2,228	2,180	2,367	2,453	2,405
Suicide and intentional self-inflicted injury	249	274	294	295	323	346	380	379	389	388	383



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Powell County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,147	1,241	1,300	1,389	1,455	1,456	1,481	1,395	1,437	1,403	1,298
Female	588	617	640	680	724	705	737	659	686	657	620
Male	559	624	660	709	731	751	744	736	751	746	678
White	1,068	1,148	1,200	1,278	1,339	1,332	1,351	1,278	1,315	1,286	1,185
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	61%	62%	64%	63%	63%	64%	61%	60%	59%	53%
Female	62%	63%	62%	64%	63%	62%	65%	60%	59%	56%	51%
Male	54%	59%	62%	63%	63%	65%	63%	62%	62%	63%	55%
White	57%	60%	61%	62%	62%	62%	63%	60%	59%	58%	52%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	364	393	393	428	450	453	449	422	433	398	358
Alcohol-related disorders	47	47	49	47	39	34	33	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	468	506	548	578	613	618	625	591	596	588	557
Attention-deficit, conduct, and disruptive behavior disorders	384	419	449	504	536	538	538	515	516	481	436
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	191	220	260	301	322	346	371	348	377	382	355
Disorders usually diagnosed in infancy, childhood, adolescence	98	116	128	145	155	156	161	154	152	145	137
Impulse control disorders, NEC	N.D.	N.D.	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	255	261	288	316	325	326	332	319	321	299	276
Mood disorders	394	387	388	377	394	374	355	306	295	265	224
Personality disorders	32	32	N.D.	32	39	36	35	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	41	39	40	45	43	41	41	35	35	N.D.	N.D.
Screening and history of mental health and substance abuse	712	748	751	763	761	743	714	627	614	551	501
Substance-related disorders	140	132	135	122	121	103	99	76	82	73	64
Suicide and intentional self-inflicted injury	39	35	38	41	39	35	31	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Powell County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,287	1,414	1,488	1,590	1,641	1,713	1,749	1,615	1,755	1,696	1,651
Female	877	985	1,032	1,098	1,133	1,181	1,208	1,101	1,191	1,138	1,087
Male	410	429	456	492	508	532	541	514	564	558	564
White	1,182	1,308	1,374	1,481	1,531	1,599	1,632	1,507	1,642	1,577	1,525
Minority	32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	69%	72%	74%	76%	75%	75%	77%	70%	76%	72%	68%
Female	73%	76%	78%	79%	78%	79%	80%	73%	78%	75%	71%
Male	63%	64%	67%	69%	69%	69%	71%	65%	71%	66%	64%
White	69%	72%	74%	75%	75%	75%	77%	70%	76%	72%	68%
Minority	23%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	212	234	258	289	304	322	338	335	347	343	347
Alcohol-related disorders	132	139	152	169	176	183	189	179	196	182	178
Anxiety disorders	777	860	933	999	1,048	1,087	1,126	1,057	1,145	1,095	1,095
Attention-deficit, conduct, and disruptive behavior disorders	60	72	73	81	82	89	99	94	104	104	102
Delirium, dementia, and amnestic and other cognitive disorders	248	252	247	244	233	224	216	193	190	174	152
Developmental disorders	107	106	115	119	121	135	137	140	138	134	139
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	381	423	451	498	519	547	578	530	559	534	513
Mood disorders	826	900	965	1,055	1,094	1,140	1,158	1,097	1,162	1,110	1,093
Personality disorders	119	133	142	154	161	168	179	169	180	169	162
Schizophrenia and other psychotic disorders	259	276	291	298	295	303	301	277	276	265	237
Screening and history of mental health and substance abuse	805	895	975	1,066	1,122	1,183	1,238	1,138	1,234	1,196	1,156
Substance-related disorders	267	300	326	354	374	407	431	403	441	415	410
Suicide and intentional self-inflicted injury	50	55	59	68	69	76	90	80	85	87	86



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Pulaski County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,869	4,281	4,472	4,714	4,790	4,824	4,957	4,696	4,829	4,737	4,363
Female	1,936	2,122	2,210	2,287	2,304	2,311	2,359	2,214	2,294	2,265	2,067
Male	1,933	2,159	2,262	2,427	2,486	2,513	2,598	2,482	2,535	2,472	2,296
White	3,586	3,965	4,124	4,336	4,366	4,379	4,471	4,220	4,320	4,238	3,894
Minority	98	103	106	103	112	112	115	121	134	133	127

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	58%	59%	59%	59%	59%	59%	56%	56%	53%	47%
Female	58%	59%	59%	58%	58%	58%	57%	53%	54%	50%	45%
Male	56%	58%	59%	60%	60%	61%	61%	58%	59%	55%	49%
White	56%	57%	58%	58%	58%	58%	58%	55%	54%	51%	46%
Minority	26%	24%	23%	20%	21%	19%	18%	18%	20%	18%	17%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	764	844	875	889	889	884	874	790	773	715	616
Alcohol-related disorders	210	202	193	187	169	151	143	111	98	76	56
Anxiety disorders	1,816	1,996	2,119	2,216	2,248	2,262	2,321	2,185	2,215	2,119	1,953
Attention-deficit, conduct, and disruptive behavior disorders	2,104	2,306	2,410	2,512	2,530	2,500	2,487	2,259	2,231	2,141	1,911
Delirium, dementia, and amnesic and other cognitive disorders	67	70	71	71	66	65	66	64	61	54	45
Developmental disorders	773	861	909	1,000	1,053	1,116	1,209	1,220	1,319	1,334	1,257
Disorders usually diagnosed in infancy, childhood, adolescence	337	389	417	433	450	450	457	408	419	416	382
Impulse control disorders, NEC	263	273	272	274	249	229	229	178	166	149	116
Miscellaneous disorders	497	522	521	510	503	476	482	413	397	351	305
Mood disorders	1,651	1,745	1,742	1,706	1,623	1,511	1,451	1,263	1,190	1,078	898
Personality disorders	121	123	111	105	95	94	86	69	62	62	40
Schizophrenia and other psychotic disorders	186	187	186	178	168	159	161	135	130	122	99
Screening and history of mental health and substance abuse	2,067	2,232	2,233	2,216	2,171	2,049	2,038	1,845	1,813	1,707	1,510
Substance-related disorders	539	533	520	495	454	415	386	314	281	244	187
Suicide and intentional self-inflicted injury	178	177	172	160	149	146	153	139	141	134	104



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Pulaski County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	5,132	5,434	5,664	5,783	5,983	6,109	6,349	6,026	6,427	6,485	6,291
Female	3,468	3,652	3,771	3,856	3,969	4,061	4,228	4,022	4,291	4,268	4,060
Male	1,664	1,782	1,893	1,927	2,014	2,048	2,121	2,004	2,136	2,217	2,231
White	4,719	4,997	5,229	5,338	5,541	5,660	5,879	5,592	5,991	6,048	5,825
Minority	163	170	156	157	139	134	129	109	116	114	127

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	69%	72%	70%	70%	70%	72%	68%	70%	69%	66%
Female	71%	72%	73%	72%	72%	72%	74%	69%	72%	71%	67%
Male	63%	65%	68%	67%	68%	67%	69%	65%	66%	65%	63%
White	68%	69%	71%	70%	70%	70%	72%	67%	70%	69%	66%
Minority	28%	29%	27%	26%	23%	21%	21%	18%	20%	18%	17%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	448	502	528	578	588	621	660	636	710	740	729
Alcohol-related disorders	447	506	530	547	594	583	599	569	593	614	590
Anxiety disorders	3,290	3,516	3,712	3,860	4,007	4,121	4,342	4,184	4,454	4,514	4,368
Attention-deficit, conduct, and disruptive behavior disorders	342	382	420	473	511	556	623	642	727	783	800
Delirium, dementia, and amnesic and other cognitive disorders	1,122	1,133	1,132	1,087	1,056	1,023	1,002	953	931	888	834
Developmental disorders	779	789	805	816	840	851	894	848	877	885	901
Disorders usually diagnosed in infancy, childhood, adolescence	186	193	206	210	214	219	230	219	238	260	263
Impulse control disorders, NEC	161	182	193	212	229	231	256	258	274	287	295
Miscellaneous disorders	1,041	1,131	1,225	1,285	1,334	1,387	1,470	1,419	1,512	1,501	1,465
Mood disorders	3,309	3,526	3,689	3,827	4,000	4,072	4,276	4,112	4,377	4,423	4,300
Personality disorders	423	439	455	461	480	474	491	455	487	466	472
Schizophrenia and other psychotic disorders	1,068	1,103	1,156	1,153	1,148	1,104	1,120	1,064	1,085	1,052	1,024
Screening and history of mental health and substance abuse	2,822	3,086	3,281	3,427	3,620	3,734	3,960	3,760	4,025	4,060	3,922
Substance-related disorders	799	878	935	981	1,064	1,092	1,199	1,141	1,255	1,301	1,299
Suicide and intentional self-inflicted injury	287	296	322	335	353	374	403	400	447	462	457



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Robertson County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	146	149	157	151	162	170	175	167	173	170	157
Female	80	79	78	80	82	82	84	82	79	76	76
Male	66	70	79	71	80	88	91	85	94	94	81
White	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	59%	59%	58%	58%	59%	62%	63%	61%	61%	56%	51%
Female	63%	62%	61%	62%	61%	65%	64%	61%	61%	54%	49%
Male	55%	56%	55%	54%	57%	60%	62%	61%	62%	58%	54%
White	61%	59%	58%	57%	58%	61%	61%	58%	59%	55%	50%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	35	36	37	35	36	40	43	50	55	51	46
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	40	38	42	40	46	47	50	49	46	44	40
Attention-deficit, conduct, and disruptive behavior disorders	62	66	77	77	85	91	91	83	79	74	67
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	44	53	62	58	65	67	69	67	71	74	70
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	38	37	37	39	42	36	39	37	32	35	N.D.
Mood disorders	56	53	52	51	50	43	45	39	36	36	N.D.
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	96	100	108	100	112	111	108	101	102	96	82
Substance-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Robertson County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	178	191	195	197	201	203	232	226	232	220	231
Female	121	132	132	136	144	143	163	159	157	151	149
Male	57	59	63	61	57	60	69	67	75	69	82
White	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	#N/A	#N/A	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	66%	68%	69%	69%	68%	74%	71%	70%	65%	65%
Female	65%	68%	71%	74%	76%	74%	77%	74%	74%	71%	70%
Male	61%	61%	64%	61%	57%	57%	68%	64%	64%	55%	57%
White	63%	64%	67%	68%	68%	67%	72%	69%	68%	64%	64%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	#N/A	#N/A	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	83	89	92	92	104	110	123	128	129	130	131
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31
Delirium, dementia, and amnestic and other cognitive disorders	69	73	69	66	70	75	78	72	75	70	66
Developmental disorders	N.D.	N.D.	N.D.	31	33	39	45	42	41	44	42
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	34	35	40	36	40	49	49	48	53	54	55
Mood disorders	89	100	104	106	114	122	134	136	138	132	139
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	31	35	33	36	36	N.D.
Schizophrenia and other psychotic disorders	63	68	69	69	71	76	77	75	76	72	66
Screening and history of mental health and substance abuse	91	104	114	119	125	127	141	142	146	143	144
Substance-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	34	38	39	40	44	49
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Rockcastle County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,133	1,262	1,311	1,334	1,385	1,404	1,469	1,449	1,518	1,520	1,401
Female	573	615	630	631	647	654	706	693	711	717	667
Male	560	647	681	703	738	750	763	756	807	803	734
White	1,083	1,202	1,245	1,263	1,304	1,321	1,383	1,366	1,429	1,430	1,317
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	54%	56%	57%	59%	60%	60%	61%	58%	59%	57%	52%
Female	56%	56%	57%	58%	58%	58%	60%	57%	57%	54%	50%
Male	52%	56%	58%	60%	62%	62%	61%	60%	62%	60%	54%
White	53%	55%	56%	58%	59%	59%	60%	57%	58%	56%	51%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	230	243	244	244	236	231	235	217	204	189	153
Alcohol-related disorders	69	76	72	70	70	63	60	56	50	38	N.D.
Anxiety disorders	500	567	579	598	627	645	687	677	685	664	620
Attention-deficit, conduct, and disruptive behavior disorders	423	495	518	534	572	578	579	559	573	564	515
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	358	402	453	488	555	571	599	593	649	658	631
Disorders usually diagnosed in infancy, childhood, adolescence	73	84	93	96	104	106	113	103	101	105	104
Impulse control disorders, NEC	N.D.	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	158	160	172	173	183	194	197	195	184	182	167
Mood disorders	381	398	377	357	342	321	306	292	270	251	213
Personality disorders	68	65	64	64	61	57	56	53	40	41	39
Schizophrenia and other psychotic disorders	40	43	43	44	40	34	31	31	32	N.D.	N.D.
Screening and history of mental health and substance abuse	566	610	605	601	597	587	581	559	560	570	510
Substance-related disorders	172	180	170	149	147	129	119	112	97	85	69
Suicide and intentional self-inflicted injury	38	39	35	39	34	32	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Rockcastle County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,497	1,615	1,660	1,682	1,730	1,745	1,899	1,785	1,901	1,942	1,863
Female	1,025	1,099	1,134	1,138	1,171	1,172	1,281	1,195	1,265	1,278	1,202
Male	472	516	526	544	559	573	618	590	636	664	661
White	1,374	1,489	1,538	1,559	1,604	1,612	1,756	1,650	1,771	1,810	1,731
Minority	36	36	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	67%	68%	69%	71%	71%	69%	71%	66%	69%	67%	64%
Female	71%	72%	73%	75%	74%	72%	76%	70%	74%	71%	68%
Male	59%	62%	61%	64%	65%	63%	63%	59%	62%	61%	59%
White	66%	67%	68%	70%	70%	68%	70%	65%	69%	67%	64%
Minority	21%	22%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	175	202	213	222	227	240	282	273	286	307	295
Alcohol-related disorders	139	151	152	162	186	187	201	184	199	200	176
Anxiety disorders	943	1,027	1,071	1,101	1,135	1,155	1,262	1,215	1,278	1,303	1,250
Attention-deficit, conduct, and disruptive behavior disorders	76	82	94	105	112	119	131	114	131	136	154
Delirium, dementia, and amnesic and other cognitive disorders	364	369	364	354	336	320	331	294	277	260	227
Developmental disorders	170	174	192	192	203	210	228	206	210	219	208
Disorders usually diagnosed in infancy, childhood, adolescence	42	43	44	45	47	49	49	43	44	48	46
Impulse control disorders, NEC	51	53	54	57	59	60	65	57	57	50	53
Miscellaneous disorders	320	360	385	395	423	427	453	442	466	483	452
Mood disorders	828	904	939	968	1,002	1,016	1,127	1,076	1,148	1,177	1,136
Personality disorders	109	124	134	141	154	154	175	159	164	161	158
Schizophrenia and other psychotic disorders	329	348	349	355	355	338	366	330	333	332	310
Screening and history of mental health and substance abuse	739	810	875	889	928	960	1,073	1,023	1,116	1,129	1,083
Substance-related disorders	257	307	332	351	370	377	422	394	434	455	409
Suicide and intentional self-inflicted injury	43	53	57	56	60	66	89	84	93	92	81



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Rowan County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,287	1,384	1,451	1,512	1,542	1,547	1,614	1,524	1,574	1,566	1,480
Female	670	712	718	738	749	732	765	716	722	701	651
Male	617	672	733	774	793	815	849	808	852	865	829
White	1,199	1,287	1,337	1,388	1,404	1,404	1,454	1,382	1,423	1,412	1,333
Minority	N.D.	N.D.	38	37	38	38	38	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	58%	60%	62%	61%	60%	61%	57%	57%	55%	51%
Female	60%	60%	60%	63%	61%	58%	59%	54%	55%	52%	48%
Male	55%	57%	60%	62%	62%	62%	62%	59%	59%	59%	54%
White	57%	58%	59%	61%	60%	59%	59%	55%	55%	54%	50%
Minority	N.D.	N.D.	26%	24%	22%	20%	20%	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	287	317	341	347	357	357	369	328	325	317	294
Alcohol-related disorders	54	62	54	54	50	44	39	33	N.D.	N.D.	N.D.
Anxiety disorders	495	520	529	568	599	591	624	598	596	588	548
Attention-deficit, conduct, and disruptive behavior disorders	534	580	613	642	661	675	721	671	674	665	620
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	330	379	426	462	477	504	539	535	543	563	557
Disorders usually diagnosed in infancy, childhood, adolescence	117	134	144	166	182	188	207	184	183	173	167
Impulse control disorders, NEC	59	62	67	70	70	72	76	64	63	63	57
Miscellaneous disorders	170	177	178	184	191	190	186	171	157	153	137
Mood disorders	551	580	569	581	580	553	555	477	458	422	366
Personality disorders	59	55	57	57	54	45	47	39	36	32	N.D.
Schizophrenia and other psychotic disorders	51	49	54	54	56	55	56	44	43	43	42
Screening and history of mental health and substance abuse	1,036	1,080	1,123	1,140	1,132	1,084	1,113	1,018	1,035	1,000	929
Substance-related disorders	152	148	142	142	132	116	113	106	89	77	54
Suicide and intentional self-inflicted injury	40	43	40	44	44	39	42	35	33	33	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Rowan County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,591	1,701	1,759	1,815	1,889	1,940	2,024	1,903	2,016	1,991	1,917
Female	1,068	1,115	1,176	1,223	1,257	1,293	1,355	1,272	1,342	1,311	1,257
Male	523	586	583	592	632	647	669	631	674	680	660
White	1,447	1,553	1,598	1,638	1,711	1,764	1,855	1,734	1,838	1,819	1,753
Minority	41	36	40	46	44	35	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	71%	73%	73%	74%	74%	73%	76%	71%	73%	70%	67%
Female	72%	75%	76%	77%	76%	76%	79%	73%	76%	73%	69%
Male	68%	70%	68%	70%	71%	69%	71%	67%	68%	66%	64%
White	70%	73%	73%	74%	74%	73%	75%	70%	72%	70%	67%
Minority	21%	19%	20%	22%	20%	16%	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	149	169	188	196	214	226	249	245	258	241	236
Alcohol-related disorders	208	228	243	242	264	263	279	264	272	255	244
Anxiety disorders	962	1,043	1,105	1,167	1,221	1,250	1,320	1,252	1,310	1,282	1,240
Attention-deficit, conduct, and disruptive behavior disorders	74	74	88	95	113	121	138	136	156	170	178
Delirium, dementia, and amnesic and other cognitive disorders	414	417	416	414	392	371	358	330	307	288	246
Developmental disorders	277	292	302	326	348	354	358	331	326	337	335
Disorders usually diagnosed in infancy, childhood, adolescence	40	43	47	47	50	56	63	58	62	67	66
Impulse control disorders, NEC	63	71	72	73	78	88	88	84	90	99	92
Miscellaneous disorders	355	385	410	423	444	465	498	462	484	473	445
Mood disorders	984	1,063	1,110	1,166	1,253	1,296	1,369	1,297	1,356	1,333	1,280
Personality disorders	237	265	282	290	307	310	323	299	309	291	296
Schizophrenia and other psychotic disorders	345	355	361	364	363	361	392	363	362	355	335
Screening and history of mental health and substance abuse	934	1,053	1,124	1,180	1,249	1,307	1,387	1,333	1,457	1,456	1,417
Substance-related disorders	279	316	336	354	388	412	438	435	459	459	452
Suicide and intentional self-inflicted injury	52	56	58	59	69	78	90	90	95	93	104



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Russell County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,357	1,503	1,553	1,629	1,668	1,673	1,690	1,588	1,639	1,612	1,499
Female	695	760	790	823	856	848	856	794	821	800	728
Male	662	743	763	806	812	825	834	794	818	812	771
White	1,281	1,410	1,455	1,511	1,547	1,549	1,554	1,451	1,498	1,464	1,362
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	63%	65%	66%	66%	66%	66%	62%	61%	58%	53%
Female	63%	65%	66%	66%	67%	67%	65%	62%	61%	58%	52%
Male	58%	62%	64%	65%	65%	66%	67%	62%	61%	58%	54%
White	60%	63%	65%	65%	65%	65%	65%	60%	60%	56%	52%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	289	313	307	311	325	313	303	276	273	260	233
Alcohol-related disorders	73	71	74	68	63	53	49	44	37	32	N.D.
Anxiety disorders	658	724	754	794	809	823	823	750	760	726	673
Attention-deficit, conduct, and disruptive behavior disorders	653	701	717	757	765	784	762	684	708	680	625
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	207	246	276	314	354	377	410	403	442	455	448
Disorders usually diagnosed in infancy, childhood, adolescence	82	91	100	106	113	115	113	109	111	107	103
Impulse control disorders, NEC	97	98	95	99	97	91	84	68	63	50	45
Miscellaneous disorders	259	270	288	273	272	258	252	224	213	196	173
Mood disorders	590	593	590	579	553	512	481	399	375	349	304
Personality disorders	41	38	44	43	43	36	34	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	53	51	55	57	55	50	47	42	40	35	33
Screening and history of mental health and substance abuse	725	773	790	791	758	710	671	609	611	600	549
Substance-related disorders	212	201	189	178	158	139	121	96	86	72	61
Suicide and intentional self-inflicted injury	45	48	48	45	45	42	35	31	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Russell County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,775	1,903	1,958	2,011	2,040	2,033	2,045	1,930	2,094	2,116	1,999
Female	1,205	1,283	1,313	1,352	1,361	1,387	1,395	1,307	1,424	1,428	1,347
Male	570	620	645	659	679	646	650	623	670	688	652
White	1,647	1,778	1,828	1,886	1,909	1,910	1,913	1,808	1,964	1,987	1,875
Minority	43	38	33	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	71%	73%	73%	73%	73%	73%	69%	72%	72%	68%
Female	71%	74%	76%	76%	75%	76%	77%	71%	74%	74%	70%
Male	63%	67%	69%	69%	69%	67%	67%	66%	68%	67%	64%
White	68%	71%	73%	73%	73%	73%	73%	69%	72%	72%	67%
Minority	23%	21%	20%	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	188	205	219	236	259	268	292	281	313	320	304
Alcohol-related disorders	157	167	176	171	185	197	196	184	202	220	214
Anxiety disorders	1,213	1,308	1,371	1,415	1,449	1,458	1,499	1,439	1,545	1,551	1,488
Attention-deficit, conduct, and disruptive behavior disorders	41	54	63	72	78	91	122	131	162	185	187
Delirium, dementia, and amnesic and other cognitive disorders	459	472	463	453	427	405	392	369	365	328	288
Developmental disorders	155	151	162	171	171	175	184	181	183	191	177
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	36	41	42	42	47	55	57	66	77	72
Miscellaneous disorders	439	480	505	547	561	568	581	550	598	611	570
Mood disorders	1,153	1,248	1,295	1,342	1,378	1,398	1,428	1,354	1,458	1,477	1,412
Personality disorders	156	166	168	166	170	167	182	175	187	181	180
Schizophrenia and other psychotic disorders	448	468	465	463	448	435	439	428	429	405	375
Screening and history of mental health and substance abuse	827	910	971	1,018	1,055	1,080	1,108	1,033	1,111	1,137	1,069
Substance-related disorders	322	354	372	394	416	430	446	422	488	517	521
Suicide and intentional self-inflicted injury	52	60	61	63	68	75	82	77	92	95	98



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Scott County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,403	1,583	1,674	1,789	1,888	1,938	2,003	1,954	2,150	2,221	2,123
Female	713	768	823	872	901	907	928	903	1,003	1,032	957
Male	690	815	851	917	987	1,031	1,075	1,051	1,147	1,189	1,166
White	1,231	1,382	1,456	1,544	1,631	1,668	1,710	1,664	1,834	1,911	1,820
Minority	118	138	150	166	165	170	181	178	197	191	190

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	56%	57%	58%	57%	57%	57%	53%	54%	49%	44%
Female	57%	57%	58%	59%	56%	55%	54%	49%	51%	46%	41%
Male	53%	56%	57%	58%	58%	59%	60%	56%	57%	52%	47%
White	56%	56%	58%	59%	57%	57%	56%	52%	53%	49%	44%
Minority	34%	38%	39%	39%	38%	38%	38%	35%	37%	30%	28%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	357	380	411	454	473	461	459	408	423	421	402
Alcohol-related disorders	72	81	79	88	79	76	70	61	58	55	47
Anxiety disorders	511	588	633	681	726	766	810	795	854	847	826
Attention-deficit, conduct, and disruptive behavior disorders	697	783	828	887	952	965	970	908	958	975	921
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	32	34	34	N.D.	N.D.	N.D.
Developmental disorders	236	296	327	369	415	460	504	546	596	612	588
Disorders usually diagnosed in infancy, childhood, adolescence	147	170	192	210	220	235	259	248	264	260	265
Impulse control disorders, NEC	55	52	53	53	55	54	48	38	35	33	N.D.
Miscellaneous disorders	236	251	255	265	269	259	272	256	272	275	252
Mood disorders	522	554	552	561	565	551	515	481	500	498	454
Personality disorders	51	54	56	56	56	55	45	35	37	34	45
Schizophrenia and other psychotic disorders	64	72	66	71	76	79	72	72	69	66	64
Screening and history of mental health and substance abuse	860	914	909	922	911	868	841	751	765	767	706
Substance-related disorders	187	204	188	180	173	156	149	135	135	126	104
Suicide and intentional self-inflicted injury	59	70	73	69	60	60	57	54	55	52	54



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Scott County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,184	1,326	1,450	1,583	1,662	1,755	1,854	1,729	1,952	1,998	1,942
Female	931	1,042	1,129	1,241	1,287	1,347	1,411	1,319	1,460	1,467	1,404
Male	253	284	321	342	375	408	443	410	492	531	538
White	1,016	1,139	1,253	1,373	1,453	1,540	1,628	1,520	1,736	1,766	1,701
Minority	110	122	127	133	127	131	133	120	124	130	127

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	68%	69%	69%	71%	72%	65%	70%	69%	65%
Female	67%	70%	71%	72%	70%	72%	73%	66%	71%	69%	65%
Male	55%	57%	59%	62%	65%	65%	70%	62%	68%	68%	65%
White	65%	68%	69%	71%	71%	72%	74%	66%	72%	70%	66%
Minority	39%	41%	40%	39%	36%	37%	37%	32%	34%	34%	31%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	117	136	152	172	190	199	230	219	239	257	231
Alcohol-related disorders	122	126	146	145	159	166	191	174	199	196	194
Anxiety disorders	671	765	841	901	979	1,022	1,091	1,044	1,143	1,147	1,109
Attention-deficit, conduct, and disruptive behavior disorders	62	78	88	108	125	138	160	159	197	230	233
Delirium, dementia, and amnestic and other cognitive disorders	323	336	342	340	334	321	332	290	288	298	263
Developmental disorders	76	78	87	92	91	103	118	121	140	140	156
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	32	39
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	34	33	42	44	44
Miscellaneous disorders	302	364	400	447	492	539	585	532	580	587	554
Mood disorders	679	771	864	966	1,002	1,055	1,133	1,062	1,175	1,194	1,165
Personality disorders	115	127	145	151	157	159	176	171	172	168	169
Schizophrenia and other psychotic disorders	243	268	287	289	286	287	297	279	271	274	249
Screening and history of mental health and substance abuse	712	827	908	1,019	1,100	1,173	1,267	1,164	1,368	1,396	1,360
Substance-related disorders	208	246	275	306	337	375	401	383	454	464	476
Suicide and intentional self-inflicted injury	54	62	71	77	88	93	106	102	111	120	115



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Shelby County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	936	1,085	1,228	1,383	1,476	1,592	1,645	1,627	1,752	1,788	1,672
Female	454	536	592	676	699	748	754	729	777	795	745
Male	482	549	636	707	777	844	891	898	975	993	927
White	704	811	920	1,063	1,125	1,238	1,276	1,255	1,345	1,381	1,298
Minority	176	213	237	242	264	263	264	268	295	293	267

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	44%	47%	51%	53%	52%	51%	49%	45%	46%	43%	39%
Female	43%	47%	50%	52%	51%	48%	45%	41%	42%	39%	35%
Male	44%	47%	52%	53%	54%	53%	52%	49%	50%	47%	42%
White	48%	50%	51%	53%	51%	50%	47%	44%	46%	42%	38%
Minority	25%	31%	39%	39%	41%	39%	40%	36%	34%	32%	29%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	260	302	332	355	355	378	367	355	365	374	333
Alcohol-related disorders	34	34	33	37	36	37	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	302	351	389	448	481	515	538	522	537	540	498
Attention-deficit, conduct, and disruptive behavior disorders	463	517	583	657	693	742	763	726	746	745	696
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	233	288	360	418	481	535	588	604	656	682	657
Disorders usually diagnosed in infancy, childhood, adolescence	91	108	121	135	155	164	179	172	173	180	167
Impulse control disorders, NEC	83	96	106	113	118	114	114	103	97	86	76
Miscellaneous disorders	79	94	103	114	123	127	133	139	143	135	129
Mood disorders	325	350	373	406	393	397	382	344	334	323	288
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	43	45	42	46	48	51	52	46	51	50	39
Screening and history of mental health and substance abuse	584	635	692	736	743	755	729	667	691	679	600
Substance-related disorders	69	71	77	85	90	91	86	70	69	70	55
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	36	37



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Shelby County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	836	902	991	1,057	1,170	1,270	1,333	1,193	1,442	1,462	1,346
Female	602	640	718	781	873	974	1,022	893	1,110	1,104	984
Male	234	262	273	276	297	296	311	300	332	358	362
White	668	709	779	819	879	964	1,045	967	1,117	1,158	1,078
Minority	135	158	172	196	249	259	239	178	270	249	213

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	54%	57%	62%	64%	65%	64%	58%	65%	64%	56%
Female	57%	54%	57%	63%	65%	67%	65%	57%	67%	65%	56%
Male	52%	54%	58%	62%	63%	58%	61%	59%	61%	59%	56%
White	59%	60%	62%	66%	66%	66%	65%	61%	67%	64%	56%
Minority	36%	32%	36%	43%	50%	51%	50%	37%	52%	50%	45%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	69	80	89	93	95	115	124	123	147	164	169
Alcohol-related disorders	50	58	64	71	77	96	108	103	107	110	108
Anxiety disorders	343	384	415	440	478	545	580	572	655	679	660
Attention-deficit, conduct, and disruptive behavior disorders	47	50	60	51	65	77	96	107	132	137	157
Delirium, dementia, and amnestic and other cognitive disorders	228	223	223	208	201	213	204	193	203	200	188
Developmental disorders	108	116	122	119	127	137	139	138	146	145	148
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	31	36
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	35	45	48	59	59	62
Miscellaneous disorders	125	133	145	156	173	194	201	194	217	222	203
Mood disorders	429	478	518	550	588	657	709	690	774	792	784
Personality disorders	53	54	61	58	63	68	70	71	74	75	74
Schizophrenia and other psychotic disorders	230	244	242	239	248	256	261	262	269	256	261
Screening and history of mental health and substance abuse	427	503	581	648	746	813	876	763	969	935	839
Substance-related disorders	63	77	90	91	95	120	131	122	153	164	169
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33	37	43	49	52



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Simpson County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	602	695	774	821	853	869	871	850	910	962	914
Female	310	353	385	409	406	415	415	385	412	437	410
Male	292	342	389	412	447	454	456	465	498	525	504
White	485	568	647	672	715	734	731	708	753	781	743
Minority	93	102	105	119	103	95	88	85	93	112	109

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	46%	47%	51%	49%	47%	45%	46%	44%	46%	44%	39%
Female	46%	47%	50%	48%	44%	43%	44%	41%	43%	41%	37%
Male	46%	48%	51%	49%	50%	48%	49%	47%	49%	47%	41%
White	47%	48%	52%	49%	48%	46%	46%	44%	46%	44%	39%
Minority	35%	37%	39%	37%	33%	29%	28%	28%	28%	27%	25%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	99	123	139	149	153	144	137	122	134	135	125
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	156	183	211	204	235	255	275	266	284	286	266
Attention-deficit, conduct, and disruptive behavior disorders	259	304	335	347	381	386	370	350	367	375	344
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	114	129	155	178	203	228	247	254	303	334	332
Disorders usually diagnosed in infancy, childhood, adolescence	57	63	81	87	90	98	100	93	95	93	94
Impulse control disorders, NEC	32	39	44	42	43	37	33	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	112	126	127	137	140	150	144	136	128	129	107
Mood disorders	188	207	220	223	228	200	198	175	163	160	133
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	379	448	495	506	476	425	402	379	375	366	339
Substance-related disorders	53	59	63	62	64	59	55	48	47	35	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Simpson County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	665	762	829	833	863	887	905	821	876	877	879
Female	503	568	609	619	643	666	676	605	651	656	631
Male	162	194	220	214	220	221	229	216	225	221	248
White	503	591	649	651	679	699	713	639	690	693	695
Minority	134	144	147	147	148	151	149	138	146	140	137

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	66%	68%	67%	65%	67%	68%	62%	66%	64%	59%
Female	64%	68%	70%	68%	67%	68%	68%	62%	67%	67%	59%
Male	51%	60%	64%	61%	62%	65%	67%	62%	61%	56%	58%
White	59%	65%	68%	67%	66%	68%	68%	62%	66%	66%	59%
Minority	54%	57%	56%	53%	50%	50%	51%	47%	50%	45%	43%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	56	58	70	74	77	87	94	91	95	94	89
Alcohol-related disorders	53	60	55	62	73	74	80	79	75	79	79
Anxiety disorders	317	378	408	423	452	468	484	473	484	507	485
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	35	36	40	44	51	56	49	68	70	83
Delirium, dementia, and amnesic and other cognitive disorders	192	206	208	207	201	200	188	184	187	174	162
Developmental disorders	53	57	56	62	60	66	70	72	74	81	76
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	227	264	298	307	340	366	407	374	385	362	357
Mood disorders	355	416	458	472	502	517	524	498	529	537	526
Personality disorders	N.D.	N.D.	N.D.	N.D.	31	32	36	34	34	39	37
Schizophrenia and other psychotic disorders	141	150	163	164	163	173	172	173	169	161	157
Screening and history of mental health and substance abuse	327	394	452	467	488	499	506	454	495	492	499
Substance-related disorders	82	98	102	111	113	113	121	122	130	133	125
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	35	41	44	54	52	53	54	53



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Spencer County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	367	423	452	500	541	562	585	555	598	617	584
Female	188	216	231	246	258	256	278	267	278	278	273
Male	179	207	221	254	283	306	307	288	320	339	311
White	N.D.	N.D.	419	460	492	513	531	504	544	562	524
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	46%	48%	49%	52%	54%	52%	52%	48%	50%	45%	41%
Female	47%	49%	51%	53%	55%	51%	52%	48%	48%	43%	40%
Male	46%	47%	46%	50%	53%	53%	51%	48%	51%	48%	42%
White	46%	48%	48%	51%	53%	52%	50%	47%	48%	44%	40%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	91	108	116	130	146	146	158	155	155	137	123
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	129	143	156	171	185	184	187	172	196	194	172
Attention-deficit, conduct, and disruptive behavior disorders	163	192	209	246	270	270	278	257	283	285	263
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	67	79	95	108	119	132	140	142	156	166	157
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	35	42	53	58	66	69	76	80	77
Impulse control disorders, NEC	N.D.	N.D.	32	31	34	N.D.	32	N.D.	31	N.D.	N.D.
Miscellaneous disorders	42	43	49	56	60	65	69	61	66	67	65
Mood disorders	153	163	168	180	201	195	192	170	179	165	145
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	195	234	240	254	271	261	257	239	251	231	215
Substance-related disorders	N.D.	N.D.	N.D.	34	35	34	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Spencer County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	414	442	457	464	503	519	553	522	547	552	522
Female	295	318	327	333	359	370	385	365	381	388	355
Male	119	124	130	131	144	149	168	157	166	164	167
White	355	382	402	407	444	462	488	468	497	508	475
Minority	49	47	43	46	45	44	46	35	31	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	59%	62%	64%	66%	67%	66%	67%	61%	65%	66%	58%
Female	60%	64%	66%	67%	68%	68%	68%	63%	67%	68%	59%
Male	56%	58%	61%	64%	65%	61%	63%	59%	60%	61%	56%
White	58%	62%	65%	66%	66%	66%	65%	60%	64%	65%	57%
Minority	52%	51%	49%	53%	56%	52%	55%	47%	46%	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	44	46	56	51	58	68	82	69	64	66	62
Alcohol-related disorders	45	48	50	52	55	55	64	53	59	56	58
Anxiety disorders	191	219	238	244	265	268	297	281	296	301	279
Attention-deficit, conduct, and disruptive behavior disorders	32	35	35	40	44	47	60	56	60	65	68
Delirium, dementia, and amnesic and other cognitive disorders	162	156	141	136	138	130	133	132	134	125	110
Developmental disorders	39	41	42	42	44	43	35	32	33	41	41
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	42	44	50	53	63	67	70	65	73	76	61
Mood disorders	253	290	310	308	333	343	363	337	345	340	318
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	33	37	31	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	144	150	147	151	152	141	149	135	129	121	110
Screening and history of mental health and substance abuse	197	233	257	266	305	323	340	314	340	340	307
Substance-related disorders	45	51	58	62	65	68	80	70	77	88	84
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Taylor County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,391	1,410	1,499	1,577	1,576	1,605	1,623	1,498	1,615	1,638	1,525
Female	709	696	741	761	753	750	750	659	731	741	677
Male	682	714	758	816	823	855	873	839	884	897	848
White	1,180	1,186	1,260	1,319	1,309	1,342	1,354	1,245	1,358	1,386	1,289
Minority	129	131	131	133	130	120	114	108	113	107	97

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	58%	61%	61%	61%	60%	61%	55%	56%	52%	47%
Female	60%	59%	62%	60%	58%	58%	58%	50%	53%	48%	43%
Male	56%	57%	61%	63%	63%	62%	64%	60%	59%	56%	50%
White	57%	56%	60%	60%	59%	59%	59%	54%	55%	50%	45%
Minority	39%	40%	39%	37%	33%	31%	30%	28%	28%	26%	23%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	323	323	369	383	380	369	369	325	335	336	297
Alcohol-related disorders	58	63	62	61	48	45	44	38	35	N.D.	N.D.
Anxiety disorders	634	654	695	712	709	717	719	660	678	655	591
Attention-deficit, conduct, and disruptive behavior disorders	665	701	775	817	806	824	837	746	772	776	724
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	326	360	412	467	473	504	530	503	560	575	552
Disorders usually diagnosed in infancy, childhood, adolescence	133	154	177	194	202	199	207	193	191	191	180
Impulse control disorders, NEC	139	151	166	166	165	163	161	134	129	124	111
Miscellaneous disorders	229	223	229	221	213	205	190	165	173	165	145
Mood disorders	599	579	592	585	555	540	518	427	422	411	354
Personality disorders	72	66	63	59	53	49	44	39	34	33	N.D.
Schizophrenia and other psychotic disorders	52	51	48	45	43	41	39	32	31	N.D.	N.D.
Screening and history of mental health and substance abuse	829	832	817	820	787	759	750	643	672	651	562
Substance-related disorders	153	137	135	131	115	108	96	74	72	67	53
Suicide and intentional self-inflicted injury	48	45	46	44	39	38	36	31	35	33	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Taylor County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,736	1,849	1,947	2,000	2,032	2,063	2,133	2,010	2,182	2,275	2,217
Female	1,193	1,256	1,313	1,351	1,383	1,411	1,471	1,391	1,504	1,548	1,483
Male	543	593	634	649	649	652	662	619	678	727	734
White	1,475	1,561	1,654	1,705	1,743	1,785	1,851	1,733	1,887	1,971	1,904
Minority	166	176	180	172	161	143	137	138	151	154	160

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	69%	72%	73%	74%	72%	73%	75%	71%	73%	72%	68%
Female	72%	75%	75%	76%	73%	74%	76%	72%	75%	74%	70%
Male	64%	68%	69%	69%	70%	72%	71%	67%	70%	69%	66%
White	70%	73%	74%	74%	73%	73%	75%	70%	73%	72%	68%
Minority	43%	43%	44%	43%	39%	37%	36%	37%	40%	36%	36%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	183	207	218	240	243	253	263	262	290	309	297
Alcohol-related disorders	228	257	266	268	284	274	282	267	284	288	272
Anxiety disorders	1,094	1,174	1,254	1,317	1,384	1,423	1,482	1,424	1,509	1,552	1,537
Attention-deficit, conduct, and disruptive behavior disorders	73	84	92	110	138	140	173	186	223	254	239
Delirium, dementia, and amnestic and other cognitive disorders	439	455	448	434	398	392	366	337	329	312	285
Developmental disorders	168	180	188	190	197	202	205	201	214	227	230
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	34	N.D.	N.D.	34	36	36	42	47	43
Impulse control disorders, NEC	44	50	55	55	60	65	76	73	80	87	93
Miscellaneous disorders	413	439	474	517	528	551	575	533	565	582	540
Mood disorders	1,125	1,220	1,283	1,357	1,395	1,415	1,477	1,417	1,504	1,557	1,497
Personality disorders	286	304	320	340	363	374	388	360	375	386	356
Schizophrenia and other psychotic disorders	455	480	503	488	484	485	504	483	478	466	432
Screening and history of mental health and substance abuse	1,033	1,117	1,205	1,266	1,333	1,383	1,447	1,374	1,523	1,589	1,556
Substance-related disorders	287	327	346	377	407	423	442	431	481	517	487
Suicide and intentional self-inflicted injury	68	75	83	87	104	110	126	119	133	132	119



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Todd County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	582	695	690	709	700	753	786	738	781	788	726
Female	294	353	353	360	340	362	377	341	357	359	330
Male	288	342	337	349	360	391	409	397	424	429	396
White	477	578	574	596	580	625	647	620	656	666	613
Minority	90	97	94	86	89	93	102	90	94	89	82

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	48%	49%	49%	48%	48%	49%	49%	45%	47%	45%	41%
Female	49%	50%	50%	49%	47%	48%	49%	43%	45%	42%	38%
Male	47%	48%	48%	48%	49%	50%	50%	47%	49%	48%	43%
White	49%	50%	50%	49%	48%	49%	50%	45%	47%	45%	41%
Minority	39%	37%	37%	34%	33%	33%	34%	34%	35%	31%	29%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	148	160	172	177	172	187	203	193	205	205	180
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	120	138	148	152	149	159	165	148	147	143	124
Attention-deficit, conduct, and disruptive behavior disorders	252	296	298	322	325	343	352	330	348	350	324
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	104	130	145	181	208	235	257	257	274	274	266
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	31	32	40	40	52	58	56	57	57	55
Impulse control disorders, NEC	54	61	61	53	47	52	55	43	50	45	32
Miscellaneous disorders	94	106	107	111	103	104	105	91	94	86	80
Mood disorders	189	206	210	215	194	195	188	154	152	144	122
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	307	344	342	299	270	271	277	233	238	234	199
Substance-related disorders	41	42	39	34	31	35	N.D.	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Todd County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	566	628	663	674	681	706	726	643	694	740	709
Female	410	452	464	474	485	501	522	464	509	533	492
Male	156	176	199	200	196	205	204	179	185	207	217
White	453	501	530	538	546	568	596	524	583	623	604
Minority	89	100	106	107	104	104	93	83	76	85	75

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	60%	62%	60%	60%	62%	65%	57%	60%	61%	58%
Female	59%	62%	62%	60%	60%	62%	66%	58%	62%	64%	60%
Male	53%	55%	60%	62%	60%	62%	62%	55%	56%	54%	56%
White	59%	61%	62%	61%	61%	63%	66%	58%	62%	62%	61%
Minority	40%	44%	46%	44%	42%	44%	41%	36%	36%	39%	34%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	52	55	64	68	70	77	79	82	87	96	100
Alcohol-related disorders	32	36	38	44	45	43	49	44	48	52	53
Anxiety disorders	262	290	313	329	334	344	372	345	386	398	404
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	33	40	44	49	64	66	66	73	84	85
Delirium, dementia, and amnestic and other cognitive disorders	169	176	181	179	167	171	163	142	137	125	116
Developmental disorders	44	48	49	46	47	48	51	45	49	53	54
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	69	88	93	105	110	116	128	109	120	132	116
Mood disorders	302	335	358	370	379	399	416	387	426	458	443
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	125	130	140	139	131	139	148	138	143	149	147
Screening and history of mental health and substance abuse	265	305	334	349	361	386	412	364	390	433	417
Substance-related disorders	33	37	43	52	57	61	75	61	73	89	87
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	33	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Trigg County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	491	531	560	595	610	652	674	598	643	659	608
Female	220	233	252	253	271	281	294	266	285	277	247
Male	271	298	308	342	339	371	380	332	358	382	361
White	399	432	448	468	492	524	549	493	522	527	482
Minority	67	73	81	90	81	84	78	64	72	80	76

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	54%	57%	58%	54%	54%	54%	48%	50%	46%	41%
Female	52%	52%	55%	54%	52%	49%	51%	46%	46%	40%	34%
Male	53%	55%	58%	62%	56%	58%	57%	51%	53%	52%	47%
White	54%	55%	57%	57%	53%	54%	55%	49%	49%	46%	40%
Minority	35%	37%	40%	44%	38%	37%	33%	28%	31%	29%	28%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	100	114	116	122	125	122	137	117	128	120	104
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	127	129	129	135	136	134	127	111	120	108	94
Attention-deficit, conduct, and disruptive behavior disorders	268	313	308	347	358	377	383	339	365	358	337
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	77	85	95	112	112	122	139	136	146	164	151
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	31	33	35	45	44	35	42	48	49
Impulse control disorders, NEC	57	64	66	69	62	62	61	50	53	47	42
Miscellaneous disorders	79	90	89	85	89	90	93	92	92	91	81
Mood disorders	167	175	176	178	182	170	162	144	137	126	110
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	252	277	288	302	295	298	305	269	269	275	242
Substance-related disorders	34	40	38	42	43	40	35	N.D.	N.D.	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Trigg County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	465	511	524	552	589	608	657	596	675	725	663
Female	333	364	369	389	414	426	458	420	464	510	458
Male	132	147	155	163	175	182	199	176	211	215	205
White	373	405	421	452	482	502	537	484	549	600	539
Minority	75	88	85	83	88	84	96	86	96	90	87

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	61%	63%	64%	63%	64%	66%	60%	66%	65%	58%
Female	60%	64%	65%	65%	63%	65%	68%	63%	67%	67%	60%
Male	54%	56%	58%	61%	61%	61%	61%	54%	64%	61%	55%
White	62%	64%	66%	66%	66%	66%	67%	61%	67%	67%	59%
Minority	39%	44%	44%	44%	43%	44%	48%	42%	46%	41%	38%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	N.D.	N.D.	37	39	45	58	65	66	67	80	63
Alcohol-related disorders	N.D.	32	31	33	42	39	42	43	51	49	45
Anxiety disorders	229	248	260	286	310	337	368	336	361	389	351
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	35	41	51	55	68	74	85	93	85
Delirium, dementia, and amnesic and other cognitive disorders	137	138	143	135	130	125	126	111	122	120	100
Developmental disorders	44	45	44	47	48	52	56	52	62	60	57
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	63	81	82	93	106	103	120	110	122	129	119
Mood disorders	254	280	296	340	376	392	432	385	438	477	427
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	93	101	109	105	119	121	126	129	134	134	120
Screening and history of mental health and substance abuse	191	224	242	273	306	313	346	315	347	373	345
Substance-related disorders	34	43	42	55	62	66	77	71	76	79	79
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	40	39



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Trimble County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	375	416	445	455	486	506	503	487	506	473	461
Female	191	203	211	216	233	248	232	222	226	201	196
Male	184	213	234	239	253	258	271	265	280	272	265
White	N.D.	384	N.D.	422	444	465	460	446	460	422	409
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	48%	49%	54%	51%	51%	52%	50%	47%	47%	43%	39%
Female	51%	50%	53%	49%	49%	51%	47%	45%	44%	40%	36%
Male	45%	48%	54%	52%	53%	54%	54%	49%	49%	45%	43%
White	47%	48%	53%	50%	49%	51%	49%	45%	45%	40%	37%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	111	119	118	122	131	136	135	129	137	122	122
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	121	143	142	149	155	157	160	149	145	135	134
Attention-deficit, conduct, and disruptive behavior disorders	174	201	221	227	237	255	262	256	269	251	234
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	72	87	99	117	132	145	146	151	167	156	156
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	33	33	38	41	45	47	50	51	46	43
Impulse control disorders, NEC	36	36	36	37	37	38	35	33	33	N.D.	N.D.
Miscellaneous disorders	34	36	36	39	37	42	41	N.D.	N.D.	N.D.	N.D.
Mood disorders	163	165	174	173	172	183	170	150	145	134	121
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	210	224	250	229	233	227	197	177	168	149	138
Substance-related disorders	34	33	39	35	34	37	35	31	32	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Trimble County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	330	329	395	413	431	429	445	427	512	515	453
Female	233	229	288	311	328	315	327	307	369	351	300
Male	97	100	107	102	103	114	118	120	143	164	153
White	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	484	421
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	57%	59%	66%	65%	64%	63%	65%	60%	66%	66%	59%
Female	58%	61%	69%	70%	68%	65%	68%	63%	70%	66%	60%
Male	54%	55%	59%	54%	54%	59%	57%	53%	59%	64%	58%
White	57%	59%	66%	65%	64%	63%	64%	59%	66%	64%	58%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	31	N.D.	38	38	38	46	54	51	64	69	65
Alcohol-related disorders	N.D.	N.D.	36	39	38	38	38	39	43	47	45
Anxiety disorders	147	158	197	215	233	242	259	259	308	309	273
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	31	35	37	45	49	45	57	68	55
Delirium, dementia, and amnestic and other cognitive disorders	106	109	110	102	99	98	92	88	82	82	67
Developmental disorders	31	31	35	34	34	37	33	34	37	38	34
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	60	59	69	82	95	95	105	101	115	100	90
Mood disorders	184	191	233	246	252	264	296	282	324	315	274
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	80	85	92	86	85	85	88	88	84	86	69
Screening and history of mental health and substance abuse	167	169	208	235	248	253	272	252	321	328	293
Substance-related disorders	N.D.	32	36	39	38	38	47	44	49	58	48
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Union County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	749	825	897	920	966	965	997	894	927	957	919
Female	351	383	415	424	437	439	454	416	423	444	428
Male	398	442	482	496	529	526	543	478	504	513	491
White	582	653	704	718	742	739	766	686	715	731	718
Minority	117	114	125	131	140	139	131	116	116	128	108

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	49%	61%	59%	59%	59%	61%	55%	54%	52%	49%
Female	51%	51%	60%	56%	56%	57%	60%	55%	52%	50%	47%
Male	53%	47%	63%	62%	62%	61%	63%	55%	56%	54%	51%
White	54%	54%	60%	58%	59%	58%	59%	53%	52%	49%	48%
Minority	33%	23%	42%	41%	38%	38%	39%	35%	34%	36%	31%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	125	134	138	133	144	146	145	122	121	123	120
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	195	206	224	228	229	226	227	196	199	195	194
Attention-deficit, conduct, and disruptive behavior disorders	427	468	523	542	570	573	586	529	539	539	505
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	169	204	246	272	306	323	347	330	347	362	356
Disorders usually diagnosed in infancy, childhood, adolescence	39	41	50	51	59	63	72	66	74	77	75
Impulse control disorders, NEC	60	66	70	68	68	72	75	62	65	62	55
Miscellaneous disorders	91	98	106	104	107	114	114	103	109	114	106
Mood disorders	357	373	385	379	394	397	377	327	318	318	295
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	400	437	456	429	412	411	412	346	347	343	321
Substance-related disorders	73	78	74	74	68	71	67	51	48	40	34
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Union County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	562	616	671	686	727	731	789	748	802	819	827
Female	444	488	518	542	566	573	607	568	610	609	598
Male	118	128	153	144	161	158	182	180	192	210	229
White	461	510	562	566	600	606	658	624	673	691	698
Minority	74	75	79	88	90	85	87	81	86	81	80

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	58%	54%	66%	66%	67%	65%	69%	64%	67%	65%	63%
Female	60%	61%	66%	68%	69%	69%	71%	65%	70%	68%	65%
Male	52%	38%	63%	59%	60%	55%	61%	60%	59%	59%	59%
White	59%	60%	65%	66%	67%	65%	69%	63%	66%	65%	64%
Minority	39%	27%	50%	49%	48%	45%	44%	43%	48%	42%	37%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	57	56	65	65	67	62	71	68	77	76	78
Alcohol-related disorders	43	39	41	46	50	56	60	58	58	69	68
Anxiety disorders	298	326	367	383	420	438	473	442	472	478	473
Attention-deficit, conduct, and disruptive behavior disorders	51	54	62	70	82	76	96	108	114	131	152
Delirium, dementia, and amnestic and other cognitive disorders	120	121	126	121	120	111	118	114	110	104	95
Developmental disorders	49	49	53	50	53	53	62	61	61	66	68
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	117	137	145	151	158	171	179	164	186	188	177
Mood disorders	359	402	442	472	499	508	555	529	574	591	594
Personality disorders	31	35	39	42	46	44	52	47	44	44	46
Schizophrenia and other psychotic disorders	106	109	120	108	116	115	117	110	111	114	102
Screening and history of mental health and substance abuse	278	311	333	360	368	357	402	376	404	415	406
Substance-related disorders	112	112	130	139	150	158	179	177	174	177	177
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	32	42	44	47	43	45



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Warren County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,465	5,005	5,297	5,467	5,562	5,621	5,692	5,414	5,775	5,947	5,645
Female	2,144	2,386	2,530	2,600	2,640	2,632	2,650	2,428	2,613	2,720	2,564
Male	2,321	2,619	2,767	2,867	2,922	2,989	3,042	2,986	3,162	3,227	3,081
White	3,287	3,694	3,870	3,976	4,000	4,041	4,080	3,887	4,129	4,282	4,087
Minority	973	1,071	1,145	1,184	1,223	1,234	1,219	1,144	1,235	1,245	1,170

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	54%	55%	55%	54%	53%	53%	49%	50%	48%	42%
Female	51%	53%	54%	54%	52%	51%	50%	46%	47%	45%	40%
Male	52%	54%	56%	56%	56%	55%	55%	52%	53%	50%	45%
White	52%	54%	55%	55%	55%	53%	52%	49%	50%	49%	44%
Minority	42%	43%	43%	43%	42%	41%	41%	38%	38%	33%	29%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	774	849	929	955	952	936	894	814	809	797	728
Alcohol-related disorders	158	160	159	156	146	143	141	107	98	83	61
Anxiety disorders	1,173	1,343	1,458	1,506	1,615	1,680	1,731	1,689	1,778	1,803	1,699
Attention-deficit, conduct, and disruptive behavior disorders	2,193	2,440	2,636	2,758	2,767	2,791	2,826	2,632	2,659	2,621	2,421
Delirium, dementia, and amnestic and other cognitive disorders	64	63	65	66	64	62	63	57	59	60	57
Developmental disorders	1,097	1,308	1,451	1,578	1,711	1,769	1,867	1,885	1,998	2,063	2,022
Disorders usually diagnosed in infancy, childhood, adolescence	393	457	518	566	591	627	655	643	677	683	656
Impulse control disorders, NEC	398	437	451	460	452	432	392	352	332	314	267
Miscellaneous disorders	642	694	728	741	729	731	698	635	633	605	557
Mood disorders	1,613	1,700	1,739	1,703	1,642	1,537	1,467	1,307	1,249	1,163	1,034
Personality disorders	93	101	95	95	91	85	68	53	56	51	39
Schizophrenia and other psychotic disorders	185	199	202	193	201	189	176	164	155	147	123
Screening and history of mental health and substance abuse	3,315	3,553	3,637	3,547	3,412	3,252	3,142	2,813	2,949	2,918	2,708
Substance-related disorders	394	415	409	406	356	335	322	264	246	222	168
Suicide and intentional self-inflicted injury	99	100	100	103	94	84	81	64	62	57	46



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Warren County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,621	5,191	5,555	5,626	5,685	5,754	5,951	5,419	5,851	6,079	5,941
Female	3,281	3,660	3,924	3,997	4,050	4,112	4,213	3,803	4,148	4,252	4,106
Male	1,340	1,531	1,631	1,629	1,635	1,642	1,738	1,616	1,703	1,827	1,835
White	3,656	4,107	4,391	4,443	4,475	4,537	4,686	4,261	4,592	4,773	4,648
Minority	757	845	905	918	926	918	938	834	904	937	917

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	64%	67%	69%	70%	70%	69%	70%	64%	68%	65%	60%
Female	65%	68%	70%	71%	71%	69%	70%	63%	68%	66%	61%
Male	62%	66%	68%	69%	68%	68%	69%	65%	67%	64%	59%
White	65%	69%	71%	72%	72%	71%	72%	66%	70%	69%	63%
Minority	47%	48%	49%	50%	49%	47%	47%	42%	43%	40%	37%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	438	476	512	524	537	545	565	522	569	602	592
Alcohol-related disorders	532	604	650	651	678	675	692	629	663	680	657
Anxiety disorders	2,431	2,693	2,944	3,067	3,154	3,231	3,365	3,195	3,412	3,519	3,426
Attention-deficit, conduct, and disruptive behavior disorders	361	424	460	522	599	636	710	705	831	899	910
Delirium, dementia, and amnestic and other cognitive disorders	1,082	1,125	1,145	1,103	1,054	1,042	1,044	964	943	895	831
Developmental disorders	480	515	532	542	547	545	611	592	611	651	636
Disorders usually diagnosed in infancy, childhood, adolescence	60	69	77	82	89	95	119	114	134	152	155
Impulse control disorders, NEC	119	124	139	142	161	160	186	184	210	220	229
Miscellaneous disorders	979	1,129	1,209	1,299	1,327	1,404	1,504	1,381	1,436	1,511	1,417
Mood disorders	2,973	3,346	3,636	3,748	3,865	3,912	4,060	3,762	4,032	4,158	4,044
Personality disorders	328	347	364	366	366	354	373	356	373	361	358
Schizophrenia and other psychotic disorders	1,089	1,166	1,224	1,237	1,244	1,246	1,283	1,233	1,265	1,215	1,142
Screening and history of mental health and substance abuse	2,940	3,373	3,717	3,804	3,880	3,877	4,019	3,655	3,978	4,112	4,010
Substance-related disorders	692	780	832	858	921	953	1,026	962	1,038	1,053	1,026
Suicide and intentional self-inflicted injury	184	201	228	248	269	273	316	310	344	346	347



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Washington County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	449	507	519	560	577	581	585	564	591	616	573
Female	205	239	248	256	261	255	260	258	265	266	248
Male	244	268	271	304	316	326	325	306	326	350	325
White	351	399	412	443	450	462	470	455	481	504	470
Minority	71	74	71	74	78	66	64	56	56	51	46

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	48%	51%	51%	53%	53%	55%	55%	53%	53%	48%	45%
Female	46%	50%	50%	50%	50%	49%	51%	50%	48%	42%	40%
Male	49%	53%	53%	55%	57%	60%	59%	56%	58%	54%	50%
White	49%	51%	51%	53%	52%	54%	54%	52%	51%	47%	44%
Minority	33%	36%	33%	33%	36%	31%	33%	31%	32%	25%	23%


CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	83	100	107	117	116	118	115	110	119	132	115
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	137	144	165	175	171	164	166	160	157	167	143
Attention-deficit, conduct, and disruptive behavior disorders	253	279	280	298	316	322	315	298	301	310	281
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	99	121	133	148	166	181	185	190	208	225	207
Disorders usually diagnosed in infancy, childhood, adolescence	51	62	68	80	88	90	86	86	84	104	103
Impulse control disorders, NEC	42	42	44	46	45	41	39	31	N.D.	31	N.D.
Miscellaneous disorders	72	81	89	96	107	99	107	103	104	102	91
Mood disorders	174	193	209	205	191	189	185	174	169	163	148
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	275	294	294	302	292	273	272	244	238	237	214
Substance-related disorders	53	51	56	51	49	45	41	41	31	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Washington County											
ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	520	570	557	600	631	640	657	643	701	720	700
Female	374	403	397	427	454	449	468	456	493	504	489
Male	146	167	160	173	177	191	189	187	208	216	211
White	429	468	452	493	521	533	543	544	595	613	600
Minority	74	83	85	85	88	81	84	72	74	75	68
ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	57%	57%	62%	65%	64%	64%	62%	66%	64%	62%
Female	55%	59%	61%	65%	67%	65%	65%	63%	66%	64%	62%
Male	48%	52%	51%	57%	59%	63%	63%	62%	64%	64%	62%
White	55%	59%	59%	64%	66%	67%	66%	64%	68%	66%	63%
Minority	37%	40%	42%	44%	47%	41%	43%	39%	39%	40%	37%
ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	47	52	47	48	51	55	59	57	65	71	78
Alcohol-related disorders	43	46	45	51	54	60	61	57	65	66	61
Anxiety disorders	252	287	301	323	354	370	383	380	419	428	409
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	31	35	44	43	49	52	48	62	66	75
Delirium, dementia, and amnestic and other cognitive disorders	182	185	173	170	167	177	177	179	177	166	152
Developmental disorders	50	54	59	66	67	70	75	78	78	76	80
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	101	102	111	117	131	135	148	149	155	159	155
Mood disorders	271	304	310	329	356	377	383	380	409	435	413
Personality disorders	49	54	56	61	70	71	76	70	69	67	62
Schizophrenia and other psychotic disorders	151	165	168	173	177	177	173	167	177	176	162
Screening and history of mental health and substance abuse	220	256	252	296	310	317	332	319	353	373	377
Substance-related disorders	45	52	55	69	72	83	83	74	91	97	97
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Wayne County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,654	1,791	1,896	2,005	2,012	2,003	2,036	1,924	1,993	1,960	1,855
Female	831	894	938	978	987	978	999	943	969	948	891
Male	823	897	958	1,027	1,025	1,025	1,037	981	1,024	1,012	964
White	1,544	1,667	1,752	1,844	1,839	1,827	1,847	1,742	1,795	1,766	1,666
Minority	39	45	48	53	51	50	51	48	52	47	43

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	55%	58%	60%	62%	62%	61%	63%	58%	60%	58%	54%
Female	56%	58%	61%	62%	62%	61%	62%	57%	58%	56%	52%
Male	54%	57%	60%	63%	62%	62%	64%	59%	61%	59%	55%
White	54%	57%	60%	61%	61%	60%	62%	57%	58%	57%	53%
Minority	26%	25%	24%	25%	22%	21%	23%	20%	21%	17%	15%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	286	307	307	322	323	321	317	274	272	258	232
Alcohol-related disorders	78	79	75	72	58	41	41	33	N.D.	N.D.	N.D.
Anxiety disorders	846	917	955	1,014	1,021	1,032	1,061	996	1,017	998	936
Attention-deficit, conduct, and disruptive behavior disorders	665	707	723	773	781	768	762	699	700	684	630
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	397	447	504	554	604	623	654	630	690	692	673
Disorders usually diagnosed in infancy, childhood, adolescence	87	100	110	124	126	124	130	125	123	116	115
Impulse control disorders, NEC	67	70	69	68	52	44	38	35	32	32	N.D.
Miscellaneous disorders	223	232	235	221	207	194	186	156	154	147	135
Mood disorders	619	627	614	604	558	521	489	429	407	385	336
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	56	56	54	58	51	48	43	42	41	41	36
Screening and history of mental health and substance abuse	826	856	894	893	861	814	778	713	720	672	635
Substance-related disorders	190	183	177	159	133	114	100	78	73	69	59
Suicide and intentional self-inflicted injury	57	55	50	49	39	37	31	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Wayne County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	2,207	2,304	2,331	2,450	2,493	2,531	2,636	2,443	2,547	2,530	2,394
Female	1,436	1,488	1,502	1,582	1,603	1,606	1,666	1,570	1,638	1,627	1,512
Male	771	816	829	868	890	925	970	873	909	903	882
White	2,002	2,104	2,126	2,245	2,290	2,324	2,415	2,228	2,339	2,318	2,186
Minority	76	70	76	70	64	60	56	52	57	54	54

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	66%	68%	69%	69%	69%	69%	71%	65%	67%	67%	63%
Female	69%	70%	71%	72%	72%	72%	73%	67%	71%	70%	66%
Male	61%	64%	65%	64%	64%	65%	67%	61%	63%	62%	60%
White	66%	68%	69%	69%	69%	69%	70%	64%	67%	67%	63%
Minority	26%	24%	26%	24%	22%	21%	19%	17%	19%	17%	17%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	221	241	260	282	278	277	303	283	299	293	268
Alcohol-related disorders	202	223	223	235	256	260	256	238	246	239	223
Anxiety disorders	1,360	1,448	1,516	1,614	1,663	1,695	1,793	1,676	1,741	1,755	1,705
Attention-deficit, conduct, and disruptive behavior disorders	71	72	86	91	102	125	149	142	174	187	192
Delirium, dementia, and amnestic and other cognitive disorders	422	420	410	389	370	357	343	310	280	256	241
Developmental disorders	204	201	212	221	227	238	242	220	230	234	239
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	38	38	51	63	67	79	79	68	75	71	68
Miscellaneous disorders	573	583	609	655	691	704	726	702	727	716	668
Mood disorders	1,265	1,319	1,393	1,509	1,538	1,577	1,660	1,558	1,610	1,598	1,548
Personality disorders	89	88	89	91	97	98	105	93	85	85	85
Schizophrenia and other psychotic disorders	448	449	455	453	437	417	422	394	383	357	340
Screening and history of mental health and substance abuse	1,116	1,210	1,248	1,318	1,343	1,343	1,416	1,308	1,386	1,390	1,307
Substance-related disorders	390	410	426	453	494	505	523	490	522	519	499
Suicide and intentional self-inflicted injury	49	64	69	79	84	87	97	94	97	97	94



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Webster County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	634	704	776	799	815	813	855	776	777	770	713
Female	308	339	384	383	383	372	396	349	344	339	315
Male	326	365	392	416	432	441	459	427	433	431	398
White	538	595	660	682	687	686	718	655	657	655	602
Minority	52	56	55	53	55	52	54	44	43	37	38

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	52%	53%	57%	55%	54%	54%	56%	50%	51%	48%	44%
Female	52%	52%	56%	53%	52%	50%	51%	45%	46%	44%	40%
Male	52%	54%	57%	56%	57%	58%	60%	55%	56%	53%	48%
White	52%	53%	57%	55%	54%	53%	54%	47%	49%	46%	42%
Minority	29%	27%	27%	25%	25%	24%	26%	25%	24%	21%	21%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	101	112	120	132	125	126	129	122	115	107	93
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	191	215	237	240	244	241	254	221	213	201	187
Attention-deficit, conduct, and disruptive behavior disorders	321	365	408	407	413	433	455	406	416	414	375
Delirium, dementia, and amnestic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	149	183	219	237	259	275	309	291	302	308	295
Disorders usually diagnosed in infancy, childhood, adolescence	41	44	58	65	76	83	88	84	87	94	83
Impulse control disorders, NEC	63	66	68	72	74	66	66	59	52	50	41
Miscellaneous disorders	88	88	101	102	103	103	113	100	100	93	90
Mood disorders	267	278	303	301	298	287	289	257	243	224	185
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	31	N.D.	31	33	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	352	367	387	377	366	347	343	297	296	272	234
Substance-related disorders	52	48	43	48	47	44	42	33	31	N.D.	N.D.
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Webster County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	634	698	743	788	852	849	848	775	851	831	798
Female	435	489	520	559	594	597	607	532	574	567	556
Male	199	209	223	229	258	252	241	243	277	264	242
White	567	626	669	706	760	760	756	691	767	749	713
Minority	45	47	50	51	53	48	41	36	37	36	40

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	62%	63%	64%	67%	67%	65%	65%	60%	66%	65%	62%
Female	63%	66%	67%	69%	69%	67%	68%	61%	67%	65%	64%
Male	61%	56%	60%	62%	64%	60%	58%	59%	63%	64%	58%
White	63%	64%	65%	68%	68%	65%	65%	60%	67%	65%	62%
Minority	39%	36%	38%	38%	34%	31%	29%	27%	26%	26%	30%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	40	46	52	58	62	62	62	50	56	63	60
Alcohol-related disorders	54	58	55	62	73	67	72	62	68	67	68
Anxiety disorders	345	389	432	461	494	506	548	491	529	520	498
Attention-deficit, conduct, and disruptive behavior disorders	53	61	71	74	87	88	107	104	116	130	120
Delirium, dementia, and amnestic and other cognitive disorders	146	139	137	135	123	120	120	118	118	117	109
Developmental disorders	62	62	65	68	73	74	76	73	72	73	70
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	122	141	145	158	178	168	173	160	165	176	165
Mood disorders	404	450	500	532	590	582	605	552	595	581	563
Personality disorders	47	51	57	55	68	66	63	62	65	69	62
Schizophrenia and other psychotic disorders	118	129	130	127	136	142	139	128	137	143	133
Screening and history of mental health and substance abuse	342	401	430	463	511	511	521	464	509	485	466
Substance-related disorders	65	82	91	104	115	113	118	115	130	142	128
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	31	42	42	46	41	47	52	54



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Whitley County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	3,798	4,154	4,377	4,663	4,791	4,846	4,860	4,594	4,747	4,590	4,292
Female	1,907	2,086	2,148	2,289	2,345	2,354	2,342	2,219	2,288	2,200	2,043
Male	1,891	2,068	2,229	2,374	2,446	2,492	2,518	2,375	2,459	2,390	2,249
White	3,497	3,807	3,998	4,257	4,366	4,396	4,382	4,152	4,284	4,132	3,856
Minority	N.D.	33	37	35	37	39	41	42	47	48	42

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	60%	63%	65%	67%	68%	69%	70%	66%	68%	64%	60%
Female	63%	65%	66%	68%	67%	68%	69%	66%	66%	63%	58%
Male	58%	60%	64%	67%	68%	70%	71%	67%	69%	66%	61%
White	59%	61%	64%	66%	66%	68%	68%	65%	66%	63%	59%
Minority	N.D.	8%	8%	7%	7%	8%	8%	8%	9%	8%	7%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	733	784	826	888	893	880	850	777	767	700	641
Alcohol-related disorders	373	407	409	418	408	392	380	326	305	270	238
Anxiety disorders	2,008	2,220	2,375	2,595	2,699	2,775	2,804	2,684	2,766	2,691	2,552
Attention-deficit, conduct, and disruptive behavior disorders	1,720	1,881	2,012	2,138	2,171	2,161	2,129	1,984	1,975	1,880	1,739
Delirium, dementia, and amnestic and other cognitive disorders	75	86	85	82	80	75	71	66	60	59	50
Developmental disorders	1,084	1,203	1,321	1,441	1,498	1,534	1,537	1,454	1,476	1,425	1,339
Disorders usually diagnosed in infancy, childhood, adolescence	384	440	476	518	555	568	565	531	538	511	488
Impulse control disorders, NEC	181	192	204	211	208	202	205	166	156	143	123
Miscellaneous disorders	750	771	777	770	771	730	712	675	665	601	542
Mood disorders	1,360	1,374	1,377	1,394	1,344	1,285	1,195	1,046	986	862	744
Personality disorders	319	326	331	347	345	332	319	272	242	229	205
Schizophrenia and other psychotic disorders	158	163	161	159	150	147	136	120	109	95	82
Screening and history of mental health and substance abuse	1,950	2,053	2,100	2,156	2,132	2,088	2,063	1,891	1,912	1,806	1,649
Substance-related disorders	719	741	746	745	700	652	595	510	465	426	355
Suicide and intentional self-inflicted injury	100	106	103	95	91	87	85	75	74	69	61



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Whitley County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	4,586	4,912	5,254	5,541	5,662	5,659	5,910	5,681	6,044	6,087	5,824
Female	3,129	3,330	3,508	3,653	3,730	3,713	3,844	3,729	3,958	3,960	3,766
Male	1,457	1,582	1,746	1,888	1,932	1,946	2,066	1,952	2,086	2,127	2,058
White	4,195	4,491	4,814	5,085	5,208	5,200	5,419	5,219	5,567	5,613	5,344
Minority	97	101	95	92	80	72	54	44	42	44	44

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	68%	71%	73%	74%	73%	73%	76%	73%	76%	74%	71%
Female	71%	75%	76%	77%	76%	76%	79%	76%	79%	78%	74%
Male	62%	64%	66%	68%	68%	69%	72%	67%	70%	69%	67%
White	68%	71%	72%	73%	73%	73%	75%	72%	75%	74%	71%
Minority	18%	18%	17%	16%	14%	13%	9%	8%	7%	7%	7%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	445	476	519	563	607	637	691	689	749	758	713
Alcohol-related disorders	540	569	625	678	707	708	760	709	737	739	742
Anxiety disorders	3,188	3,440	3,743	3,950	4,075	4,103	4,307	4,173	4,369	4,354	4,178
Attention-deficit, conduct, and disruptive behavior disorders	228	275	306	350	413	446	506	517	559	610	619
Delirium, dementia, and amnesic and other cognitive disorders	867	900	900	901	869	843	849	805	795	757	686
Developmental disorders	424	444	475	503	527	545	605	581	611	653	657
Disorders usually diagnosed in infancy, childhood, adolescence	66	67	68	76	81	87	94	103	114	118	121
Impulse control disorders, NEC	106	117	125	137	150	151	163	163	176	180	185
Miscellaneous disorders	1,524	1,660	1,781	1,922	1,991	2,028	2,127	2,059	2,133	2,153	2,045
Mood disorders	2,788	3,002	3,224	3,436	3,540	3,566	3,773	3,658	3,832	3,837	3,695
Personality disorders	496	538	559	618	654	673	724	694	714	716	682
Schizophrenia and other psychotic disorders	981	1,048	1,069	1,102	1,099	1,101	1,172	1,099	1,088	1,042	962
Screening and history of mental health and substance abuse	2,564	2,789	3,043	3,276	3,399	3,451	3,701	3,551	3,823	3,872	3,773
Substance-related disorders	1,172	1,277	1,401	1,533	1,609	1,654	1,777	1,708	1,849	1,873	1,833
Suicide and intentional self-inflicted injury	197	221	235	269	277	286	336	321	336	334	319



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Wolfe County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	924	966	946	989	983	995	1,028	971	1,020	1,004	944
Female	473	491	485	502	483	481	496	446	485	474	444
Male	451	475	461	487	500	514	532	525	535	530	500
White	835	873	850	886	877	884	902	853	892	877	828
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	66%	69%	68%	69%	67%	66%	68%	65%	68%	67%	62%
Female	69%	71%	70%	71%	68%	66%	67%	62%	66%	66%	60%
Male	63%	67%	65%	68%	66%	66%	70%	68%	70%	67%	63%
White	64%	67%	65%	67%	65%	64%	65%	62%	65%	64%	59%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	212	232	239	257	272	282	287	272	277	259	251
Alcohol-related disorders	93	92	85	87	80	73	71	62	58	51	43
Anxiety disorders	471	486	481	521	511	517	531	490	486	466	429
Attention-deficit, conduct, and disruptive behavior disorders	314	333	335	371	372	386	403	377	376	364	335
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	324	356	365	400	408	423	433	418	428	415	398
Disorders usually diagnosed in infancy, childhood, adolescence	87	89	100	105	110	111	110	100	107	108	95
Impulse control disorders, NEC	86	91	91	100	99	95	90	84	83	71	56
Miscellaneous disorders	111	105	108	108	98	95	94	81	80	78	73
Mood disorders	366	353	328	322	301	294	286	253	246	229	192
Personality disorders	54	52	49	51	46	43	42	41	38	39	N.D.
Schizophrenia and other psychotic disorders	46	47	44	38	38	39	40	35	33	32	N.D.
Screening and history of mental health and substance abuse	587	602	572	567	544	523	542	495	507	489	451
Substance-related disorders	143	140	126	115	104	100	96	92	91	81	64
Suicide and intentional self-inflicted injury	31	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Wolfe County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	1,366	1,405	1,508	1,592	1,637	1,628	1,689	1,650	1,738	1,753	1,685
Female	853	877	946	990	1,023	1,025	1,062	1,028	1,086	1,084	1,024
Male	513	528	562	602	614	603	627	622	652	669	661
White	1,253	1,290	1,393	1,476	1,516	1,511	1,565	1,526	1,612	1,628	1,560
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	73%	75%	77%	78%	78%	79%	81%	79%	81%	80%	75%
Female	77%	79%	82%	83%	83%	84%	86%	82%	86%	85%	80%
Male	67%	68%	71%	71%	71%	71%	74%	73%	74%	73%	68%
White	72%	74%	77%	78%	78%	78%	81%	78%	81%	80%	74%
Minority	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	131	143	148	166	176	185	198	193	199	212	205
Alcohol-related disorders	192	199	233	249	250	253	255	248	255	255	248
Anxiety disorders	943	993	1,079	1,155	1,196	1,204	1,272	1,246	1,296	1,323	1,274
Attention-deficit, conduct, and disruptive behavior disorders	33	38	41	47	58	60	68	67	93	103	106
Delirium, dementia, and amnestic and other cognitive disorders	275	272	272	259	258	236	233	228	226	222	200
Developmental disorders	154	164	167	182	191	190	213	207	223	223	214
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	32	33	33	41	47	43
Miscellaneous disorders	319	352	375	409	416	426	428	423	422	413	392
Mood disorders	847	899	979	1,041	1,084	1,099	1,157	1,139	1,193	1,205	1,176
Personality disorders	121	127	143	161	170	167	171	166	162	160	161
Schizophrenia and other psychotic disorders	231	244	253	264	275	261	270	268	269	276	267
Screening and history of mental health and substance abuse	834	882	974	1,037	1,073	1,091	1,142	1,129	1,195	1,193	1,149
Substance-related disorders	288	308	343	388	409	419	444	430	458	478	466
Suicide and intentional self-inflicted injury	38	41	52	55	58	61	63	65	69	73	75



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Woodford County

CHILDREN: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	560	637	730	744	772	799	824	756	809	828	796
Female	284	304	341	327	348	356	357	321	337	349	337
Male	276	333	389	417	424	443	467	435	472	479	459
White	482	550	630	645	669	689	699	638	686	705	675
Minority	58	61	74	71	75	77	79	71	73	73	72

CHILDREN: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	53%	51%	55%	54%	51%	49%	50%	46%	47%	43%	37%
Female	54%	50%	54%	51%	50%	47%	45%	40%	41%	37%	32%
Male	52%	52%	57%	57%	52%	52%	54%	51%	53%	48%	42%
White	55%	52%	57%	56%	53%	50%	49%	45%	46%	42%	37%
Minority	33%	32%	36%	32%	31%	31%	33%	32%	32%	27%	25%

CHILDREN: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	170	195	229	237	250	253	262	235	252	239	225
Alcohol-related disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Anxiety disorders	184	200	225	214	203	211	227	209	213	204	191
Attention-deficit, conduct, and disruptive behavior disorders	269	305	350	367	383	387	416	373	398	383	359
Delirium, dementia, and amnesic and other cognitive disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Developmental disorders	110	137	180	206	221	247	278	266	293	313	306
Disorders usually diagnosed in infancy, childhood, adolescence	52	58	76	81	84	87	101	89	100	101	103
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	95	104	127	113	111	111	108	95	99	94	83
Mood disorders	208	225	245	234	226	232	230	194	205	194	174
Personality disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Schizophrenia and other psychotic disorders	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Screening and history of mental health and substance abuse	354	378	417	397	396	388	369	335	322	301	263
Substance-related disorders	63	66	75	67	63	59	66	62	57	49	42
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."


A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

Woodford County

ADULTS: Number	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	416	504	562	624	638	642	663	630	719	768	762
Female	314	381	431	483	493	497	502	474	540	573	554
Male	102	123	131	141	145	145	161	156	179	195	208
White	341	412	470	519	537	537	556	530	615	652	638
Minority	57	71	72	76	68	72	71	68	72	84	88

ADULTS: Percent	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
All	59%	63%	64%	63%	64%	64%	66%	62%	67%	64%	61%
Female	59%	63%	65%	65%	66%	65%	67%	62%	68%	65%	62%
Male	56%	63%	60%	59%	61%	61%	65%	61%	62%	61%	59%
White	60%	65%	67%	65%	67%	66%	68%	63%	69%	67%	62%
Minority	41%	43%	39%	40%	36%	37%	39%	38%	40%	38%	39%

ADULTS: Number with Disorder	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Adjustment disorders	48	68	86	99	91	93	109	109	140	151	151
Alcohol-related disorders	43	44	59	67	69	69	80	77	79	80	85
Anxiety disorders	203	258	305	347	358	363	396	384	426	443	443
Attention-deficit, conduct, and disruptive behavior disorders	N.D.	N.D.	32	41	43	45	48	53	71	93	100
Delirium, dementia, and amnesic and other cognitive disorders	87	87	90	83	84	86	81	74	70	65	65
Developmental disorders	N.D.	N.D.	31	35	38	43	46	44	49	57	62
Disorders usually diagnosed in infancy, childhood, adolescence	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Impulse control disorders, NEC	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Miscellaneous disorders	102	133	160	200	198	211	217	180	216	222	227
Mood disorders	237	297	337	381	386	387	424	404	450	460	480
Personality disorders	37	44	54	56	49	57	61	54	63	63	62
Schizophrenia and other psychotic disorders	75	80	86	82	85	86	93	92	91	85	89
Screening and history of mental health and substance abuse	223	296	351	414	416	423	450	440	504	534	513
Substance-related disorders	79	94	111	139	133	147	161	158	190	202	201
Suicide and intentional self-inflicted injury	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	35	34



Note: N.D. indicates that the number is not disclosable because the sample size is too small while #N/A indicates the value is zero and therefore "not applicable."

A PROFILE OF KENTUCKY MEDICAID MENTAL HEALTH DIAGNOSES, 2000 TO 2010

